

Albena T Dinkova-Kostova

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7332352/albena-t-dinkova-kostova-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

159
papers

18,557
citations

67
h-index

136
g-index

171
ext. papers

21,837
ext. citations

6.9
avg, IF

7.29
L-index

#	Paper	IF	Citations
159	Direct evidence that sulfhydryl groups of Keap1 are the sensors regulating induction of phase 2 enzymes that protect against carcinogens and oxidants. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 11908-13	11.5	1519
158	The Nrf2 regulatory network provides an interface between redox and intermediary metabolism. <i>Trends in Biochemical Sciences</i> , 2014 , 39, 199-218	10.3	1157
157	Protection against electrophile and oxidant stress by induction of the phase 2 response: fate of cysteines of the Keap1 sensor modified by inducers. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 2040-5	11.5	805
156	The cytoprotective role of the Keap1-Nrf2 pathway. <i>Archives of Toxicology</i> , 2011 , 85, 241-72	5.8	687
155	Itaconate is an anti-inflammatory metabolite that activates Nrf2 via alkylation of KEAP1. <i>Nature</i> , 2018 , 556, 113-117	50.4	609
154	The emerging role of Nrf2 in mitochondrial function. <i>Free Radical Biology and Medicine</i> , 2015 , 88, 179-188	8.8	493
153	Mechanisms of activation of the transcription factor Nrf2 by redox stressors, nutrient cues, and energy status and the pathways through which it attenuates degenerative disease. <i>Free Radical Biology and Medicine</i> , 2015 , 88, 108-146	7.8	483
152	NAD(P)H:quinone acceptor oxidoreductase 1 (NQO1), a multifunctional antioxidant enzyme and exceptionally versatile cytoprotector. <i>Archives of Biochemistry and Biophysics</i> , 2010 , 501, 116-23	4.1	481
151	Therapeutic targeting of the NRF2 and KEAP1 partnership in chronic diseases. <i>Nature Reviews Drug Discovery</i> , 2019 , 18, 295-317	64.1	476
150	Extremely potent triterpenoid inducers of the phase 2 response: correlations of protection against oxidant and inflammatory stress. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005 , 102, 4584-9	11.5	445
149	Cellular stress responses, the hormesis paradigm, and vitagenes: novel targets for therapeutic intervention in neurodegenerative disorders. <i>Antioxidants and Redox Signaling</i> , 2010 , 13, 1763-811	8.4	434
148	Cancer chemoprevention mechanisms mediated through the Keap1-Nrf2 pathway. <i>Antioxidants and Redox Signaling</i> , 2010 , 13, 1713-48	8.4	413
147	Glucosinolates and isothiocyanates in health and disease. <i>Trends in Molecular Medicine</i> , 2012 , 18, 337-47	11.5	410
146	Oxidative Stress in Cancer. <i>Cancer Cell</i> , 2020 , 38, 167-197	24.3	402
145	Nrf2 regulates ROS production by mitochondria and NADPH oxidase. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2015 , 1850, 794-801	4	319
144	The role of Keap1 in cellular protective responses. <i>Chemical Research in Toxicology</i> , 2005 , 18, 1779-91	4	318
143	Quantitative determination of dithiocarbamates in human plasma, serum, erythrocytes and urine: pharmacokinetics of broccoli sprout isothiocyanates in humans. <i>Clinica Chimica Acta</i> , 2002 , 316, 43-53	6.2	296

142	Cellular stress responses, hormetic phytochemicals and vitagenes in aging and longevity. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2012 , 1822, 753-83	6.9	286
141	Nrf2 impacts cellular bioenergetics by controlling substrate availability for mitochondrial respiration. <i>Biology Open</i> , 2013 , 2, 761-70	2.2	266
140	Safety, tolerance, and metabolism of broccoli sprout glucosinolates and isothiocyanates: a clinical phase I study. <i>Nutrition and Cancer</i> , 2006 , 55, 53-62	2.8	265
139	Preclinical and clinical evaluation of sulforaphane for chemoprevention in the breast. <i>Carcinogenesis</i> , 2007 , 28, 1485-90	4.6	256
138	Importance of phase 2 gene regulation in protection against electrophile and reactive oxygen toxicity and carcinogenesis. <i>Advances in Enzyme Regulation</i> , 2003 , 43, 121-34		230
137	Regulatory flexibility in the Nrf2-mediated stress response is conferred by conformational cycling of the Keap1-Nrf2 protein complex. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 15259-64	11.5	228
136	Persuasive evidence that quinone reductase type 1 (DT diaphorase) protects cells against the toxicity of electrophiles and reactive forms of oxygen. <i>Free Radical Biology and Medicine</i> , 2000 , 29, 231-40	7.8	202
135	Relation of structure of curcumin analogs to their potencies as inducers of Phase 2 detoxification enzymes. <i>Carcinogenesis</i> , 1999 , 20, 911-4	4.6	202
134	Direct and indirect antioxidant properties of inducers of cytoprotective proteins. <i>Molecular Nutrition and Food Research</i> , 2008 , 52 Suppl 1, S128-38	5.9	199
133	The multifaceted role of Nrf2 in mitochondrial function. <i>Current Opinion in Toxicology</i> , 2016 , 1, 80-91	4.4	198
132	Loss of Nrf2 markedly exacerbates nonalcoholic steatohepatitis. <i>Free Radical Biology and Medicine</i> , 2010 , 48, 357-71	7.8	190
131	Protection against UV-light-induced skin carcinogenesis in SKH-1 high-risk mice by sulforaphane-containing broccoli sprout extracts. <i>Cancer Letters</i> , 2006 , 240, 243-52	9.9	183
130	Keap1, the sensor for electrophiles and oxidants that regulates the phase 2 response, is a zinc metalloprotein. <i>Biochemistry</i> , 2005 , 44, 6889-99	3.2	176
129	Sulforaphane mobilizes cellular defenses that protect skin against damage by UV radiation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 17500-5	11.5	165
128	Nitric oxide in cell survival: a janus molecule. <i>Antioxidants and Redox Signaling</i> , 2009 , 11, 2717-39	8.4	160
127	Regiochemical control of monolignol radical coupling: a new paradigm for lignin and lignan biosynthesis. <i>Chemistry and Biology</i> , 1999 , 6, 143-51		157
126	Chemoprotective properties of phenylpropenoids, bis(benzylidene)cycloalkanones, and related Michael reaction acceptors: correlation of potencies as phase 2 enzyme inducers and radical scavengers. <i>Journal of Medicinal Chemistry</i> , 1998 , 41, 5287-96	8.3	155
125	(+)-Pinoresinol/(+)-lariciresinol reductase from <i>Forsythia intermedia</i> . Protein purification, cDNA cloning, heterologous expression and comparison to isoflavone reductase. <i>Journal of Biological Chemistry</i> , 1996 , 271, 29473-82	5.4	152

124	Keap1, the cysteine-based mammalian intracellular sensor for electrophiles and oxidants. <i>Archives of Biochemistry and Biophysics</i> , 2017 , 617, 84-93	4.1	151
123	Nrf2 affects the efficiency of mitochondrial fatty acid oxidation. <i>Biochemical Journal</i> , 2014 , 457, 415-24	3.8	148
122	Susceptibility of Nrf2-null mice to steatohepatitis and cirrhosis upon consumption of a high-fat diet is associated with oxidative stress, perturbation of the unfolded protein response, and disturbance in the expression of metabolic enzymes but not with insulin resistance. <i>Molecular and Cellular Biology</i> , 2014 , 34, 3305-20	4.8	141
121	The role of Nrf2 signaling in counteracting neurodegenerative diseases. <i>FEBS Journal</i> , 2018 , 285, 3576-3590	3.9	137
120	Induction of the phase 2 response in mouse and human skin by sulforaphane-containing broccoli sprout extracts. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2007 , 16, 847-51	4	134
119	Electrophilic tuning of the chemoprotective natural product sulforaphane. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 9590-5	11.5	133
118	KEAP1 and Done? Targeting the NRF2 Pathway with Sulforaphane. <i>Trends in Food Science and Technology</i> , 2017 , 69, 257-269	15.3	128
117	Protection against electrophile and oxidative stress by induction of phase 2 genes: the quest for the elusive sensor that responds to inducers. <i>Advances in Enzyme Regulation</i> , 2004 , 44, 335-67		119
116	Targeting the CoREST complex with dual histone deacetylase and demethylase inhibitors. <i>Nature Communications</i> , 2018 , 9, 53	17.4	116
115	Curcumin and the cellular stress response in free radical-related diseases. <i>Molecular Nutrition and Food Research</i> , 2008 , 52, 1062-73	5.9	115
114	The "Prochaska" microtiter plate bioassay for inducers of NQO1. <i>Methods in Enzymology</i> , 2004 , 382, 243-58	5.8	115
113	Dual regulation of transcription factor Nrf2 by Keap1 and by the combined actions of I κ B α and GSK-3. <i>Biochemical Society Transactions</i> , 2015 , 43, 611-20	5.1	104
112	Broccoli or Sulforaphane: Is It the Source or Dose That Matters?. <i>Molecules</i> , 2019 , 24,	4.8	103
111	Experimental Nonalcoholic Steatohepatitis and Liver Fibrosis Are Ameliorated by Pharmacologic Activation of Nrf2 (NF-E2 p45-Related Factor 2). <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2018 , 5, 367-398	7.9	101
110	Coordinate regulation of enzyme markers for inflammation and for protection against oxidants and electrophiles. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 15926-31	11.5	101
109	Phytochemicals as protectors against ultraviolet radiation: versatility of effects and mechanisms. <i>Planta Medica</i> , 2008 , 74, 1548-59	3.1	97
108	Chemical structures of inducers of nicotinamide quinone oxidoreductase 1 (NQO1). <i>Methods in Enzymology</i> , 2004 , 382, 423-48	1.7	97
107	Monitoring Keap1-Nrf2 interactions in single live cells. <i>Biotechnology Advances</i> , 2014 , 32, 1133-44	17.8	92

106	Oxidative stress and chronic inflammation in osteoarthritis: can NRF2 counteract these partners in crime?. <i>Annals of the New York Academy of Sciences</i> , 2017 , 1401, 114-135	6.5	90
105	An exceptionally potent inducer of cytoprotective enzymes: elucidation of the structural features that determine inducer potency and reactivity with Keap1. <i>Journal of Biological Chemistry</i> , 2010 , 285, 33747-55	5.4	90
104	Can Activation of NRF2 Be a Strategy against COVID-19?. <i>Trends in Pharmacological Sciences</i> , 2020 , 41, 598-610	13.2	90
103	Nrf2 activation in the treatment of neurodegenerative diseases: a focus on its role in mitochondrial bioenergetics and function. <i>Biological Chemistry</i> , 2016 , 397, 383-400	4.5	89
102	Regulation of the mammalian heat shock factor 1. <i>FEBS Journal</i> , 2017 , 284, 1606-1627	5.7	86
101	Design, Synthesis, and Evaluation of Triazole Derivatives That Induce Nrf2 Dependent Gene Products and Inhibit the Keap1-Nrf2 Protein-Protein Interaction. <i>Journal of Medicinal Chemistry</i> , 2015 , 58, 7186-94	8.3	86
100	Chlorophyll, chlorophyllin and related tetrapyrroles are significant inducers of mammalian phase 2 cytoprotective genes. <i>Carcinogenesis</i> , 2005 , 26, 1247-55	4.6	85
99	Activation of the NRF2 signaling pathway by copper-mediated redox cycling of para- and ortho-hydroquinones. <i>Chemistry and Biology</i> , 2010 , 17, 75-85		84
98	A Defective Pentose Phosphate Pathway Reduces Inflammatory Macrophage Responses during Hypercholesterolemia. <i>Cell Reports</i> , 2018 , 25, 2044-2052.e5	10.6	84
97	Transcription factors Hsf1 and Nrf2 engage in crosstalk for cytoprotection. <i>Trends in Pharmacological Sciences</i> , 2015 , 36, 6-14	13.2	82
96	HSF1-dependent upregulation of Hsp70 by sulfhydryl-reactive inducers of the KEAP1/NRF2/ARE pathway. <i>Chemistry and Biology</i> , 2011 , 18, 1355-61		78
95	Cross-talk between transcription factors AhR and Nrf2: lessons for cancer chemoprevention from dioxin. <i>Toxicological Sciences</i> , 2009 , 111, 199-201	4.4	76
94	Reprogramming of keratin biosynthesis by sulforaphane restores skin integrity in epidermolysis bullosa simplex. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 14460-5	11.5	74
93	Nrf2 Activation Protects against Solar-Simulated Ultraviolet Radiation in Mice and Humans. <i>Cancer Prevention Research</i> , 2015 , 8, 475-86	3.2	71
92	Vitagenes, cellular stress response, and acetylcarnitine: relevance to hormesis. <i>BioFactors</i> , 2009 , 35, 146-60		67
91	Role of nicotinamide quinone oxidoreductase 1 (NQO1) in protection against toxicity of electrophiles and reactive oxygen intermediates. <i>Methods in Enzymology</i> , 2004 , 382, 355-64	1.7	66
90	KEAP1-modifying small molecule reveals muted NRF2 signaling responses in neural stem cells from Huntington's disease patients. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017 , 114, E4676-E4685	11.5	65
89	KEAP1 inhibition is neuroprotective and suppresses the development of epilepsy. <i>Brain</i> , 2018 , 141, 1390-1403	14.03	58

88	The spatiotemporal regulation of the Keap1-Nrf2 pathway and its importance in cellular bioenergetics. <i>Biochemical Society Transactions</i> , 2015 , 43, 602-10	5.1	58
87	Induction of the Keap1/Nrf2/ARE pathway by oxidizable diphenols. <i>Chemico-Biological Interactions</i> , 2011 , 192, 101-6	5	58
86	SIRT2- and NRF2-Targeting Thiazole-Containing Compound with Therapeutic Activity in Huntington@ Disease Models. <i>Cell Chemical Biology</i> , 2016 , 23, 849-861	8.2	54
85	Neuroprotective effects of sulforaphane after contusive spinal cord injury. <i>Journal of Neurotrauma</i> , 2012 , 29, 2576-86	5.4	53
84	Novel semisynthetic analogues of betulinic acid with diverse cytoprotective, antiproliferative, and proapoptotic activities. <i>Molecular Cancer Therapeutics</i> , 2007 , 6, 2113-9	6.1	53
83	Activation of Nrf2 Signaling Augments Vesicular Stomatitis Virus Oncolysis via Autophagy-Driven Suppression of Antiviral Immunity. <i>Molecular Therapy</i> , 2017 , 25, 1900-1916	11.7	48
82	Synthesis, chemical reactivity as Michael acceptors, and biological potency of monocyclic cyanoenones, novel and highly potent anti-inflammatory and cytoprotective agents. <i>Journal of Medicinal Chemistry</i> , 2012 , 55, 4837-46	8.3	47
81	Two-step mechanism of induction of the gene expression of a prototypic cancer-protective enzyme by diphenols. <i>Chemical Research in Toxicology</i> , 2008 , 21, 805-12	4	44
80	A novel acetylenic tricyclic bis-(cyano enone) potently induces phase 2 cytoprotective pathways and blocks liver carcinogenesis induced by aflatoxin. <i>Cancer Research</i> , 2008 , 68, 6727-33	10.1	44
79	Heat Shock Factor 1 Is a Substrate for p38 Mitogen-Activated Protein Kinases. <i>Molecular and Cellular Biology</i> , 2016 , 36, 2403-17	4.8	44
78	Pathogenic p62/SQSTM1 mutations impair energy metabolism through limitation of mitochondrial substrates. <i>Scientific Reports</i> , 2017 , 7, 1666	4.9	43
77	Chemoprotection against cancer by isothiocyanates: a focus on the animal models and the protective mechanisms. <i>Topics in Current Chemistry</i> , 2013 , 329, 179-201		42
76	Obesity and NRF2-mediated cytoprotection: Where is the missing link?. <i>Pharmacological Research</i> , 2020 , 156, 104760	10.2	39
75	Diffusion dynamics of the Keap1-Cullin3 interaction in single live cells. <i>Biochemical and Biophysical Research Communications</i> , 2013 , 433, 58-65	3.4	39
74	The indirect antioxidant sulforaphane protects against thiopurine-mediated photooxidative stress. <i>Carcinogenesis</i> , 2012 , 33, 2457-66	4.6	34
73	Dietary glucoraphanin-rich broccoli sprout extracts protect against UV radiation-induced skin carcinogenesis in SKH-1 hairless mice. <i>Photochemical and Photobiological Sciences</i> , 2010 , 9, 597-600	4.2	33
72	C151 in KEAP1 is the main cysteine sensor for the cyanoenone class of NRF2 activators, irrespective of molecular size or shape. <i>Scientific Reports</i> , 2018 , 8, 8037	4.9	33
71	Redox ranking of inducers of a cancer-protective enzyme via the energy of their highest occupied molecular orbital. <i>Free Radical Biology and Medicine</i> , 2004 , 36, 1418-23	7.8	31

70	KEAP1, a cysteine-based sensor and a drug target for the prevention and treatment of chronic disease. <i>Open Biology</i> , 2020 , 10, 200105	7	31
69	Practical approaches to investigate redox regulation of heat shock protein expression and intracellular glutathione redox state. <i>Methods in Enzymology</i> , 2008 , 441, 83-110	1.7	30
68	Highly potent activation of Nrf2 by topical tricyclic bis(cyano enone): implications for protection against UV radiation during thiopurine therapy. <i>Cancer Prevention Research</i> , 2012 , 5, 973-81	3.2	29
67	Transcription factors NRF2 and HSF1 have opposing functions in autophagy. <i>Scientific Reports</i> , 2017 , 7, 11023	4.9	26
66	New Monocyclic, Bicyclic, and Tricyclic Ethynylcyanodienones as Activators of the Keap1/Nrf2/ARE Pathway and Inhibitors of Inducible Nitric Oxide Synthase. <i>Journal of Medicinal Chemistry</i> , 2015 , 58, 4738-48	8.3	26
65	Rhodiola rosea L.: from golden root to green cell factories. <i>Phytochemistry Reviews</i> , 2016 , 15, 515-536	7.7	25
64	Biomarker Exploration in Human Peripheral Blood Mononuclear Cells for Monitoring Sulforaphane Treatment Responses in Autism Spectrum Disorder. <i>Scientific Reports</i> , 2020 , 10, 5822	4.9	25
63	Flavonolignan 2,3-dehydrosilydianin activates Nrf2 and upregulates NAD(P)H:quinone oxidoreductase 1 in Hepa1c1c7 cells. <i>Phytotherapy Research</i> , 2017 , 119, 115-120	3.2	24
62	Phenethyl Isothiocyanate, a Dual Activator of Transcription Factors NRF2 and HSF1. <i>Molecular Nutrition and Food Research</i> , 2018 , 62, e1700908	5.9	24
61	Nrf2-Mediated Neuroprotection Against Recurrent Hypoglycemia Is Insufficient to Prevent Cognitive Impairment in a Rodent Model of Type 1 Diabetes. <i>Diabetes</i> , 2016 , 65, 3151-60	0.9	24
60	Loss of Nrf2 abrogates the protective effect of Keap1 downregulation in a preclinical model of cutaneous squamous cell carcinoma. <i>Scientific Reports</i> , 2016 , 6, 25804	4.9	23
59	Oxidative stress management in the hair follicle: Could targeting NRF2 counter age-related hair disorders and beyond?. <i>BioEssays</i> , 2017 , 39, 1700029	4.1	20
58	The Role of Sulfhydryl Reactivity of Small Molecules for the Activation of the KEAP1/NRF2 Pathway and the Heat Shock Response. <i>Scientifica</i> , 2012 , 2012, 606104	2.6	20
57	NRF2 and the Ambiguous Consequences of Its Activation during Initiation and the Subsequent Stages of Tumourigenesis. <i>Cancers</i> , 2020 , 12,	6.6	20
56	Bis(2-hydroxybenzylidene)acetone, a potent inducer of the phase 2 response, causes apoptosis in mouse leukemia cells through a p53-independent, caspase-mediated pathway. <i>Cancer Letters</i> , 2007 , 245, 341-9	9.9	19
55	Chemical tuning enhances both potency toward nrf2 and in vitro therapeutic index of triterpenoids. <i>Toxicological Sciences</i> , 2014 , 140, 462-9	4.4	17
54	Molecular basis for the disruption of Keap1-Nrf2 interaction via Hinge & Latch mechanism. <i>Communications Biology</i> , 2021 , 4, 576	6.7	17
53	Activation of transcription factor Nrf2 to counteract mitochondrial dysfunction in Parkinson's disease. <i>Medicinal Research Reviews</i> , 2021 , 41, 785-802	14.4	17

52	Pharmacokinetics and pharmacodynamics of orally administered acetylenic tricyclic bis(cyanoenone), a highly potent Nrf2 activator with a reversible covalent mode of action. <i>Biochemical and Biophysical Research Communications</i> , 2015 , 465, 402-7	3.4	16
51	Synthesis and biological evaluation of 1-[2-cyano-3,12-dioxooleana-1,9(11)-dien-28-oyl]-4-ethynylimidazole. A novel and highly potent anti-inflammatory and cytoprotective agent. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2011 , 21, 2188-91	2.9	15
50	Oncogene-Stimulated Congestion at the KEAP1 Stress Signaling Hub Allows Bypass of NRF2 and Induction of NRF2-Target Genes that Promote Tumor Survival. <i>Cancer Cell</i> , 2017 , 32, 539-541	24.3	14
49	Oral azathioprine leads to higher incorporation of 6-thioguanine in DNA of skin than liver: the protective role of the Keap1/Nrf2/ARE pathway. <i>Cancer Prevention Research</i> , 2011 , 4, 1665-74	3.2	14
48	Epigenetic Control of NRF2-Directed Cellular Antioxidant Status in Dictating Life-Death Decisions. <i>Molecular Cell</i> , 2017 , 68, 5-7	17.6	14
47	Whole-Exome Sequencing Validates a Preclinical Mouse Model for the Prevention and Treatment of Cutaneous Squamous Cell Carcinoma. <i>Cancer Prevention Research</i> , 2017 , 10, 67-75	3.2	13
46	A dicyanotriterpenoid induces cytoprotective enzymes and reduces multiplicity of skin tumors in UV-irradiated mice. <i>Biochemical and Biophysical Research Communications</i> , 2008 , 367, 859-65	3.4	13
45	Regulation of the CNC-bZIP transcription factor Nrf2 by Keap1 and the axis between GSK-3 and ETrCP. <i>Current Opinion in Toxicology</i> , 2016 , 1, 92-103	4.4	12
44	High NRF2 Levels Correlate with Poor Prognosis in Colorectal Cancer Patients and with Sensitivity to the Kinase Inhibitor AT9283 In Vitro. <i>Biomolecules</i> , 2020 , 10,	5.9	12
43	The isothiocyanate sulforaphane inhibits mTOR in an NRF2-independent manner. <i>Phytomedicine</i> , 2021 , 86, 153062	6.5	12
42	Precise determination of the erythema response of human skin to ultraviolet radiation and quantification of effects of protectors. <i>Photodermatology Photoimmunology and Photomedicine</i> , 2009 , 25, 45-50	2.4	11
41	Semisynthetic flavonoid 7-O-galloylquercetin activates Nrf2 and induces Nrf2-dependent gene expression in RAW264.7 and Hepa1c1c7 cells. <i>Chemico-Biological Interactions</i> , 2016 , 260, 58-66	5	10
40	NAD(P)H : quinone oxidoreductase 1 inducer activity of some Saudi Arabian medicinal plants. <i>Planta Medica</i> , 2013 , 79, 459-64	3.1	10
39	Potency ranking of triterpenoids as inducers of a cytoprotective enzyme and as inhibitors of a cellular inflammatory response via their electron affinity and their electrophilicity index. <i>Chemico-Biological Interactions</i> , 2010 , 186, 118-26	5	10
38	Rapid body weight gain increases the risk of UV radiation-induced skin carcinogenesis in SKH-1 hairless mice. <i>Nutrition Research</i> , 2008 , 28, 539-43	4	10
37	Isomeric O-methyl cannabidiolquinones with dual BACH1/NRF2 activity. <i>Redox Biology</i> , 2020 , 37, 101689	11.3	9
36	Nrf2 is activated by disruption of mitochondrial thiol homeostasis but not by enhanced mitochondrial superoxide production. <i>Journal of Biological Chemistry</i> , 2021 , 296, 100169	5.4	9
35	Downregulation of Keap1 Confers Features of a Fasted Metabolic State. <i>IScience</i> , 2020 , 23, 101638	6.1	8

34	Potency of extracts from selected Egyptian plants as inducers of the Nrf2-dependent chemopreventive enzyme NQO1. <i>Journal of Natural Medicines</i> , 2016 , 70, 683-8	3.3	8
33	Investigation into the use of histone deacetylase inhibitor MS-275 as a topical agent for the prevention and treatment of cutaneous squamous cell carcinoma in an SKH-1 hairless mouse model. <i>PLoS ONE</i> , 2019 , 14, e0213095	3.7	7
32	Measuring the Interaction of Transcription Factor Nrf2 with Its Negative Regulator Keap1 in Single Live Cells by an Improved FRET/FLIM Analysis. <i>Chemical Research in Toxicology</i> , 2019 , 32, 500-512	4	6
31	Filaggrin genotype does not determine the skin threshold to UV-induced erythema. <i>Journal of Allergy and Clinical Immunology</i> , 2016 , 137, 1280-1282.e3	11.5	6
30	Synthesis, molecular modeling and NAD(P)H:quinone oxidoreductase 1 inducer activity of novel cyanoenone and enone benzenesulfonamides. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2014 , 29, 840-5	5.6	6
29	Phenolics in Aging and Neurodegenerative Disorders 427-451		6
28	Radiomodulatory effect of a non-electrophilic NQO1 inducer identified in a screen of new 6, 8-diiodoquinazolin-4(3H)-ones carrying a sulfonamide moiety. <i>European Journal of Medicinal Chemistry</i> , 2020 , 200, 112467	6.8	6
27	NAD(P)H:quinone oxidoreductase 1 inducer activity of some novel anilinoquinazoline derivatives. <i>Drug Design, Development and Therapy</i> , 2016 , 10, 2515-24	4.4	5
26	The stress-responsive kinase DYRK2 activates heat shock factor 1 promoting resistance to proteotoxic stress. <i>Cell Death and Differentiation</i> , 2021 , 28, 1563-1578	12.7	5
25	Studies on the mechanism of anti-inflammatory action of swietenine, a tetranortriterpenoid isolated from <i>Swietenia macrophylla</i> seeds. <i>Phytomedicine Plus</i> , 2021 , 1, 100018		5
24	The Cell-Permeable Derivative of the Immunoregulatory Metabolite Itaconate, 4-Octyl Itaconate, Is Anti-Fibrotic in Systemic Sclerosis. <i>Cells</i> , 2021 , 10,	7.9	5
23	Nrf2 activation reprograms macrophage intermediary metabolism and suppresses the type I interferon response.. <i>iScience</i> , 2022 , 25, 103827	6.1	4
22	Novel iodinated quinazolinones bearing sulfonamide as new scaffold targeting radiation induced oxidative stress. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2021 , 42, 128002	2.9	4
21	Synthesis, molecular modeling and NAD(P)H:quinone oxidoreductase 1 inducer activity of novel 2-phenylquinazolin-4-amine derivatives. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2016 , 31, 1612-8	5.6	4
20	Synthesis and biological evaluation of novel 2-phenylquinazoline-4-amine derivatives: identification of 6-phenyl-8H-benzo[g]quinazolino[4,3-b]quinazolin-8-one as a highly potent inducer of NAD(P)H quinone oxidoreductase 1. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2016 , 31, 34-39	5.6	4
19	Chemoprotection against cancer: an idea whose time has come. <i>Alternative Therapies in Health and Medicine</i> , 2007 , 13, S122-7	2.5	4
18	NAD(P)H: quinone oxidoreductase 1 inducer activity of novel 4-aminoquinazoline derivatives. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2016 , 31, 1369-74	5.6	3
17	The isoquinoline PRL-295 increases the thermostability of Keap1 and disrupts its interaction with Nrf2.. <i>iScience</i> , 2022 , 25, 103703	6.1	3

16	Assessment of ROS Production in the Mitochondria of Live Cells. <i>Methods in Molecular Biology</i> , 2021 , 2202, 33-42	1.4	3
15	Electron affinity of tricyclic, bicyclic, and monocyclic compounds containing cyanoenones correlates with their potency as inducers of a cytoprotective enzyme. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2016 , 26, 4345-9	2.9	2
14	Synthesis and biological evaluation of biotin conjugates of (E)-(4bS,8aR,10aS)-10a-ethynyl-4b,8,8-trimethyl-3,7-dioxo-3,4b,7,8,8a,9,10,10a-octahydro-phenanthrene-2,6-dicarbonitril as an activator of the Keap1/Nrf2/ARE pathway, for the isolation of its protein targets. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2013 , 23, 5540-3	2.9	2
13	Novel Thioureido Derivatives Carrying Thione and Sulfonamide Moieties Induce the Cytoprotective Enzyme NAD(P)H:Quinone Oxidoreductase 1. <i>Asian Journal of Chemistry</i> , 2014 , 26, 8501-8504	0.4	2
12	Synthesis of (13) C2 (15) N2 -labeled anti-inflammatory and cytoprotective tricyclic bis(cyanoenone) ([[(13) C2 (15) N2]-TBE-31) as an internal standard for quantification by stable isotope dilution LC-MS method. <i>Journal of Labelled Compounds and Radiopharmaceuticals</i> , 2014 , 57, 606-10	1.9	2
11	The Isothiocyanate Sulforaphane Induces the Phase 2 Response by Signaling of the Keap1/Nrf2/ARE Pathway. <i>Oxidative Stress and Disease</i> , 2008 ,		2
10	The Chemopreventive Power of Isothiocyanates 2020 , 271-318		2
9	DYRK2 activates heat shock factor 1 promoting resistance to proteotoxic stress in triplenegative breast cancer		2
8	Sulfhydryl-Reactive Phytochemicals as Dual Activators of Transcription Factors NRF2 and HSF1 2013 , 95-119		1
7	Sulfoxythiocarbamate S-4 inhibits HSP90 in human cutaneous squamous cell carcinoma cells. <i>European Journal of Pharmacology</i> , 2020 , 889, 173609	5.3	1
6	Measuring Changes in Keap1-Nrf2 Protein Complex Conformation in Individual Cells by FLIM-FRET. <i>Current Protocols in Toxicology / Editorial Board, Mahin D Maines (editor-in-chief) [et Al]</i> , 2020 , 85, e96	1	1
5	Clinically relevant aberrant Filip1l DNA methylation detected in a murine model of cutaneous squamous cell carcinoma. <i>EBioMedicine</i> , 2021 , 67, 103383	8.8	1
4	Application of the in vivo oxidative stress reporter Hmox1 as mechanistic biomarker of arsenic toxicity. <i>Environmental Pollution</i> , 2021 , 270, 116053	9.3	1
3	Nrf2 activation does not affect adenoma development in a mouse model of colorectal cancer. <i>Communications Biology</i> , 2021 , 4, 1081	6.7	0
2	The synthetic triterpenoids CDDO-TFEA and CDDO-Me, but not CDDO, promote nuclear exclusion of BACH1 impairing its activity.. <i>Redox Biology</i> , 2022 , 51, 102291	11.3	0
1	Detection of thermal shift in cellular Keap1 by protein-protein interaction inhibitors using immunoblot- and fluorescence microplate-based assays.. <i>STAR Protocols</i> , 2022 , 3, 101265	1.4	0