

Jing Wu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7334421/publications.pdf>

Version: 2024-02-01

16
papers

916
citations

567281

15
h-index

940533

16
g-index

16
all docs

16
docs citations

16
times ranked

1694
citing authors

#	ARTICLE	IF	CITATIONS
1	Klotho is a target gene of PPAR- β . <i>Kidney International</i> , 2008, 74, 732-739.	5.2	142
2	CD70 Exacerbates Blood Pressure Elevation and Renal Damage in Response to Repeated Hypertensive Stimuli. <i>Circulation Research</i> , 2016, 118, 1233-1243.	4.5	128
3	Antihypertensive effects of selective prostaglandin E2 receptor subtype 1 targeting. <i>Journal of Clinical Investigation</i> , 2007, 117, 2496-2505.	8.2	94
4	Thyroid hormone-responsive SPOT 14 homolog promotes hepatic lipogenesis, and its expression is regulated by Liver X receptor β through a sterol regulatory element-binding protein 1c-dependent mechanism in mice. <i>Hepatology</i> , 2013, 58, 617-628.	7.3	72
5	Peroxisome Proliferator-Activated Receptor β / β Dual Agonist Tesaglitazar Attenuates Diabetic Nephropathy in db/db Mice. <i>Diabetes</i> , 2007, 56, 2036-2045.	0.6	70
6	Proteomics analysis reveals diabetic kidney as a ketogenic organ in type 2 diabetes. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2011, 300, E287-E295.	3.5	67
7	Pancreatic-derived factor promotes lipogenesis in the mouse liver: Role of the Forkhead box 1 signaling pathway. <i>Hepatology</i> , 2011, 53, 1906-1916.	7.3	53
8	Liver X receptor β mediates cholesterol efflux in glomerular mesangial cells. <i>American Journal of Physiology - Renal Physiology</i> , 2004, 287, F886-F895.	2.7	48
9	Liver X receptor agonist TO-901317 upregulates SCD1 expression in renal proximal straight tubule. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 290, F1065-F1073.	2.7	42
10	Expression profiling of hepatic genes associated with lipid metabolism in nephrotic rats. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 295, F662-F671.	2.7	37
11	Induction of MIF expression by oxidized LDL via activation of NF- κ B in vascular smooth muscle cells. <i>Atherosclerosis</i> , 2009, 207, 428-433.	0.8	35
12	Liver X receptor activation increases hepatic fatty acid desaturation by the induction of SCD1 expression through an LXR- β -dependent mechanism (LXR β -dependent SREBP1c). <i>Journal of Diabetes</i> (2010), 1(2), 1-6.	1.8	35
13	Plasma Periostin Levels Are Increased in Chinese Subjects with Obesity and Type 2 Diabetes and Are Positively Correlated with Glucose and Lipid Parameters. <i>Mediators of Inflammation</i> , 2016, 2016, 1-6.	3.0	31
14	Expression of mouse membrane-associated prostaglandin E2 synthase-2 (mPGES-2) along the urogenital tract. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2006, 1761, 1459-1468.	2.4	27
15	Peroxisome proliferator-activated receptors and renal diseases. <i>Frontiers in Bioscience - Landmark</i> , 2009, Volume, 995.	3.0	24
16	The effect of the EP3 antagonist DG-041 on male mice with diet-induced obesity. <i>Prostaglandins and Other Lipid Mediators</i> , 2019, 144, 106353.	1.9	11