

# Yingying Ding

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/733837/publications.pdf>

Version: 2024-02-01

50  
papers

669  
citations

623734

14  
h-index

642732

23  
g-index

52  
all docs

52  
docs citations

52  
times ranked

1077  
citing authors

#	ARTICLE	IF	CITATIONS
1	Low willingness and actual uptake of pre-exposure prophylaxis for HIV-1 prevention among men who have sex with men in Shanghai, China. <i>BioScience Trends</i> , 2016, 10, 113-119.	3.4	57
2	Adverse childhood experiences and interaction with methamphetamine use frequency in the risk of methamphetamine-associated psychosis. <i>Drug and Alcohol Dependence</i> , 2014, 142, 295-300.	3.2	56
3	A multilevel intervention for HIV-affected families in China: Together for Empowerment Activities (TEA). <i>Social Science and Medicine</i> , 2011, 73, 1214-1221.	3.8	42
4	Illicit Heroin and Methamphetamine Use among Methadone Maintenance Treatment Patients in Dehong Prefecture of Yunnan Province, China. <i>PLoS ONE</i> , 2015, 10, e0133431.	2.5	39
5	Higher Prevalence of Frailty Among a Sample of HIV-Infected Middle-aged and Older Chinese Adults Is Associated With Neurocognitive Impairment and Depressive Symptoms. <i>Journal of Infectious Diseases</i> , 2017, 215, 687-692.	4.0	36
6	Social support and depressive symptoms among 'money' boys and general men who have sex with men in Shanghai, China. <i>Sexual Health</i> , 2014, 11, 285.	0.9	24
7	Interaction Effects between HIV and Aging on Selective Neurocognitive Impairment. <i>Journal of NeuroImmune Pharmacology</i> , 2017, 12, 661-669.	4.1	24
8	Factors influencing uptake of pre-exposure prophylaxis: some qualitative insights from an intervention study of men who have sex with men in China. <i>Sexual Health</i> , 2018, 15, 39.	0.9	24
9	Circumstances of initiation into new-type drug use among adults in Shanghai: Are there differences by types of first new-type drug used?. <i>Drug and Alcohol Dependence</i> , 2013, 131, 278-283.	3.2	23
10	Age-specific associations between HIV infection and carotid artery intima-media thickness in China: a cross-sectional evaluation of baseline data from the CHART cohort. <i>Lancet HIV</i> , 2019, 6, e860-e868.	4.7	23
11	Evolving HIV Epidemiology in Mainland China: 2009–2018. <i>Current HIV/AIDS Reports</i> , 2019, 16, 423-430.	3.1	23
12	More improvement than progression of liver fibrosis following antiretroviral therapy in a longitudinal cohort of HIV-infected patients with or without HBV and HCV co-infections. <i>Journal of Viral Hepatitis</i> , 2017, 24, 412-420.	2.0	22
13	Prevalence and genotypes of anal human papillomavirus infection among HIV-positive vs HIV-negative men in Taizhou, China. <i>Epidemiology and Infection</i> , 2019, 147, e117.	2.1	19
14	Tobacco use among HIV-infected individuals in a rural community in Yunnan Province, China. <i>Drug and Alcohol Dependence</i> , 2014, 134, 144-150.	3.2	18
15	Tobacco and alcohol use among drug users receiving methadone maintenance treatment: a cross-sectional study in a rural prefecture of Yunnan Province, Southwest China. <i>BMJ Open</i> , 2017, 7, e014643.	1.9	17
16	Cross-sectional comparison of various sleep disturbances among sex- and age-matched HIV-infected versus HIV-uninfected individuals in China. <i>Sleep Medicine</i> , 2020, 65, 18-25.	1.6	15
17	Epidemiological and molecular characteristics of HIV infection among money boys and general men who have sex with men in Shanghai, China. <i>Infection, Genetics and Evolution</i> , 2015, 31, 135-141.	2.3	14
18	Intimate partner violence victimization and HIV infection among men who have sex with men in Shanghai, China. <i>BioScience Trends</i> , 2018, 12, 142-148.	3.4	13

#	ARTICLE	IF	CITATIONS
19	Hypertension in HIV-Infected Adults Compared with Similar but Uninfected Adults in China: Body Mass Index-Dependent Effects of Nadir CD4 Count. <i>AIDS Research and Human Retroviruses</i> , 2017, 33, 1117-1125.	1.1	12
20	Comprehensive metabolomics profiling reveals common metabolic alterations underlying the four major non-communicable diseases in treated HIV infection. <i>EBioMedicine</i> , 2021, 71, 103548.	6.1	12
21	High Prevalence of HIV Infection and Bisexual Networks among a Sample of Men Who Have Sex with Men in Eastern China. <i>PLoS ONE</i> , 2015, 10, e0129300.	2.5	11
22	Sex differences in neurocognitive screening among adults living with HIV in China. <i>Journal of NeuroVirology</i> , 2019, 25, 363-371.	2.1	11
23	Severity of club drug dependence and perceived need for treatment among a sample of adult club drug users in Shanghai, China. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2014, 49, 395-404.	3.1	10
24	Timing of antiretroviral therapy initiation after diagnosis of recent human immunodeficiency virus infection and CD4+ T-cell recovery. <i>Clinical Microbiology and Infection</i> , 2016, 22, 290.e5-290.e8.	6.0	9
25	Stronger Association between Insomnia Symptoms and Shorter Telomere Length in Old HIV-Infected Patients Compared with Uninfected Individuals. , 2018, 9, 1010.		9
26	HIV and Aging in Mainland China: Implications for Control and Prevention Research. <i>Current HIV/AIDS Reports</i> , 2019, 16, 439-447.	3.1	9
27	HIV infection and electrocardiogram abnormalities: baseline assessment from the CHART cohort. <i>Clinical Microbiology and Infection</i> , 2020, 26, 1689.e1-1689.e7.	6.0	9
28	Mitochondrial Haplogroups N9 and G Are Associated with Metabolic Syndrome Among Human Immunodeficiency Virus-Infected Patients in China. <i>AIDS Research and Human Retroviruses</i> , 2019, 35, 536-543.	1.1	8
29	Differential prevalence and correlates of whole blood Epstein-Barr virus DNA between HIV-positive and HIV-negative men who have sex with men in Shanghai, China. <i>Epidemiology and Infection</i> , 2017, 145, 2330-2340.	2.1	7
30	Effects of aging, baseline renal function and stage of HIV infection on post-treatment changes in renal function among HIV-infected patients: a retrospective cohort study. <i>HIV Medicine</i> , 2019, 20, 591-600.	2.2	7
31	Risk assessment and identification of HIV infection among men who have sex with men: a cross-sectional study in Southwest China. <i>BMJ Open</i> , 2020, 10, e039557.	1.9	7
32	HIV drug resistance in newly diagnosed adults in a rural prefecture of eastern China. <i>Epidemiology and Infection</i> , 2015, 143, 663-672.	2.1	6
33	Liver fibrosis after antiretroviral therapy in a longitudinal cohort of sexually infected HIV patients in eastern China. <i>BioScience Trends</i> , 2017, 11, 274-281.	3.4	6
34	Sex with older partners, condomless anal sex and unrecognized HIV infection among Chinese men who have sex with men. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2018, 30, 305-311.	1.2	5
35	Fibrosis-4 index predicts mortality in HIV/HCV co-infected patients receiving combination antiretroviral therapy in rural China. <i>BioScience Trends</i> , 2019, 13, 32-39.	3.4	4
36	General and abdominal obesity and incident hypertension among people living with HIV on antiretroviral therapy. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2021, 33, 1616-1620.	1.2	4

#	ARTICLE	IF	CITATIONS
37	Electrocardiographic abnormalities among people with HIV in Shanghai, China. <i>BioScience Trends</i> , 2020, 14, 9-15.	3.4	4
38	Effects of Docosahexanoic Acid on Gut Microbiota and Fecal Metabolites in HIV-Infected Patients With Neurocognitive Impairment: A 6-Month Randomized, Double-Blind, Placebo-Controlled Trial. <i>Frontiers in Nutrition</i> , 2021, 8, 756720.	3.7	4
39	Health-care access and utilization among HIV-infected men who have sex with men in two Chinese municipalities with or without lockdown amidst early COVID-19 pandemic. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2022, 34, 1390-1399.	1.2	4
40	Adolescent Sexual Debut and Initiation into New-Type Drug Use among a Sample of Young Adults. <i>Journal of Psychoactive Drugs</i> , 2015, 47, 182-186.	1.7	3
41	Risk factors for incident HSV-2 infections among a prospective cohort of HIV-1-discordant couples in China. <i>Sexually Transmitted Infections</i> , 2016, 92, 76-82.	1.9	3
42	Incidence of impaired kidney function among people with HIV: a systematic review and meta-analysis. <i>BMC Nephrology</i> , 2022, 23, 107.	1.8	3
43	Increased CD4+ T cell count is associated with lower anal human papillomavirus prevalence among HIV-positive male cohort in Taizhou, China: a cross-sectional study. <i>BMC Infectious Diseases</i> , 2022, 22, 250.	2.9	3
44	Low body mass index and efavirenz use are independently associated with self-reported fatigue in HIV-infected patients. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2019, 31, 513-518.	1.2	2
45	Differences in hypertension and prehypertension among people living with and without HIV in China: role of HIV infection and antiretroviral therapy. <i>HIV Medicine</i> , 2021, 22, 409-417.	2.2	2
46	Association between inflammation and coagulation biomarkers and carotid atherosclerosis among treated people with HIV. <i>Open Forum Infectious Diseases</i> , 0, , .	0.9	2
47	The prevalence of HSV-2 infection in HIV-1 discordant couples. <i>Epidemiology and Infection</i> , 2016, 144, 97-105.	2.1	1
48	Association of HIV infection with metabolic syndrome among normal or underweight young adults: evidence from the CHART cohort. <i>BioScience Trends</i> , 2020, 14, 450-456.	3.4	1
49	Interaction of sex and HIV infection on renal impairment: baseline evidence from the CHART cohort. <i>International Journal of Infectious Diseases</i> , 2022, 116, 182-188.	3.3	1
50	Baseline Characteristics of Mitochondrial DNA and Mutations Associated With Short-Term Posttreatment CD4+T-Cell Recovery in Chinese People With HIV. <i>Frontiers in Immunology</i> , 2021, 12, 793375.	4.8	0