
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7342375/publications.pdf Version: 2024-02-01



Μάρτα Βάτιις

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Optimal Management of Metastatic Melanoma: Current Strategies and Future Directions. American Journal of Clinical Dermatology, 2013, 14, 179-194. | 3.3 | 89 |
| 2 | Treatment of advanced squamous cell carcinoma of the lung: a review. Translational Lung Cancer Research, 2015, 4, 524-32. | 1.3 | 74 |
| 3 | Patterns of locoregional failure in stage III non-small cell lung cancer treated with definitive chemoradiation therapy. Practical Radiation Oncology, 2014, 4, 342-348. | 1.1 | 26 |
| 4 | Neutrophil-to-Lymphocyte Ratio Is a Predictive Biomarker in Patients with Epidermal Growth Factor Receptor (EGFR) Mutated Advanced Non-Small Cell Lung Cancer (NSCLC) Treated with Tyrosine Kinase Inhibitor (TKI) Therapy. Cancers, 2021, 13, 1426. | 1.7 | 17 |
| 5 | Docetaxel, bevacizumab, and gemcitabine for very high risk sarcomas in adolescents and young adults: A single-center experience. Pediatric Blood and Cancer, 2017, 64, e26265. | 0.8 | 14 |
| 6 | Neutrophil to lymphocyte ratio as predictive of prolonged progression free survival (PFS) and overall survival (OS) in patients with metastatic non-small cell lung cancer (NSCLC) treated with second-line PD-1 immune checkpoint inhibitors Journal of Clinical Oncology, 2017, 35, e14530-e14530. | 0.8 | 12 |
| 7 | Survival Prediction in Ambulatory Patients With Stage III/IV Non-Small Cell Lung Cancer Using the Palliative Performance Scale, ECOG, and Lung Cancer Symptom Scale. American Journal of Hospice and Palliative Medicine, 2016, 33, 374-380. | 0.8 | 10 |
| 8 | Pulmonary metastasectomy in bone and soft tissue sarcoma with metastasis to the lung. Interactive Cardiovascular and Thoracic Surgery, 2021, 33, 879-884. | 0.5 | 9 |
| 9 | Primary and secondary therapeutic strategies for EGF receptor pathway inhibition in non-small-cell lung cancer. Expert Review of Anticancer Therapy, 2010, 10, 1589-1599. | 1.1 | 8 |
| 10 | Differential expression of circulating biomarkers of tumor phenotype and outcomes in previously treated non-small cell lung cancer patients receiving erlotinib vs. cytotoxic chemotherapy. Oncotarget, 2017, 8, 58108-58121. | 0.8 | 7 |
| 11 | Heart dose and coronary artery calcification in patients receiving thoracic irradiation for lung cancer. Journal of Thoracic Disease, 2020, 12, 223-231. | 0.6 | 7 |
| 12 | Outcomes of brain metastasis in high-grade bone and soft tissue sarcoma: An analysis of clinicopathological characteristics and survival data. Rare Tumors, 2021, 13, 203636132110261. | 0.3 | 6 |
| 13 | Practical and theoretical implications of weight gain in advanced non-small cell lung cancer patients. Annals of Translational Medicine, 2017, 5, 152-152. | 0.7 | 6 |
| 14 | The clinical utility of next-generation sequencing for bone and soft tissue sarcoma. Acta Oncológica, 2022, 61, 38-44. | 0.8 | 6 |
| 15 | Relationship between circulating tumor-associated autoantibodies and clinical outcomes in advanced-stage NSCLC patients receiving PD-1/â°'L1 directed immune checkpoint inhibition Journal of Immunological Methods, 2021, 490, 112956. | 0.6 | 5 |
| 16 | IMpower150: Exploratory efficacy analysis in patients (pts) with bulky disease Journal of Clinical Oncology, 2020, 38, e21637-e21637. | 0.8 | 4 |
| 17 | What is the clinical impact of staging and surveillance PET T scan findings in patients with bone and soft tissue sarcoma?. Journal of Surgical Oncology, 2022, , . | 0.8 | 3 |
| 18 | Potential predictive value of hepatocyte growth factor (HGF) in advanced non-small cell lung cancer (NSCLC) treated with a platinum doublet and bevacizumab Journal of Clinical Oncology, 2014, 32, e22000-e22000. | 0.8 | 2 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Clinical benefit of next generation sequencing in soft tissue and bone sarcoma: Rush University Medical Center's experience Journal of Clinical Oncology, 2019, 37, e22552-e22552. | 0.8 | 2 |
| 20 | Does synovial sarcoma grade predict oncologic outcomes, and does a lowâ€grade variant exist?. Journal of Surgical Oncology, 2022, , . | 0.8 | 2 |
| 21 | Associations between longitudinal pretreatment BMI and neutrophil/lymphocyte ratio(NLR) and progression-free(PFS) and overall survival(OS) in advanced NSCLC patients treated with single agent anti-PD-1/anti-PDL1 monoclonal antibodies(mAbs) Journal of Clinical Oncology, 2021, 39, e21189-e21189. | 0.8 | 1 |
| 22 | Association of KRAS mutations detected via liquid biopsy in metastatic non-small cell lung cancer patients with high levels of FDG-PET SUV Journal of Clinical Oncology, 2017, 35, e20594-e20594. | 0.8 | 1 |
| 23 | Association of baseline and longitudinal low neutrophil-lymphocyte ratio (NLR) and high lymphocyte counts (LCs) with progression-free survival (PFS) and overall survival (OS) in real world advanced non-small cell lung cancer (aNSCLC) patients (pts) treated with nivolumab (nivo) or pembrolizumab (pembro) lournal of Clinical Oncology, 2018, 36, e21020-e21020. | 0.8 | 1 |
| 24 | Associations between baseline serum biomarker levels and cachexia/precachexia in pretreated non-small cell lung cancer (NSCLC) patients Journal of Clinical Oncology, 2019, 37, 3054-3054. | 0.8 | 1 |
| 25 | Differences in circulating angiogenic biomarkers as prognosticator for outcome in bevacizumab-treated nonsquamous non-small cell lung cancer (NSCLC) patients Journal of Clinical Oncology, 2014, 32, 11037-11037. | 0.8 | 1 |
| 26 | Real-world (rw) clinical outcomes for advanced/metastatic non-small cell lung cancer (aNSCLC) patients (pts) treated with second line (2L) ramucirumab plus docetaxel (R+D) post frontline (1L) platinum based chemotherapy plus immune checkpoint inhibitors (Pt + ICI) Journal of Clinical Oncology, 2019, 37, e20727-e20727. | 0.8 | 1 |
| 27 | Real-world tumor response (rwTR) to ramucirumab plus docetaxel (R+D) post platinum-based (Pt) and immune checkpoint inhibitor (ICI) therapy in advanced non-small cell lung cancer (aNSCLC) patients (pts) Journal of Clinical Oncology, 2019, 37, e20725-e20725. | 0.8 | 1 |
| 28 | The prognostic value of neutrophil-to-lymphocyte ratio in patients with epidermal growth factor receptor mutated advanced non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2020, 38, e21675-e21675. | 0.8 | 1 |
| 29 | Psychosocial distress in patients with driver-mutant lung cancer Journal of Clinical Oncology, 2021, 39, e24125-e24125. | 0.8 | 0 |
| 30 | A Pilot Study To Test the Efficacy of a Combination of Gleevec with Thalidomide in Patients with Idiopathic Primary Myelofibrosis, Myelofibrosis with Myeloid Metaplasia and Myelodysplastic Syndromes Who Present with Myelofibrosis Blood, 2006, 108, 4841-4841. | 0.6 | 0 |
| 31 | Hematopoietic Cell Transplantation - Comorbidity Index (HCT-CI) as a Surrgoate Marker for Predition of Transplant Outcome for Pateints (pts) with Multiple Myeloma (MM) Undergoing Autologous Stem Cell Transplantation (ASCT): High HCT-CI Score Predicts for Poor Outcome Blood, 2007, 110, 5104-5104. | 0.6 | 0 |
| 32 | Autologous (Auto) and Allogeneic (Allo) Stem Cell Transplantation (SCT) for Pateints (pts) with Advanced Chronic Lymphoctic Leukemia (CLL): A Comparative Study Blood, 2007, 110, 5044-5044. | 0.6 | 0 |
| 33 | Preoperative survivin, ERCC1, and PTEN expression in stage III non-small cell lung cancer (NSCLC) patients (pts) treated with neoadjuvant and definitive chemoradiation and association with overall survival (OS) Journal of Clinical Oncology, 2012, 30, 7067-7067. | 0.8 | 0 |
| 34 | EGFR gene mutation and epithelial to mensenchymal transition (EMT) markers in advanced NSCLC patients treated with erlotinib Journal of Clinical Oncology, 2012, 30, e18117-e18117. | 0.8 | 0 |
| 35 | Thyroid transcription factor 1 (TTF-1) and overall survival in wild type EGFR patients treated with erlotinib Journal of Clinical Oncology, 2013, 31, e19113-e19113. | 0.8 | 0 |
| 36 | Epithelial to mesenchymal markers and clinical outcomes on erlotinib in stage IV non-small cell lung cancer patients Journal of Clinical Oncology, 2013, 31, e19117-e19117. | 0.8 | 0 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Association of overall survival with weight gain during treatment, initial body mass index (BMI), and glucose intolerance (GI) in advanced non-small cell lung cancer (NSCLC) patients Journal of Clinical Oncology, 2014, 32, e19033-e19033. | 0.8 | 0 |
| 38 | Serial neutrophil to lymphocyte ratios (NLR) in relation to weight change and overall survival in patients (Pts) with metastatic non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2015, 33, e19143-e19143. | 0.8 | 0 |
| 39 | Association of biomarkers of insulin resistance and inflammation with PET CT SUV values in stage IV non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2016, 34, e20589-e20589. | 0.8 | 0 |
| 40 | KRAS and P53 mutations and FDG-PET as a measure of glycolytic activity in metastatic non-small cell lung cancer patients Journal of Clinical Oncology, 2016, 34, 9086-9086. | 0.8 | 0 |
| 41 | Two year progression free survival in stage IV NSCLC patients treated with pemetrexed continuation maintenance Journal of Clinical Oncology, 2016, 34, e20654-e20654. | 0.8 | 0 |
| 42 | Association of biomarkers of insulin resistance and inflammation with skeletal muscle index (SMI) in stage IV non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2016, 34, 10013-10013. | 0.8 | 0 |
| 43 | Association of acute phase proteins with a serum proteomic test in previously treated patients with NSCLC Journal of Clinical Oncology, 2016, 34, e20649-e20649. | 0.8 | 0 |
| 44 | Circulating biomarkers and outcomes in advanced non-small cell lung cancer patients treated with anti-PD1 (program death 1 receptor) monoclonal antibodies Journal of Clinical Oncology, 2017, 35, e20592-e20592. | 0.8 | 0 |
| 45 | Mature progression-free survival (PFS) milestones in real world stage IV, non-squamous, non-small cell lung cancer patients (nsqNSCLC) treated with first line pemetrexed(Pem)/platinum(Plat) +/-bevacizumab(Bev) followed by pem +/-bev maintenance Journal of Clinical Oncology, 2018, 36, e21063-e21063. | 0.8 | 0 |
| 46 | Ramucirumab (ram), immune checkpoint inhibitors (ICIs), and single-agent chemotherapy (chemo) usage in real-world advanced non-small cell lung cancer (aNSCLC) patients (pts) after rapid disease progression (RDP) on platinum (Pt) Journal of Clinical Oncology, 2018, 36, e21162-e21162. | 0.8 | 0 |
| 47 | Sequencing of ramucirumab (ram) and immune checkpoint inhibitors (ICIs) in platinum (Pt)-treated real-world patients (pts) with advanced non-small cell lung cancer (aNSCLC) Journal of Clinical Oncology, 2018, 36, e21209-e21209. | 0.8 | 0 |
| 48 | Prognostic value of neutrophil-to-lymphocyte ratio (NLR), serum albumin and sequence of immunotherapy (Immuno.) on overall survival (OS), and progression free survival (PFS) in patients with metastatic non-small cell lung cancer (NSCLC) treated with ramucirumab plus docetaxel (RD) Journal of Clinical Oncology, 2018, 36, e21163-e21163. | 0.8 | 0 |
| 49 | Real-world economic burden of rapid disease progression (RDP) in patients (pts) with advanced/metastatic non-small cell lung cancer (aNSCLC) Journal of Clinical Oncology, 2019, 37, e20716. | 0.8 | 0 |
| 50 | Treatment of soft tissue sarcoma with a novel cold plasma jet Journal of Clinical Oncology, 2019, 37, e22562-e22562. | 0.8 | 0 |
| 51 | Real-world clinical burden of aggressive disease (AD) in advanced/metastatic non small cell lung cancer (aNSCLC) Journal of Clinical Oncology, 2019, 37, e20723-e20723. | 0.8 | 0 |
| 52 | Correlating the effects of VeriStrat result, anti-PD1 therapy, and neutrophil-to-lymphocyte ratio (NLR) on progression-free survival (PFS) in patients with stage IV non-small cell lung cancer (NSCLC) Journal of Clinical Oncology, 2019, 37, e20660-e20660. | 0.8 | 0 |
| 53 | Real-world characterization of advanced/metastatic non-small cell lung cancer (aNSCLC) patients (pts) with rapid disease progression (RDP) Journal of Clinical Oncology, 2019, 37, e20706-e20706. | 0.8 | 0 |
| 54 | Decreasing BMI/weight immediately prior to starting anti-PD-1/PDL-1 monoclonal antibodies for treatment for stage IV non-small cell lung cancer is associated with shorter progression-free survival Journal of Clinical Oncology, 2019, 37, e20710-e20710. | 0.8 | 0 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Mature, real world progression-free survival (PFS) and overall survival (OS) milestones in stage IV, non-squamous, non-small cell lung cancer patients (nsqNSCLC) treated with first line pemetrexed(Pem)/platinum(Plat) followed by pem+/-bevacizumab(Bev) maintenance Journal of Clinical Oncology, 2019, 37, e20721-e20721. | 0.8 | 0 |
| 56 | Changes in skeletal muscle mass during PD-1 and PD-L1 checkpoint inhibitor therapy in advanced-stage non-small cell lung cancer patients Journal of Clinical Oncology, 2019, 37, e14061-e14061. | 0.8 | 0 |
| 57 | RNA pathway enrichment in serum-based mass spectroscopy prognostic analyses Journal of Clinical Oncology, 2020, 38, e15197-e15197. | 0.8 | 0 |
| 58 | PET-CT staging affects time to treatment in sarcoma. Surgical Oncology, 2022, 41, 101732. | 0.8 | 0 |
| 59 | Gender and weight change, skeletal muscle index(SMI) change, and survival in advanced non-small cell lung cancer(NSCLC) patients(pts.) receiving platinum chemotherapy Journal of Clinical Oncology, 2022, 40, e24080-e24080. | 0.8 | 0 |
| 60 | NRG-DT001 phase lb trial of neoadjuvant navtemadlin (previously AMG232 and KRT232) concurrent with preoperative radiotherapy in wild-type p53 soft tissue sarcoma of the extremity and body wall Journal of Clinical Oncology, 2022, 40, 11521-11521. | 0.8 | 0 |
| 61 | Clinical and laboratory parameters associated with rapid progression in advanced NSCLC patients treated with second or third-line single agent immune checkpoint inhibitors (ICIs) Journal of Clinical Oncology, 2022, 40, e21087-e21087. | 0.8 | 0 |