
Jingyi Chen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7353667/publications.pdf Version: 2024-02-01



LINCYL CHEN

#	Article	IF	CITATIONS
1	Dexmedetomidine Preconditioning Protects Cardiomyocytes Against Hypoxia/Reoxygenation-Induced Necroptosis by Inhibiting HMGB1-Mediated Inflammation. Cardiovascular Drugs and Therapy, 2019, 33, 45-54.	1.3	35
2	LncRNA MALAT1 sponges miR-203 to promote inflammation in myocardial ischemia-reperfusion injury. International Journal of Cardiology, 2018, 268, 245.	0.8	29
3	LncRNA MALAT1 Promotes Oxygen-Glucose Deprivation and Reoxygenation Induced Cardiomyocytes Injury Through Sponging miR-20b to Enhance beclin1-Mediated Autophagy. Cardiovascular Drugs and Therapy, 2019, 33, 675-686.	1.3	23
4	Circular RNA DLGAP4 ameliorates cardiomyocyte apoptosis through regulating BCL2 via targeting miR-143 in myocardial ischemia-reperfusion injury. International Journal of Cardiology, 2019, 279, 147.	0.8	23
5	MALAT1/miR-204/LC3-II: A potential regulated axis of autophagy in myocardial ischemia-reperfusion injury. International Journal of Cardiology, 2019, 277, 222.	0.8	10
6	TUG1/miR-421/PINK1: A potential mechanism for treating myocardial ischemia-reperfusion injury. International Journal of Cardiology, 2019, 292, 197.	0.8	8
7	Circulating HOTAIR/miR-126 axis is negatively associated with disease risk of incident myocardial infarction. International Journal of Cardiology, 2020, 298, 121.	0.8	5
8	MALAT1/miR-144/Brg1: A potential regulated axis of inflammation in myocardial ischemia-reperfusion injury. International Journal of Cardiology, 2019, 283, 151.	0.8	4
9	The circulating LncRNA SNHG15/miR-346 axis may be a potential biomarker of cardiomyocyte apoptosis during myocardial ischemia/reperfusion injury. International Journal of Cardiology, 2021, 334, 30.	0.8	4
10	MEG3/miR-223: A potentially reliable risk factor predictor for myocardial ischemia-reperfusion injury. International Journal of Cardiology, 2019, 293, 259.	0.8	3
11	LncRNA XIST may exert a profibrotic role via sponging miR-133a through SOCS2-activated autophagy in myocardial infarction. International Journal of Cardiology, 2021, 337, 100.	0.8	3