## Chen Jianhuai

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7367066/publications.pdf Version: 2024-02-01



<u>CHEN ΙΙΛΝΗΙΙΛΙ</u>

#	Article	IF	CITATIONS
1	Upregulated expression of NMDA receptor in the paraventricular nucleus shortens ejaculation latency in rats with experimental autoimmune prostatitis. Andrology, 2021, 9, 352-360.	1.9	4
2	Impaired frontalâ€parietal control network in chronic prostatitis/chronic pelvic pain syndrome revealed by graph theoretical analysis: A DTI study. European Journal of Neuroscience, 2021, 53, 1060-1071.	1.2	16
3	Aberrant default mode network and auditory network underlying the sympathetic skin response of the penis (PSSR) of patients with premature ejaculation: A restingâ€state fMRI study. Andrology, 2021, 9, 277-287.	1.9	6
4	Graph theory analysis reveals premature ejaculation is a brain disorder with altered structural connectivity and depressive symptom: A DTIâ€based connectome study. European Journal of Neuroscience, 2021, 53, 1905-1921.	1.2	11
5	Abnormal Functional Connectivity Between the Left Medial Superior Frontal Gyrus and Amygdala Underlying Abnormal Emotion and Premature Ejaculation: A Resting State fMRI Study. Frontiers in Neuroscience, 2021, 15, 704920.	1.4	4
6	Increased attentional network activity in premature ejaculation patients with anxiety revealed by restingâ€state functional magnetic resonance imaging. European Journal of Neuroscience, 2021, 54, 5417-5426.	1.2	4
7	Altered Structural and Functional Connectivity Contribute to Rapid Ejaculation: Insights from a Multimodal Neuroimaging Study. Neuroscience, 2021, 471, 93-101.	1.1	1
8	Disrupted topological properties of brain networks in erectile dysfunction patients owing predominantly to psychological factors: a structural and functional neuroimaging study. Andrology, 2020, 8, 381-391.	1.9	15
9	Graph analysis of <scp>DTI</scp> â€based connectome: decreased local efficiency of subcortical regions in <scp>PE</scp> patients with high sympathetic activity. Andrology, 2020, 8, 400-406.	1.9	18
10	Reduced segregation and integration of structural brain network associated with sympathetic and dorsal penile nerve activity in anejaculation patients: a graphâ€based connectome study. Andrology, 2020, 8, 392-399.	1.9	9
11	Variation in Brain Subcortical Network Topology Between Men with and Without PE: A Diffusion Tensor Imaging Study. Journal of Sexual Medicine, 2020, 17, 48-59.	0.3	13
12	Dopamine D2 receptors in the basolateral amygdala modulate erectile function in a rat model of nonorganic erectile dysfunction. Andrologia, 2019, 51, e13160.	1.0	13
13	Changes in Male Rat Sexual Behavior and Brain Activity Revealed by Functional Magnetic Resonance Imaging in Response to Chronic Mild Stress. Journal of Sexual Medicine, 2018, 15, 136-147.	0.3	18
14	Brain structural network topological alterations of the left prefrontal and limbic cortex in psychogenic erectile dysfunction. International Journal of Neuroscience, 2018, 128, 393-403.	0.8	23
15	Impaired Prefrontal-Amygdala Pathway, Self-Reported Emotion, and Erection in Psychogenic Erectile Dysfunction Patients With Normal Nocturnal Erection. Frontiers in Human Neuroscience, 2018, 12, 157.	1.0	22
16	Altered brain networks in psychogenic erectile dysfunction: a restingâ€state <scp>fMRI</scp> study. Andrology, 2017, 5, 1073-1081.	1.9	23
17	Altered anatomical patterns of depression in relation to antidepressant treatment: Evidence from a pattern recognition analysis on the topological organization of brain networks. Journal of Affective Disorders, 2015, 180, 129-137.	2.0	46
18	Abnormal hubs of white matter networks in the frontal-parieto circuit contribute to depression discrimination via pattern classification. Magnetic Resonance Imaging, 2014, 32, 1314-1320.	1.0	27