

# Hiroaki Hagiwara

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7370596/publications.pdf>

Version: 2024-02-01

16

papers

452

citations

933447

10

h-index

940533

16

g-index

16

all docs

16

docs citations

16

times ranked

519

citing authors

#	ARTICLE	IF	CITATIONS
1	Thermosalience coupled to abrupt spin crossover with dynamic ligand motion in an iron( $\text{L}_2\text{-}2\text{Ph}$ ) molecular crystal. <i>CrystEngComm</i> , 2022, 24, 4224-4234.	2.6	4
2	Alternative Route Triggering Multistep Spin Crossover with Hysteresis in an Iron(II) Family Mediated by Flexible Anion Ordering. <i>Inorganic Chemistry</i> , 2020, 59, 9866-9880.	4.0	10
3	Jumping Crystals of Stacked Planar Cobalt Complexes: Thermosalient Effect Promoted by Hydrogen-bonded Lattice Solvent Release. <i>Chemistry Letters</i> , 2019, 48, 1077-1080.	1.3	6
4	Iron(II) Spin Crossover Complex with the 1,2,3-Triazole-Containing Linear Pentadentate Schiff-Base Ligand and the MeCN Monodentate Ligand. <i>Crystals</i> , 2019, 9, 276.	2.2	4
5	High-temperature Spin Crossover of a Solvent-Free Iron(II) Complex with the Linear Hexadentate Ligand $[\text{Fe}(\text{L}_2\text{-}2\text{Ph})](\text{AsF}_6)_2$ ( $\text{L}_2\text{-}2\text{Ph}$ = bis[N-(1-Phenyl-1H-1,2,3-triazol-4-yl)methylidene-2-aminoethyl]-1,3-) Tj ETQq1 1 0.784314	1.0	14
6	Neutral Molecular Iron(II) Complexes Showing Tunable Bistability at Above, Below, and Just Room Temperature by a Crystal Engineering Approach: Ligand Mobility into a Three-Dimensional Flexible Supramolecular Network. <i>Crystal Growth and Design</i> , 2017, 17, 6006-6019.	3.0	26
7	High-Temperature Wide Thermal Hysteresis of an Iron(II) Dinuclear Double Helicate. <i>Inorganics</i> , 2017, 5, 49.	2.7	15
8	Synthesis, structure, and spin crossover above room temperature of a mononuclear and related dinuclear double helicate iron( $\text{L}_2\text{-}2\text{Ph}$ ) complexes. <i>Dalton Transactions</i> , 2016, 45, 17132-17140.	3.3	33
9	A polymorphism-dependent $T_{1/2}$ shift of 100 K in a hysteretic spin-crossover complex related to differences in intermolecular weak $\text{CH}\cdots\text{X}$ hydrogen bonds ( $\text{X} = \text{S}$ vs. $\text{S}$ and $\text{N}$ ). <i>Chemical Communications</i> , 2016, 52, 815-818.	4.1	41
10	Synthesis, Structure, and Magnetic Property of a New Mononuclear Iron(II) Spin Crossover Complex with a Tripodal Ligand Containing Three 1,2,3-Triazole Groups. <i>Chemistry Letters</i> , 2014, 43, 950-952.	1.3	21
11	Synthesis, structures, and magnetic properties of iron(II) complexes, $[\text{Fe}(\text{HLMe})_2](\text{ClO}_4)_2$ and its ethanol adduct $[\text{Fe}(\text{HLMe})_2](\text{ClO}_4)_2\cdot\text{EtOH}$ ( $\text{HLMe}$ =2-methylimidazol-4-yl-methylideneamino-2-ethylpyridine): Their structural distortion and spin states. <i>Polyhedron</i> , 2012, 48, 110-116.	2.2	2
12	Layered iron(II) spin crossover complex constructed by $\text{NH}\cdots\text{Br}^\bullet$ hydrogen bonds with 2K wide thermal hysteresis, $[\text{Fe}(\text{H3LMe})\text{Br}\cdot\text{CF}_3\text{SO}_3]$ ( $\text{H3LMe}$ =tris[2-((2-methylimidazol-4-yl)methylidene)amino)ethyl]amine). <i>Inorganica Chimica Acta</i> , 2011, 366, 283-289.	2.4	24
13	Conformational effect of a spin crossover iron(II) complex: Bis[N-(2-methylimidazol-4-yl)methylidene-2-aminoethyl]propanediamineiron(II) perchlorate. <i>Inorganica Chimica Acta</i> , 2011, 367, 141-150.	2.4	12
14	Two-Dimensional Iron(II) Spin Crossover Complex Constructed of Bifurcated $\text{NH}\cdots\text{O}$ -Hydrogen Bonds and $\text{C}\cdots\text{O}$ Interactions: $[\text{Fe}(\text{HLH},\text{Me})_2](\text{ClO}_4)_2\cdot1.5\text{MeCN}$ ( $\text{HLH},\text{Me}$ ) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 2174Td (Imidazol-4-yl-methylideneamino-2-ethylpyridine)	1.0	10
15	Structural-Electronic Correlation in the First-Order Phase Transition of $[\text{FeH}_2\text{L}_2\text{-Me}](\text{ClO}_4)_2$ ( $\text{H}_2\text{L}_2\text{-Me}$ = Bis[((2-methylimidazol-4-yl)methylidene)-3-aminopropyl]ethylenediamine). <i>Inorganic Chemistry</i> , 2006, 45, 8126-8135.	4.0	43
16	A Variety of Spin-Crossover Behaviors Depending on the Counter Anion: Two-Dimensional Complexes Constructed by $\text{NH}\cdots\text{Cl}^\bullet$ Hydrogen Bonds, $[\text{Fe}(\text{H3LMe})\text{Cl}\cdots\text{X}]$ ( $\text{X} = \text{PF}_6^\bullet$ , $\text{AsF}_6^\bullet$ , $\text{SbF}_6^\bullet$ , $\text{CF}_3\text{SO}_3^\bullet$ ; $\text{H}_3\text{LMe}$ =) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 2174Td (Imidazol-4-yl-methylideneamino-2-ethylpyridine)	1.0	160
	12, 4536-4549.		