
Georgios Nikolakaros

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7371424/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Recurrent depression in childhood and adolescence and low childhood socioeconomic status predict low cardiorespiratory fitness in early adulthood. Journal of Affective Disorders, 2020, 266, 782-792.	2.0	7
2	<p>A patient with Korsakoff syndrome of psychiatric and alcoholic etiology presenting as DSM-5 mild neurocognitive disorder</p> . Neuropsychiatric Disease and Treatment, 2019, Volume 15, 1311-1320.	1.0	2
3	Korsakoff Syndrome in Non-alcoholic Psychiatric Patients. Variable Cognitive Presentation and Impaired Frontotemporal Connectivity. Frontiers in Psychiatry, 2018, 9, 204.	1.3	4
4	Obesity, Underweight, and Smoking Are Associated with Worse Cardiorespiratory Fitness in Finnish Healthy Young Men: A Population-Based Study. Frontiers in Public Health, 2017, 5, 206.	1.3	12
5	Non-alcoholic Korsakoff syndrome in psychiatric patients with a history of undiagnosed Wernicke's encephalopathy. Journal of the Neurological Sciences, 2016, 370, 296-302.	0.3	15
6	Obstructive sleep apnea in psychiatric outpatients. A clinic-based study. Journal of Psychiatric Research, 2015, 69, 126-134.	1.5	8
7	Factors associated with reported childhood depressive symptoms at age 8 and later self-reported depressive symptoms among boys at age 18. Social Psychiatry and Psychiatric Epidemiology, 2011, 46, 207-218.	1.6	11
8	Child Mental Health Problems and Obesity in Early Adulthood. Journal of Pediatrics, 2010, 156, 93-97.	0.9	47
9	Life events, self-reported psychopathology and sense of coherence among young men—A population-based study. Nordic Journal of Psychiatry, 2008, 62, 464-471.	0.7	25
10	Childhood Predictors of Military Fitness: A Prospective, Community-Based, Follow-up Study from Age 8 to Age 18. Military Medicine, 2008, 173, 146-154.	0.4	9
11	Recognition of psychiatric disorders, and self-perceived problems. A follow-up study from age 8 to age 18. Journal of Child Psychology and Psychiatry and Allied Disciplines, 2005, 46, 1124-1134.	3.1	39
12	Factors associated with ideation and acts of deliberate self-harm among 18-year-old boys. Social Psychiatry and Psychiatric Epidemiology, 2005, 40, 912-921.	1.6	72
13	Sense of coherence among Finnish young men – A cross-sectional study at military call-up. Nordic Journal of Psychiatry, 2005, 59, 473-480.	0.7	17
14	Childhood Predictors of Psychiatric Disorders Among Boys: A Prospective Community-Based Follow-up Study From Age 8 Years to Early Adulthood. Journal of the American Academy of Child and Adolescent Psychiatry, 2005, 44, 756-767.	0.3	89
15	Factors associated with depressive symptoms among 18-year-old boys: a prospective 10-year follow-up study. Journal of Affective Disorders, 2004, 83, 143-154.	2.0	48
16	Mental Health Service Use Among 18-Year-Old Adolescent Boys: A Prospective 10-Year Follow-up Study. Journal of the American Academy of Child and Adolescent Psychiatry, 2004, 43, 1250-1258.	0.3	60