

Mitsuhiko Noda

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7378634/mitsuhiko-noda-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

280
papers

9,173
citations

47
h-index

86
g-index

297
ext. papers

10,602
ext. citations

4.3
avg, IF

5.88
L-index

#	Paper	IF	Citations
280	Employment status and diabetic outpatient appointment non-attendance in middle to senior working generation with type 2 diabetes: the Japan diabetes outcome intervention trial-2 large-scale trial 005 (J-DOIT2-LT005).. <i>Acta Diabetologica</i> , 2022 , 1	3.9	
279	Dietary intake and physical activity in Japanese patients with type 2 diabetes: the Japan Diabetes Complication and its Prevention prospective study (JDCP study 8).. <i>Diabetology International</i> , 2022 , 13, 344-357	2.3	
278	Cross-sectional associations between the types/amounts of beverages consumed and the glycemia status: The Japan Public Health Center-based Prospective Diabetes study.. <i>Metabolism Open</i> , 2022 , 14, 100185	2.8	
277	Higher Serum Soluble TREM2 as a Potential Indicative Biomarker for Cognitive Impairment in Inadequately Controlled Type 2 Diabetes Without Obesity: The DOR-KyotoJ-1.. <i>Frontiers in Endocrinology</i> , 2022 , 13, 880148	5.7	0
276	Differences in the birthweight of infants born to patients with early- or mid-to-late-detected gestational diabetes mellitus who underwent guideline-based glycemetic control. <i>Journal of Diabetes and Its Complications</i> , 2021 , 35, 107850	3.2	2
275	Impact of telephone support programme using telemonitoring on stage of change towards healthy eating and active exercise in people with prediabetes. <i>Journal of Telemedicine and Telecare</i> , 2021 , 27, 307-313	6.8	1
274	Dietary glycemic index, glycemic load and mortality: Japan Public Health Center-based prospective study. <i>European Journal of Nutrition</i> , 2021 , 60, 4607-4620	5.2	
273	Low carbohydrate diet and all cause and cause-specific mortality. <i>Clinical Nutrition</i> , 2021 , 40, 2016-2024	5.9	5
272	Effective coverage of medical treatment for hypertension, diabetes and dyslipidaemia in Japan: An analysis of National Health and Nutrition Surveys 2003-2017. <i>Journal of Health Services Research and Policy</i> , 2021 , 26, 106-114	2.4	1
271	Multifactorial intervention has a significant effect on diabetic kidney disease in patients with type 2 diabetes. <i>Kidney International</i> , 2021 , 99, 256-266	9.9	14
270	Effect of a Multifactorial Intervention on Fracture in Patients With Type 2 Diabetes: Subanalysis of the J-DOIT3 Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 , 106, e2116-e2128	5.6	0
269	Characteristics Associated with Early Worsening of Retinopathy in Patients with Type 2 Diabetes Diagnosed with Retinopathy at Their First Visit: A Retrospective Observational Study. <i>Journal of Diabetes Research</i> , 2021 , 2021, 7572326	3.9	0
268	Association of sugary drink consumption with all-cause and cause-specific mortality: the Japan Public Health Center-based Prospective Study. <i>Preventive Medicine</i> , 2021 , 148, 106561	4.3	2
267	Synergistic association of the copper/zinc ratio under inflammatory conditions with diabetic kidney disease in patients with type 2 diabetes: The Asahi Diabetes Complications Study. <i>Journal of Diabetes Investigation</i> , 2021 ,	3.9	1
266	A Comparison of the Association of Fasting Plasma Glucose and HbA1c Levels with Diabetic Retinopathy in Japanese Men. <i>Journal of Diabetes Research</i> , 2020 , 2020, 3214676	3.9	0
265	Body mass index and mortality among middle-aged Japanese individuals with diagnosed diabetes: The Japan Public Health Center-based prospective study (JPHC study). <i>Diabetes Research and Clinical Practice</i> , 2020 , 164, 108198	7.4	
264	Effects of obesity, metabolic syndrome, and non-alcoholic or alcoholic elevated liver enzymes on incidence of diabetes following lifestyle intervention: A subanalysis of the J-DOIT1. <i>Journal of Occupational Health</i> , 2020 , 62, e12109	2.3	0

263	A cluster-randomized trial of the effectiveness of a triple-faceted intervention promoting adherence to primary care physician visits by diabetes patients. <i>Scientific Reports</i> , 2020 , 10, 2842	4.9	1
262	Association of soy and fermented soy product intake with total and cause specific mortality: prospective cohort study. <i>BMJ, The</i> , 2020 , 368, m34	5.9	28
261	Dietary fiber intake and total and cause-specific mortality: the Japan Public Health Center-based prospective study. <i>American Journal of Clinical Nutrition</i> , 2020 , 111, 1027-1035	7	13
260	Clinical characterization of patients with primary aldosteronism plus subclinical Cushing@ syndrome. <i>BMC Endocrine Disorders</i> , 2020 , 20, 9	3.3	3
259	Prevalence of diabetes in Japanese patients with cancer. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1159-1162	3.9	162
258	Passive smoking and type2 diabetes among never-smoking women: The Japan Public Health Center-based Prospective Study. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1352-1358	3.9	2
257	Intensity-specific validity and reliability of the Japan Public Health Center-based prospective study-physical activity questionnaire. <i>Preventive Medicine Reports</i> , 2020 , 20, 101169	2.6	3
256	Japanese Clinical Practice Guideline for Diabetes 2019. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1020-1076	3.9	46
255	Japanese Clinical Practice Guideline for Diabetes 2019. <i>Diabetology International</i> , 2020 , 11, 165-223	2.3	75
254	Self-weighing frequency and the incidence of type 2 diabetes: post hoc analysis of a cluster-randomized controlled trial. <i>BMC Research Notes</i> , 2020 , 13, 375	2.3	0
253	Risk perception, self-efficacy, trust for physician, depression, and behavior modification in diabetic patients. <i>Journal of Health Psychology</i> , 2020 , 25, 350-360	3.1	9
252	Association Between Okinawan Vegetables Consumption and Risk of Type 2 Diabetes in Japanese Communities: The JPHC Study. <i>Journal of Epidemiology</i> , 2020 , 30, 227-235	3.4	1
251	Serum 25-hydroxyvitamin D and risk of type 2 diabetes among Japanese adults: the Hitachi Health Study. <i>Clinical Nutrition</i> , 2020 , 39, 1218-1224	5.9	5
250	Diabetes and cancer risk: A Mendelian randomization study. <i>International Journal of Cancer</i> , 2020 , 146, 712-719	7.5	14
249	Soy food and isoflavones are not associated with changes in serum lipids and glycohemoglobin concentrations among Japanese adults: a cohort study. <i>European Journal of Nutrition</i> , 2020 , 59, 2075-2087	5.2	4
248	Bodyweight threshold for sudden onset of ketosis might exist in ketosis-prone type2 diabetes patients. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 499-501	3.9	1
247	Detailed Time Course of Decline in Serum C-Peptide Levels in Anti-Programmed Cell Death-1 Therapy-Induced Fulminant Type 1 Diabetes. <i>Diabetes Care</i> , 2019 , 42, e40-e41	14.6	7
246	Recommended configuration for personal health records by standardized data item sets for diabetes mellitus and associated chronic diseases: A report from Collaborative Initiative by six Japanese Associations. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 868-875	3.9	3

245	Prevention of Metabolic Syndrome by Telephone-Delivered Lifestyle Intervention in a Real-World Setting: Sub-Analysis of a Cluster-Randomized Trial. <i>Metabolic Syndrome and Related Disorders</i> , 2019 , 17, 355-361	2.6	1
244	Serum amino acid profiles and risk of type 2 diabetes among Japanese adults in the Hitachi Health Study. <i>Scientific Reports</i> , 2019 , 9, 7010	4.9	31
243	Clinical characteristics of anti-glutamic acid decarboxylase antibody-positive fulminant type 1 diabetes. <i>Endocrine Journal</i> , 2019 , 66, 329-336	2.9	3
242	Menstrual and reproductive factors and type 2 diabetes risk: The Japan Public Health Center-based Prospective Study. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 147-153	3.9	9
241	Significance of peripheral mononuclear cells producing interferon- γ in response to insulin B:9-23-related peptides in subtypes of type 1 diabetes. <i>Clinical Immunology</i> , 2019 , 208, 108260	9	1
240	Plasma C-peptide and glycated albumin and subsequent risk of cancer: From a large prospective case-cohort study in Japan. <i>International Journal of Cancer</i> , 2019 , 144, 718-729	7.5	2
239	Weight control before and during pregnancy for patients with gestational diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 1075-1082	3.9	6
238	Being underweight in adolescence is independently associated with adult-onset diabetes among women: The Japan NursesQHealth Study. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 827-836	3.9	6
237	Urinary afamin levels are associated with the progression of diabetic nephropathy. <i>Diabetes Research and Clinical Practice</i> , 2019 , 147, 37-46	7.4	8
236	Cruciferous vegetable intake and mortality in middle-aged adults: A prospective cohort study. <i>Clinical Nutrition</i> , 2019 , 38, 631-643	5.9	11
235	Intrinsic insulin secretion capacity might be preserved by discontinuing anti-programmed cell death protein 1 antibody treatment in Qnti-programmed cell death protein 1 antibody-inducedQfulminant type 1 diabetes. <i>Journal of Diabetes Investigation</i> , 2018 , 9, 448-449	3.9	8
234	Changes in Antidiabetic Drug Prescription and Glycemic Control Trends in Elderly Patients with Type 2 Diabetes Mellitus from 2005-2013: An Analysis of the National Center Diabetes Database (NCDD-03). <i>Internal Medicine</i> , 2018 , 57, 1229-1240	1.1	14
233	Japanese Clinical Practice Guideline for Diabetes 2016. <i>Journal of Diabetes Investigation</i> , 2018 , 9, 657	3.9	136
232	Circulating odd-chain saturated fatty acids were associated with arteriosclerosis among patients with diabetes, dyslipidemia, or hypertension in Sri Lanka but not Japan. <i>Nutrition Research</i> , 2018 , 50, 82-93	4	6
231	Administration of thiamazole for GravesQisease might trigger the onset of type 1 diabetes. <i>Journal of Diabetes Investigation</i> , 2018 , 9, 1228-1229	3.9	0
230	Japanese Clinical Practice Guideline for Diabetes 2016. <i>Diabetology International</i> , 2018 , 9, 1-45	2.3	114
229	Evaluation of Tenueligliptin Effects on Transcriptional Activity of PPAR α in Cell-Based Assays. <i>Journal of Nippon Medical School</i> , 2018 , 85, 95-101	1.3	
228	Long-Term Effects of Ipragliflozin on Diabetic Nephropathy and Blood Pressure in Patients With Type 2 Diabetes: 104-Week Follow-up of an Open-Label Study. <i>Journal of Clinical Medicine Research</i> , 2018 , 10, 679-687	2.9	5

227	Effects of Coffee and Tea Consumption on Glucose Metabolism: A Systematic Review and Network Meta-Analysis. <i>Nutrients</i> , 2018 , 11,	6.7	26
226	Coffee Consumption and Lung Cancer Risk: The Japan Public Health Center-Based Prospective Study. <i>Journal of Epidemiology</i> , 2018 , 28, 207-213	3.4	7
225	Luminal plant sterol promotes brush border membrane-to-lumen cholesterol efflux in the small intestine. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2018 , 63, 102-105	3.1	8
224	Elevated haemoglobin A1c but not fasting plasma glucose conveys risk of chronic kidney disease in non-diabetic individuals. <i>Diabetes Research and Clinical Practice</i> , 2018 , 146, 233-239	7.4	6
223	Adult height and all-cause and cause-specific mortality in the Japan Public Health Center-based Prospective Study (JPHC). <i>PLoS ONE</i> , 2018 , 13, e0197164	3.7	10
222	The annual rate of coronary artery calcification with combination therapy with a PCSK9 inhibitor and a statin is lower than that with statin monotherapy. <i>Npj Aging and Mechanisms of Disease</i> , 2018 , 4, 7	5.5	21
221	Case of type 1 diabetes associated with less-dose nivolumab therapy in a melanoma patient. <i>Journal of Dermatology</i> , 2017 , 44, 605-606	1.6	37
220	Circulating ferritin concentrations and risk of type 2 diabetes in Japanese individuals. <i>Journal of Diabetes Investigation</i> , 2017 , 8, 462-470	3.9	23
219	Proteomic analysis of serum biomarkers for prediabetes using the Long-Evans Agouti rat, a spontaneous animal model of type 2 diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2017 , 8, 661-673	3.9	15
218	Plasma adiponectin levels, ADIPOQ variants, and incidence of type 2 diabetes: A nested case-control study. <i>Diabetes Research and Clinical Practice</i> , 2017 , 127, 254-264	7.4	11
217	Effects of the Activation of Three Major Hepatic Akt Substrates on Glucose Metabolism in Male Mice. <i>Endocrinology</i> , 2017 , 158, 2659-2671	4.8	6
216	Dietary acid load and mortality among Japanese men and women: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2017 , 106, 146-154	7	22
215	Risk of Cardiovascular Events in Patients With Diabetes Mellitus on β -Blockers. <i>Hypertension</i> , 2017 , 70, 103-110	8.5	41
214	Trends in prevalence and management of diabetes and related vascular risks in Japanese adults: Japan National Health and Nutrition Surveys 2003-2012. <i>Diabetes Research and Clinical Practice</i> , 2017 , 127, 115-122	7.4	11
213	Effect of an intensified multifactorial intervention on cardiovascular outcomes and mortality in type 2 diabetes (J-DOIT3): an open-label, randomised controlled trial. <i>Lancet Diabetes and Endocrinology</i> , 2017 , 5, 951-964	18.1	141
212	Blood pressure-lowering effect of Shinrin-yoku (Forest bathing): a systematic review and meta-analysis. <i>BMC Complementary and Alternative Medicine</i> , 2017 , 17, 409	4.7	46
211	Comparison of Ipragliflozin and Pioglitazone Effects on Nonalcoholic Fatty Liver Disease in Patients With Type 2 Diabetes: A Randomized, 24-Week, Open-Label, Active-Controlled Trial. <i>Diabetes Care</i> , 2017 , 40, 1364-1372	14.6	154
210	Impact of birth weight on adult-onset diabetes mellitus in relation to current body mass index: The Japan Nurses' Health Study. <i>Journal of Epidemiology</i> , 2017 , 27, 428-434	3.4	24

209	High Serum Phospholipid Dihomo- Δ -Linoleic Acid Concentration and Low Δ -5-Desaturase Activity Are Associated with Increased Risk of Type 2 Diabetes among Japanese Adults in the Hitachi Health Study. <i>Journal of Nutrition</i> , 2017 , 147, 1558-1566	4.1	15
208	Postprandial Glucose Surges after Extremely Low Carbohydrate Diet in Healthy Adults. <i>Tohoku Journal of Experimental Medicine</i> , 2017 , 243, 35-39	2.4	5
207	A positive family history of hypertension might be associated with an accelerated onset of type 2 diabetes: Results from the National Center Diabetes Database (NCDD-02). <i>Endocrine Journal</i> , 2017 , 64, 515-520	2.9	6
206	Dietary patterns and all-cause, cancer, and cardiovascular disease mortality in Japanese men and women: The Japan public health center-based prospective study. <i>PLoS ONE</i> , 2017 , 12, e0174848	3.7	71
205	Effects of Ipragliflozin on Diabetic Nephropathy and Blood Pressure in Patients With Type 2 Diabetes: An Open-Label Study. <i>Journal of Clinical Medicine Research</i> , 2017 , 9, 154-162	2.9	10
204	The New Molecular Entity Evolocumab, One Kind of PCSK9 Inhibitor, Reduce Plasma Small Size LDL-Cholesterol Levels by Using a New Standardized Method of Measuring LDL Size. <i>Open Journal of Molecular and Integrative Physiology</i> , 2017 , 07, 1-23	1	2
203	Cross-Over Study Comparing Postprandial Glycemic Increase After Addition of a Fixed-Dose Mitiglinide/Voglibose Combination or a Dipeptidyl Peptidase-4 Inhibitor to Basal Insulin Therapy in Patients with Type 2 Diabetes Mellitus. <i>Medical Science Monitor Basic Research</i> , 2017 , 23, 36-44	3.2	
202	Time Spent Walking and Risk of Diabetes in Japanese Adults: The Japan Public Health Center-Based Prospective Diabetes Study. <i>Journal of Epidemiology</i> , 2016 , 26, 224-32	3.4	5
201	Body mass index and the risk of cancer incidence in patients with type 2 diabetes in Japan: Results from the National Center Diabetes Database. <i>Journal of Diabetes Investigation</i> , 2016 , 7, 908-914	3.9	9
200	Reduced serum level of leukocyte cell-derived chemotaxin 2 is associated with the presence of diabetic retinopathy. <i>Clinica Chimica Acta</i> , 2016 , 463, 145-149	6.2	8
199	Quality of diet and mortality among Japanese men and women: Japan Public Health Center based prospective study. <i>BMJ, The</i> , 2016 , 352, i1209	5.9	95
198	Design of and rationale for the Japan Diabetes Optimal Integrated Treatment study for 3 major risk factors of cardiovascular diseases (J-DOIT3): a multicenter, open-label, randomized, parallel-group trial. <i>BMJ Open Diabetes Research and Care</i> , 2016 , 4, e000123	4.5	19
197	Continuous Glucose Monitoring in Patients with Abnormal Glucose Tolerance during Pregnancy: A Case Series. <i>Japanese Clinical Medicine</i> , 2016 , 7, 1-8	1	6
196	Beneficial effects through aggressive coronary screening for type 2 diabetes patients with advanced vascular complications. <i>Medicine (United States)</i> , 2016 , 95, e4307	1.8	3
195	Report of the Japan Diabetes Society (JDS)/Japanese Cancer Association (JCA) Joint Committee on Diabetes and Cancer, Second Report. <i>Diabetology International</i> , 2016 , 7, 12-15	2.3	7
194	Accelerated decline of renal function in type 2 diabetes following severe hypoglycemia. <i>Journal of Diabetes and Its Complications</i> , 2016 , 30, 681-5	3.2	3
193	Association Between Severe Hypoglycemia and Cardiovascular Disease Risk in Japanese Patients With Type 2 Diabetes. <i>Journal of the American Heart Association</i> , 2016 , 5, e002875	6	40
192	A Patient-Held Medical Record Integrating Depression Care into Diabetes Care. <i>Japanese Clinical Medicine</i> , 2016 , 7, 19-22	1	1

191	Factors complicating the diabetes management of visitors to Japan: advices from a Japanese National Center for overseas medical staff. <i>Journal of Medical Investigation</i> , 2016 , 63, 15-8	1.2	1
190	Constipation, hard stools, fecal urgency, and incomplete evacuation, but not diarrhea is associated with diabetes and its related factors. <i>World Journal of Gastroenterology</i> , 2016 , 22, 3252-60	5.6	16
189	Visceral Fat Accumulation, Insulin Resistance, and Elevated Depressive Symptoms in Middle-Aged Japanese Men. <i>PLoS ONE</i> , 2016 , 11, e0149436	3.7	10
188	Increasing Number of People with Diabetes in Japan: Is This Trend Real?. <i>Internal Medicine</i> , 2016 , 55, 1827-30	1.1	29
187	Long-term Low-carbohydrate Diets and Type 2 Diabetes Risk: A Systematic Review and Meta-analysis of Observational Studies. <i>Journal of General and Family Medicine</i> , 2016 , 17, 60-70	1	8
186	A large-scale observational study to investigate the current status of diabetes complications and their prevention in Japan: research outline and baseline data for type 1 diabetes-JDCP study 2. <i>Diabetology International</i> , 2016 , 7, 4-11	2.3	16
185	Intensive Glycemic Therapy in Patients With Type 2 Diabetes on β Blockers. <i>Diabetes Care</i> , 2016 , 39, 1818-26	1.6	13
184	High hemoglobin A1c levels within the non-diabetic range are associated with the risk of all cancers. <i>International Journal of Cancer</i> , 2016 , 138, 1741-53	7.5	25
183	Report of the Japan diabetes society/Japanese cancer association joint committee on diabetes and cancer, Second report. <i>Cancer Science</i> , 2016 , 107, 369-71	6.9	6
182	Burden of cancer associated with type 2 diabetes mellitus in Japan, 2010-2030. <i>Cancer Science</i> , 2016 , 107, 521-7	6.9	11
181	Cluster-randomized trial to improve the quality of diabetes management: The study for the efficacy assessment of the standard diabetes manual (SEAS-DM). <i>Journal of Diabetes Investigation</i> , 2016 , 7, 539-43	3.9	5
180	Additive effects of miglitol and anagliptin on insulin-treated type 2 diabetes mellitus: a case study. <i>Clinical Drug Investigation</i> , 2015 , 35, 141-7	3.2	1
179	Abdominal Fat Accumulation, as Measured by Computed Tomography, Increases the Risk of Ischemic Colitis: A Retrospective Case-Control Study. <i>Digestive Diseases and Sciences</i> , 2015 , 60, 2104-11	4	4
178	Prediction of 90-day mortality in patients without diabetes by severe hypoglycemia: blood glucose level as a novel marker of severity of underlying disease. <i>Acta Diabetologica</i> , 2015 , 52, 307-14	3.9	29
177	Higher daily energy expenditure by locomotive activities is favorably associated with cardiac autonomic nervous function and arterial stiffness. <i>International Journal of Cardiology</i> , 2015 , 194, 70-1	3.2	1
176	Abdominal visceral fat accumulation measured by computed tomography associated with an increased risk of gallstone disease. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2015 , 30, 1323-31	4	9
175	Association of coffee intake with total and cause-specific mortality in a Japanese population: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2015 , 101, 1029-37	7	50
174	Association of green tea consumption with mortality due to all causes and major causes of death in a Japanese population: the Japan Public Health Center-based Prospective Study (JPHC Study). <i>Annals of Epidemiology</i> , 2015 , 25, 512-518.e3	6.4	54

173	Evidence-based practice guideline for the treatment for diabetes in Japan 2013. <i>Diabetology International</i> , 2015 , 6, 151-187	2.3	57
172	Visceral fat accumulation affects risk of colonic diverticular hemorrhage. <i>International Journal of Colorectal Disease</i> , 2015 , 30, 1399-406	3	12
171	The association between daily physical activity and plasma B-type natriuretic peptide in patients with glucose intolerance: a cross-sectional study. <i>BMJ Open</i> , 2015 , 5, e006276	3	16
170	A large-scale, observational study to investigate the current status of diabetes complications and their prevention in Japan: research outline and baseline data for type 2 diabetes JDCP study 1. <i>Diabetology International</i> , 2015 , 6, 243-251	2.3	12
169	Insulin receptor substrate-2 (Irs2) in endothelial cells plays a crucial role in insulin secretion. <i>Diabetes</i> , 2015 , 64, 876-86	0.9	28
168	Hemoglobin a1c levels and the risk of cardiovascular disease in people without known diabetes: a population-based cohort study in Japan. <i>Medicine (United States)</i> , 2015 , 94, e785	1.8	20
167	Impact of population aging on trends in diabetes prevalence: A meta-regression analysis of 160,000 Japanese adults. <i>Journal of Diabetes Investigation</i> , 2015 , 6, 533-42	3.9	75
166	Acute Multiple Arteriovenous Thromboses in a Patient with Diabetic Ketoacidosis. <i>Internal Medicine</i> , 2015 , 54, 2025-8	1.1	4
165	2) Diabetes and Its Treatment and Cancer. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2015 , 104, 1897-1900	0	
164	Analyzing the Factors Contributing to Withdrawal from Insulin Therapy Following Additional Administration of Alogliptin: Retrospective Study after Removing Glucotoxicity with Insulin. <i>Japanese Clinical Medicine</i> , 2015 , 6, 15-20	1	
163	Impact of medication adherence on renal function in comorbid patients with type 2 diabetes and depression: protocol for a cohort study. <i>BMC Family Practice</i> , 2015 , 16, 124	2.6	0
162	Visceral Abdominal Obesity Measured by Computed Tomography is Associated With Increased Risk of Colonic Diverticulosis. <i>Journal of Clinical Gastroenterology</i> , 2015 , 49, 816-22	3	23
161	Emerging Link between Diabetes and Cancer. <i>Journal of General and Family Medicine</i> , 2015 , 16, 170-176	1	
160	Verification of glycemic profiles using continuous glucose monitoring: cases with steroid use, liver cirrhosis, enteral nutrition, or late dumping syndrome. <i>Journal of Medical Investigation</i> , 2015 , 62, 1-10	1.2	10
159	Factors associated with untreated diabetes: analysis of data from 20,496 participants in the Japanese National Health and Nutrition Survey. <i>PLoS ONE</i> , 2015 , 10, e0118749	3.7	8
158	Low-carbohydrate diet and type 2 diabetes risk in Japanese men and women: the Japan Public Health Center-Based Prospective Study. <i>PLoS ONE</i> , 2015 , 10, e0118377	3.7	50
157	Daily Physical Activity Assessed by a Triaxial Accelerometer Is Beneficially Associated with Waist Circumference, Serum Triglycerides, and Insulin Resistance in Japanese Patients with Prediabetes or Untreated Early Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2015 , 2015, 526201	3.9	25
156	Possible discrepancy of HbA1c values and its assessment among patients with chronic renal failure, hemodialysis and other diseases. <i>Clinical and Experimental Nephrology</i> , 2015 , 19, 1179-83	2.5	16

155	High risk of abnormal QT prolongation in the early morning in diabetic and non-diabetic patients with severe hypoglycemia. <i>Annals of Medicine</i> , 2015 , 47, 238-44	1.5	12
154	Effectiveness of Prior Use of Beta-Blockers for Preventing Adverse Influences of Severe Hypoglycemia in Patients With Diabetes: An Observational Study. <i>Medicine (United States)</i> , 2015 , 94, e1629	1.8	10
153	Plasma insulin, C-peptide and blood glucose and the risk of gastric cancer: the Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2015 , 136, 1402-10	7.5	32
152	Effects of telephone-delivered lifestyle support on the development of diabetes in participants at high risk of type 2 diabetes: J-DOIT1, a pragmatic cluster randomised trial. <i>BMJ Open</i> , 2015 , 5, e007316	3	23
151	Diagnosed diabetes and premature death among middle-aged Japanese: results from a large-scale population-based cohort study in Japan (JPHC study). <i>BMJ Open</i> , 2015 , 5, e007736	3	20
150	Identification of a Hashimoto thyroiditis susceptibility locus via a genome-wide comparison with Graves disease. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, E319-24	5.6	18
149	Effect of body mass index on insulin secretion or sensitivity and diabetes. <i>American Journal of Preventive Medicine</i> , 2015 , 48, 128-135	6.1	24
148	Body Weight Gain and Hyperphagia After Administration of SGLT-2 Inhibitor: A Case Report. <i>American Journal of Case Reports</i> , 2015 , 16, 863-7	1.3	4
147	Effect of body mass index and intra-abdominal fat measured by computed tomography on the risk of bowel symptoms. <i>PLoS ONE</i> , 2015 , 10, e0123993	3.7	6
146	Constructing the National Center Diabetes Database. <i>Diabetology International</i> , 2014 , 5, 234-243	2.3	5
145	How can waist circumference predict the body composition?. <i>Diabetology and Metabolic Syndrome</i> , 2014 , 6, 11	5.6	12
144	Identification of independent susceptible and protective HLA alleles in Japanese autoimmune thyroid disease and their epistasis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, E379-83	5.6	43
143	The validity of the non-exercise activity thermogenesis questionnaire evaluated by objectively measured daily physical activity by the triaxial accelerometer. <i>BMC Sports Science, Medicine and Rehabilitation</i> , 2014 , 6, 27	2.4	10
142	Glucose management in diabetic patients undergoing hemodialysis. <i>Diabetology International</i> , 2014 , 5, 84-91	2.3	2
141	Descriptive epidemiology of diabetes prevalence and HbA1c distributions based on a self-reported questionnaire and a health checkup in the JPHC diabetes study. <i>Journal of Epidemiology</i> , 2014 , 24, 460-8	3.4	9
140	A newer conversion equation for the correlation between HbA1c and glycated albumin. <i>Endocrine Journal</i> , 2014 , 61, 553-60	2.9	20
139	Increase in homeostasis model assessment of insulin resistance (HOMA-IR) had a strong impact on the development of type 2 diabetes in Japanese individuals with impaired insulin secretion: the Saku study. <i>PLoS ONE</i> , 2014 , 9, e105827	3.7	20
138	The difficulties of interprofessional teamwork in diabetes care: a questionnaire survey. <i>Journal of Multidisciplinary Healthcare</i> , 2014 , 7, 333-9	2.8	12

137	The factors that limit activities of certified diabetes educators in Japan: a questionnaire survey. <i>SpringerPlus</i> , 2014 , 3, 611		4
136	Cholesterol and egg intakes and the risk of type 2 diabetes: the Japan Public Health Center-based Prospective Study. <i>British Journal of Nutrition</i> , 2014 , 112, 1636-43	3.6	33
135	Importance of high-density lipoprotein cholesterol levels in elderly diabetic individuals with type IIb dyslipidemia: A 2-year survey of cardiovascular events. <i>Geriatrics and Gerontology International</i> , 2014 , 14, 806-10	2.9	2
134	Efficacy of HMG-CoA reductase inhibitors in the prevention of cerebrovascular attack in 1016 patients older than 75 years among 4014 type 2 diabetic individuals. <i>International Journal of Cardiology</i> , 2014 , 177, 860-6	3.2	1
133	Visceral abdominal fat measured by computed tomography is associated with an increased risk of colorectal adenoma. <i>International Journal of Cancer</i> , 2014 , 135, 2273-81	7.5	45
132	Adiponectin and visceral fat associate with cardiovascular risk factors. <i>Obesity</i> , 2014 , 22, 287-91	8	20
131	Improvement of both fasting and postprandial glycemic control by the two-step addition of miglitol and mitiglinide to basal insulin therapy: a pilot study. <i>Diabetology and Metabolic Syndrome</i> , 2014 , 6, 48	5.6	5
130	Prediction of response to GLP-1 receptor agonist therapy in Japanese patients with type 2 diabetes. <i>Diabetology and Metabolic Syndrome</i> , 2014 , 6, 110	5.6	13
129	Predictors for cecal insertion time: the impact of abdominal visceral fat measured by computed tomography. <i>Diseases of the Colon and Rectum</i> , 2014 , 57, 1213-9	3.1	15
128	Seasonal variations of severe hypoglycemia in patients with type 1 diabetes mellitus, type 2 diabetes mellitus, and non-diabetes mellitus: clinical analysis of 578 hypoglycemia cases. <i>Medicine (United States)</i> , 2014 , 93, e148	1.8	20
127	Comparison of characteristics and healing course of diabetic foot ulcers by etiological classification: neuropathic, ischemic, and neuro-ischemic type. <i>Journal of Diabetes and Its Complications</i> , 2014 , 28, 528-35	3.2	36
126	Low-molecular-weight adiponectin and high-molecular-weight adiponectin levels in relation to diabetes. <i>Obesity</i> , 2014 , 22, 401-7	8	31
125	Vital signs, QT prolongation, and newly diagnosed cardiovascular disease during severe hypoglycemia in type 1 and type 2 diabetic patients. <i>Diabetes Care</i> , 2014 , 37, 217-25	14.6	74
124	Effects of exenatide in a morbidly obese patient with type 2 diabetes. <i>Diabetes Therapy</i> , 2014 , 5, 323-32	3.6	4
123	Correlations of non-exercise activity thermogenesis to metabolic parameters in Japanese patients with type 2 diabetes. <i>Diabetology and Metabolic Syndrome</i> , 2013 , 5, 26	5.6	25
122	A report on diabetes seminars for medical staff: findings from a questionnaire survey. <i>Diabetology International</i> , 2013 , 4, 261-265	2.3	2
121	Report of the JDS/JCA Joint Committee on Diabetes and Cancer. <i>Diabetology International</i> , 2013 , 4, 81-96	2.6	26
120	Japan Diabetes Outcome Intervention Trial-1 (J-DOIT1), a nationwide cluster randomized trial of type 2 diabetes prevention by telephone-delivered lifestyle support for high-risk subjects detected at health checkups: rationale, design, and recruitment. <i>BMC Public Health</i> , 2013 , 13, 81	4.1	6

119	Metabolic predictors of ischemic heart disease and cerebrovascular attack in elderly diabetic individuals: difference in risk by age. <i>Cardiovascular Diabetology</i> , 2013 , 12, 10	8.7	7
118	Latest insights into the risk of cancer in diabetes. <i>Journal of Diabetes Investigation</i> , 2013 , 4, 225-32	3.9	60
117	Report of the Japan Diabetes Society/Japanese Cancer Association Joint Committee on Diabetes and Cancer. <i>Cancer Science</i> , 2013 , 104, 965-76	6.9	56
116	The long-term coronary heart disease risk of previously obese patients with type 2 diabetes mellitus. <i>BMC Endocrine Disorders</i> , 2013 , 13, 38	3.3	3
115	Soft drink, 100% fruit juice, and vegetable juice intakes and risk of diabetes mellitus. <i>Clinical Nutrition</i> , 2013 , 32, 300-8	5.9	74
114	Non-exercise activity thermogenesis is associated with markers for diabetic microangiopathy in Japanese female patients with type 2 diabetes. <i>International Journal of Cardiology</i> , 2013 , 168, 4836-7	3.2	6
113	Severe hypoglycaemia and cardiovascular disease: systematic review and meta-analysis with bias analysis. <i>BMJ, The</i> , 2013 , 347, f4533	5.9	331
112	Impact of impaired insulin secretion and insulin resistance on the incidence of diabetes in a Japanese cohort. Reply to Yamauchi K and Aizawa T [letter]. <i>Diabetologia</i> , 2013 , 56, 2546-7	10.3	2
111	Red meat consumption is associated with the risk of type 2 diabetes in men but not in women: a Japan Public Health Center-based Prospective Study. <i>British Journal of Nutrition</i> , 2013 , 110, 1910-8	3.6	33
110	Effect of longitudinal changes in visceral fat area on incidence of metabolic risk factors: the Hitachi health study. <i>Obesity</i> , 2013 , 21, 2126-9	8	16
109	Diabetes care: After the Great East Japan Earthquake. <i>Journal of Diabetes Investigation</i> , 2013 , 4, 97-102	3.9	10
108	Effects of walking on medical cost: A quantitative evaluation by simulation focusing on diabetes. <i>Journal of Diabetes Investigation</i> , 2013 , 4, 667-72	3.9	9
107	A case of malignant insulinoma: successful control of glycemic fluctuation by replacing octreotide injections with octreotide LAR injections. <i>Endocrine Journal</i> , 2013 , 60, 951-7	2.9	3
106	Validity of diabetes self-reports in the Saku diabetes study. <i>Journal of Epidemiology</i> , 2013 , 23, 295-300	3.4	25
105	Vegetable and fruit intake and risk of type 2 diabetes: Japan Public Health Center-based Prospective Study. <i>British Journal of Nutrition</i> , 2013 , 109, 709-17	3.6	35
104	Dietary glycemic index, glycemic load and incidence of type 2 diabetes in Japanese men and women: the Japan Public Health Center-based Prospective Study. <i>Nutrition Journal</i> , 2013 , 12, 165	4.3	38
103	Dietary patterns and suicide in Japanese adults: the Japan Public Health Center-based Prospective Study. <i>British Journal of Psychiatry</i> , 2013 , 203, 422-7	5.4	34
102	Successful treatment of chronic intractable itching using oral pregabalin in a patient with diabetes and systemic prurigo nodularis: a case report of an iliopsoas muscle abscess. <i>Internal Medicine</i> , 2013 , 52, 2629-33	1.1	8

101	Effect of calcium channel blockers on incidence of diabetes: a meta-analysis. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2013 , 6, 257-61	3.4	21
100	History of having a macrosomic infant and the risk of diabetes: the Japan public health center-based prospective diabetes study. <i>PLoS ONE</i> , 2013 , 8, e84542	3.7	2
99	Low-carbohydrate diets and all-cause mortality: a systematic review and meta-analysis of observational studies. <i>PLoS ONE</i> , 2013 , 8, e55030	3.7	130
98	Reduction in adiposity, β cell function, insulin sensitivity, and cardiovascular risk factors: a prospective study among Japanese with obesity. <i>PLoS ONE</i> , 2013 , 8, e57964	3.7	11
97	Incidence of type 2 diabetes in Japan: a systematic review and meta-analysis. <i>PLoS ONE</i> , 2013 , 8, e74699	3.7	30
96	A morbid obese Japanese woman with a body mass index of 83.2 kg/m ² : before and after sleeve gastrectomy. <i>Internal Medicine</i> , 2012 , 51, 969-75	1.1	1
95	Significantly increased risk of cancer in diabetes mellitus patients: A meta-analysis of epidemiological evidence in Asians and non-Asians. <i>Journal of Diabetes Investigation</i> , 2012 , 3, 24-33	3.9	49
94	International clinical harmonization of glycated hemoglobin in Japan: From Japan Diabetes Society to National Glycohemoglobin Standardization Program values. <i>Journal of Diabetes Investigation</i> , 2012 , 3, 39-40	3.9	624
93	Associations of sex hormone-binding globulin and testosterone with diabetes among men and women (the Saku Diabetes study): a case control study. <i>Cardiovascular Diabetology</i> , 2012 , 11, 130	8.7	25
92	Dietary glycemic index and glycemic load in relation to HbA1c in Japanese obese adults: a cross-sectional analysis of the Saku Control Obesity Program. <i>Nutrition and Metabolism</i> , 2012 , 9, 79	4.6	5
91	Visceral fat area cutoff for the detection of multiple risk factors of metabolic syndrome in Japanese: the Hitachi Health Study. <i>Obesity</i> , 2012 , 20, 1744-9	8	19
90	Effect of the addition of sitagliptin and miglitol on insulin-treated type 2 diabetes. <i>Diabetes Therapy</i> , 2012 , 3, 11	3.6	7
89	International clinical harmonization of glycated hemoglobin in Japan: From Japan Diabetes Society to National Glycohemoglobin Standardization Program values. <i>Diabetology International</i> , 2012 , 3, 8-10	2.3	192
88	Adult mortality attributable to preventable risk factors for non-communicable diseases and injuries in Japan: a comparative risk assessment. <i>PLoS Medicine</i> , 2012 , 9, e1001160	11.6	144
87	Effect of longitudinal changes in visceral fat area and other anthropometric indices to the changes in metabolic risk factors in Japanese men: the Hitachi Health Study. <i>Diabetes Care</i> , 2012 , 35, 1139-43	14.6	23
86	Severely fluctuating blood glucose levels associated with a somatostatin-producing ovarian neuroendocrine tumor. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012 , 97, 3845-50	5.6	7
85	Improvement of glycemic control after periodontal treatment by resolving gingival inflammation in type 2 diabetic patients with periodontal disease. <i>Journal of Diabetes Investigation</i> , 2012 , 3, 402-9	3.9	13
84	FDG uptake by a condylomata acuminata in an HIV-infected patient mimicked urine contamination. <i>Clinical Nuclear Medicine</i> , 2012 , 37, 420-1	1.7	5

83	The Great East Japan Earthquake: Experiences and Suggestions for Survivors with Diabetes (perspective). <i>PLOS Currents</i> , 2012 , 4, e4fac9d99b997		8
82	Smoking cessation increases short-term risk of type 2 diabetes irrespective of weight gain: the Japan Public Health Center-Based Prospective Study. <i>PLoS ONE</i> , 2012 , 7, e17061	3.7	45
81	Cancer risk in diabetic patients treated with metformin: a systematic review and meta-analysis. <i>PLoS ONE</i> , 2012 , 7, e33411	3.7	401
80	ApoE isoforms, treatment of diabetes and the risk of coronary heart disease. <i>World Journal of Diabetes</i> , 2012 , 3, 54-9	4.7	5
79	Diabetes mellitus defined by hemoglobin A1c value: Risk characterization for incidence among Japanese subjects in the JPHC Diabetes Study. <i>Journal of Diabetes Investigation</i> , 2011 , 2, 359-65	3.9	3
78	What has made the population of Japan healthy?. <i>Lancet, The</i> , 2011 , 378, 1094-105	4.0	284
77	Efficacy of ezetimibe as monotherapy or combination therapy in hypercholesterolemic patients with and without diabetes. <i>Journal of Medical Investigation</i> , 2011 , 58, 86-94	1.2	15
76	Correlation between baseline serum 1,5-anhydroglucitol levels and 2-hour post-challenge glucose levels during oral glucose tolerance tests. <i>Endocrine Journal</i> , 2011 , 58, 13-7	2.9	19
75	Congener-specific polychlorinated biphenyls and the prevalence of diabetes in the Saku Control Obesity Program (SCOP). <i>Endocrine Journal</i> , 2011 , 58, 589-96	2.9	33
74	Significantly increased risk of cancer in patients with diabetes mellitus: a systematic review and meta-analysis. <i>Endocrine Practice</i> , 2011 , 17, 616-28	3.2	96
73	A patient with diabetes and breast cancer in whom virilization was caused by a testosterone-producing mature cystic teratoma containing a Brenner tumor. <i>American Journal of the Medical Sciences</i> , 2011 , 341, 74-7	2.2	7
72	Associations of smoking cessation with visceral fat area and prevalence of metabolic syndrome in men: the Hitachi health study. <i>Obesity</i> , 2011 , 19, 647-51	8	33
71	Social support and suicide in Japanese men and women - the Japan Public Health Center (JPHC)-based prospective study. <i>Journal of Psychiatric Research</i> , 2011 , 45, 1545-50	5.2	32
70	A pilot study of the efficacy of miglitol and sitagliptin for type 2 diabetes with a continuous glucose monitoring system and incretin-related markers. <i>Cardiovascular Diabetology</i> , 2011 , 10, 115	8.7	27
69	Long chain n-3 fatty acids intake, fish consumption and suicide in a cohort of Japanese men and women--the Japan Public Health Center-based (JPHC) prospective study. <i>Journal of Affective Disorders</i> , 2011 , 129, 282-8	6.6	33
68	Differences in suicide risk according to living arrangements in Japanese men and women--the Japan Public Health Center-based (JPHC) prospective study. <i>Journal of Affective Disorders</i> , 2011 , 131, 113-9	6.6	15
67	Insulin secretion and insulin sensitivity in Japanese patients with type 2 diabetes: a cross-sectional study comparing the homeostasis model assessment-2 (HOMA2) indexes and indexes derived from the oral glucose tolerance test. <i>Diabetology International</i> , 2011 , 2, 72-78	2.3	6
66	Validity and applicability of a simple questionnaire for the estimation of total and domain-specific physical activity. <i>Diabetology International</i> , 2011 , 2, 47-54	2.3	29

65	Japan Prevention Trial of Diabetes by Pitavastatin in Patients with Impaired Glucose Tolerance (the J-PREDICT study): rationale, study design, and clinical characteristics of 1269 patients. <i>Diabetology International</i> , 2011 , 2, 134-140	2.3	18
64	Fish intake and type 2 diabetes in Japanese men and women: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2011 , 94, 884-91	7	79
63	Asymptomatic coronary heart disease in patients with type 2 diabetes with vascular complications: a cross-sectional study. <i>BMJ Open</i> , 2011 , 1, e000139	3	10
62	Association of weight change in different periods of adulthood with risk of type 2 diabetes in Japanese men and women: the Japan Public Health Center-Based Prospective Study. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, 1104-10	5.1	38
61	Immunohistochemical findings in the pancreatic islets of a patient with transfusional iron overload and diabetes: case report. <i>Journal of Medical Investigation</i> , 2010 , 57, 345-9	1.2	14
60	Associations of visceral and subcutaneous fat areas with the prevalence of metabolic risk factor clustering in 6,292 Japanese individuals: the Hitachi Health Study. <i>Diabetes Care</i> , 2010 , 33, 2117-9	14.6	60
59	Soy product and isoflavone intakes are associated with a lower risk of type 2 diabetes in overweight Japanese women. <i>Journal of Nutrition</i> , 2010 , 140, 580-6	4.1	91
58	Autoimmune diabetes in HIV-infected patients on highly active antiretroviral therapy. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, 4056-60	5.6	23
57	Rice intake and type 2 diabetes in Japanese men and women: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2010 , 92, 1468-77	7	154
56	4. Iron Overload and Pancreatic Insufficiency/diabetes.. <i>The Journal of the Japanese Society of Internal Medicine</i> , 2010 , 99, 1255-1260	0	
55	Development of a database of health insurance claims: standardization of disease classifications and anonymous record linkage. <i>Journal of Epidemiology</i> , 2010 , 20, 413-9	3.4	211
54	Decreased insulin secretion and accumulation of triglyceride in beta cells overexpressing a dominant-negative form of AMP-activated protein kinase. <i>Endocrine Journal</i> , 2010 , 57, 141-52	2.9	11
53	Fasting plasma glucose and 5-year incidence of diabetes in the JPHC diabetes study - suggestion for the threshold for impaired fasting glucose among Japanese. <i>Endocrine Journal</i> , 2010 , 57, 629-37	2.9	27
52	Relationship between periodontitis and diabetes - importance of a clinical study to prove the vicious cycle. <i>Internal Medicine</i> , 2010 , 49, 881-5	1.1	24
51	Multifaceted intervention to promote the regular visiting of patients with diabetes to primary care physicians: rationale, design and conduct of a cluster-randomized controlled trial. The Japan Diabetes Outcome Intervention Trial-2 study protocol. <i>Diabetology International</i> , 2010 , 1, 83-89	2.3	10
50	Substantially increased risk of cancer in patients with diabetes mellitus: a systematic review and meta-analysis of epidemiologic evidence in Japan. <i>Journal of Diabetes and Its Complications</i> , 2010 , 24, 345-53	3.2	83
49	4. ??????????????????. <i>Japanese Journal of Clinical Pharmacology and Therapeutics</i> , 2010 , 41, 281-286	0	0
48	Fasting plasma glucose and incidence of diabetes --- implication for the threshold for impaired fasting glucose: results from the population-based Omiya MA cohort study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2009 , 16, 857-61	4	16

47	Metabolic factors and subsequent risk of hepatocellular carcinoma by hepatitis virus infection status: a large-scale population-based cohort study of Japanese men and women (JPHC Study Cohort II). <i>Cancer Causes and Control</i> , 2009 , 20, 741-50	2.8	46
46	Impact of metabolic factors on subsequent cancer risk: results from a large-scale population-based cohort study in Japan. <i>European Journal of Cancer Prevention</i> , 2009 , 18, 240-7	2	104
45	Primary aldosteronism associated with severe rhabdomyolysis due to profound hypokalemia. <i>Internal Medicine</i> , 2009 , 48, 219-23	1.1	25
44	Hemoglobin variant HbE found in two South Asian diabetic patients. <i>Internal Medicine</i> , 2009 , 48, 1397-401	1.1	3
43	Psychological factors, coffee and risk of diabetes mellitus among middle-aged Japanese: a population-based prospective study in the JPHC study cohort. <i>Endocrine Journal</i> , 2009 , 56, 459-68	2.9	90
42	Proteomic Studies on Investigations of Diabetes. <i>Journal of the Mass Spectrometry Society of Japan</i> , 2009 , 57, 201-206	0.2	0
41	BMI nomogram [corrected]. <i>Clinical Nutrition</i> , 2008 , 27, 168-9	5.9	5
40	A case of fulminant type 1 diabetes associated with significant elevation of mumps titers. <i>Endocrine Journal</i> , 2008 , 55, 561-4	2.9	27
39	Distribution of blood glucose and the correlation between blood glucose and hemoglobin A1c levels in diabetic outpatients. <i>Endocrine Journal</i> , 2008 , 55, 913-23	2.9	10
38	Hypopituitarism caused by bilateral internal carotid artery aneurysms with a carotid-cavernous fistula. <i>Internal Medicine</i> , 2008 , 47, 815-6	1.1	4
37	Anthropometric and Clinical Findings in Obese Japanese: The Saku Control Obesity Program (SCOP). <i>Anti-aging Medicine</i> , 2008 , 5, 13-16		8
36	Alcohol consumption and risk of type 2 diabetes mellitus in Japanese: a systematic review. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2008 , 17, 545-51	1	45
35	Extracellular proteome of human hepatoma cell, HepG2 analyzed using two-dimensional liquid chromatography coupled with tandem mass spectrometry. <i>Molecular and Cellular Biochemistry</i> , 2007 , 298, 83-92	4.2	38
34	An elderly case of type 2 diabetes which developed in association with oral and esophageal candidiasis. <i>Internal Medicine</i> , 2007 , 46, 387-90	1.1	7
33	Genetic variations of the ABC transporter gene ABCC3 in a Japanese population. <i>Drug Metabolism and Pharmacokinetics</i> , 2007 , 22, 129-35	2.2	19
32	Genetic variations and haplotype structures of transcriptional factor Nrf2 and its cytosolic reservoir protein Keap1 in Japanese. <i>Drug Metabolism and Pharmacokinetics</i> , 2007 , 22, 212-9	2.2	19
31	Possible aggravation and recovery of slowly progressive type 1 diabetes by onset and resolution of oral and esophageal candidiasis. <i>Internal Medicine</i> , 2007 , 46, 1629	1.1	1
30	Rapid-onset type 1 diabetes associated with cytomegalovirus infection and islet autoantibody synthesis. <i>Internal Medicine</i> , 2007 , 46, 873-7	1.1	13

29	Use of insulin glargine in Japanese patients with type 1 diabetes. <i>Internal Medicine</i> , 2007 , 46, 937-43	1.1	5
28	Glucokinase and IRS-2 are required for compensatory beta cell hyperplasia in response to high-fat diet-induced insulin resistance. <i>Journal of Clinical Investigation</i> , 2007 , 117, 246-57	15.9	262
27	Role of IRS and PHIP on insulin-induced tyrosine phosphorylation and distribution of IRS proteins. <i>Cell Structure and Function</i> , 2007 , 32, 69-78	2.2	6
26	Insulin allergy and immunologic insulin resistance caused by interleukin-6 in a patient with lung cancer. <i>Diabetes Care</i> , 2006 , 29, 1711-2	14.6	20
25	Hepatocyte nuclear factor-4alpha P2 promoter haplotypes are associated with type 2 diabetes in the Japanese population. <i>Diabetes</i> , 2006 , 55, 1260-4	0.9	28
24	Diabetes mellitus and the risk of cancer: results from a large-scale population-based cohort study in Japan. <i>Archives of Internal Medicine</i> , 2006 , 166, 1871-7		396
23	Role of uncoupling protein-2 up-regulation and triglyceride accumulation in impaired glucose-stimulated insulin secretion in a beta-cell lipotoxicity model overexpressing sterol regulatory element-binding protein-1c. <i>Endocrinology</i> , 2004 , 145, 3566-77	4.8	68
22	Pioglitazone reduces islet triglyceride content and restores impaired glucose-stimulated insulin secretion in heterozygous peroxisome proliferator-activated receptor-gamma-deficient mice on a high-fat diet. <i>Diabetes</i> , 2004 , 53, 2844-54	0.9	81
21	Chronological characterization of diabetes development in male Spontaneously Diabetic Torii rats. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 314, 870-7	3.4	100
20	Glucose metabolism and glutamate analog acutely alkalinize pH of insulin secretory vesicles of pancreatic beta-cells. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2003 , 285, E262-71 ⁶		37
19	Coffee consumption and risk of type 2 diabetes mellitus. <i>Lancet, The</i> , 2003 , 361, 703-4	4.0	47
18	The role of PPARgamma in high-fat diet-induced obesity and insulin resistance. <i>Journal of Diabetes and Its Complications</i> , 2002 , 16, 41-5	3.2	47
17	Switch to anaerobic glucose metabolism with NADH accumulation in the beta-cell model of mitochondrial diabetes. Characteristics of betaHC9 cells deficient in mitochondrial DNA transcription. <i>Journal of Biological Chemistry</i> , 2002 , 277, 41817-26	5.4	51
16	Genetic manipulations of fatty acid metabolism in beta-cells are associated with dysregulated insulin secretion. <i>Diabetes</i> , 2002 , 51 Suppl 3, S414-20	0.9	45
15	Phosphatidylinositol 3-kinase suppresses glucose-stimulated insulin secretion by affecting post-cytosolic [Ca(2+)] elevation signals. <i>Diabetes</i> , 2002 , 51, 87-97	0.9	69
14	Maturity-onset diabetes of the young resulting from a novel mutation in the HNF-4alpha gene. <i>Internal Medicine</i> , 2002 , 41, 848-52	1.1	7
13	A link between Cdc42 and syntaxin is involved in mastoparan-stimulated insulin release. <i>Biochemistry</i> , 2002 , 41, 9663-71	3.2	56
12	Hexamminecobalt(III) chloride inhibits glucose-induced insulin secretion at the exocytotic process. <i>Journal of Biological Chemistry</i> , 2001 , 276, 2979-85	5.4	6

11	Importance of standardization of hemoglobin A1c in the analysis of factors that predict hemoglobin A1c levels in non-diabetic residents of three distinct areas of Japan. <i>Diabetes Research and Clinical Practice</i> , 2001 , 53, 91-7	7.4	3
10	NADH shuttle system regulates K(ATP) channel-dependent pathway and steps distal to cytosolic Ca(2+) concentration elevation in glucose-induced insulin secretion. <i>Journal of Biological Chemistry</i> , 1999 , 274, 25386-92	5.4	46
9	Calbindin-D(28k) controls [Ca(2+)](i) and insulin release. Evidence obtained from calbindin-d(28k) knockout mice and beta cell lines. <i>Journal of Biological Chemistry</i> , 1999 , 274, 34343-9	5.4	95
8	Ethidium bromide-induced inhibition of mitochondrial gene transcription suppresses glucose-stimulated insulin release in the mouse pancreatic beta-cell line betaHC9. <i>Journal of Biological Chemistry</i> , 1998 , 273, 20300-7	5.4	83
7	Nutrient augmentation of Ca2+-dependent and Ca2+-independent pathways in stimulus-coupling to insulin secretion can be distinguished by their guanosine triphosphate requirements: studies on rat pancreatic islets. <i>Endocrinology</i> , 1998 , 139, 1172-83	4.8	34
6	Establishment of two substrains, diabetes-prone and non-diabetic, from Long-Evans Tokushima Lean (LETL) rats. <i>Endocrine Journal</i> , 1998 , 45, 737-44	2.9	40
5	Augmentation of insulin release by glucose in the absence of extracellular Ca2+: new insights into stimulus-secretion coupling. <i>Diabetes</i> , 1997 , 46, 1928-38	0.9	76
4	Pancreatic beta-cell-specific targeted disruption of glucokinase gene. Diabetes mellitus due to defective insulin secretion to glucose. <i>Journal of Biological Chemistry</i> , 1995 , 270, 30253-6	5.4	167
3	Nutrient Augmentation of Ca2+-Dependent and Ca2+-Independent Pathways in Stimulus-Coupling to Insulin Secretion Can Be Distinguished by Their Guanosine Triphosphate Requirements: Studies on Rat Pancreatic Islets		9
2	A Case of Type 2 Diabetes with Hemochromatosis: Efficacy of Changes in Insulin by Switching From Glargine to Degludec. <i>Medical Science Case Reports</i> , 2 , 72-77		
1	Short-Term Effect of Twice-Daily Dosage of Repaglinide in Elderly Patients with Type 2 Diabetes. <i>Medical Science Case Reports</i> , 2 , 53-57		