

# Torkell Ellingsen

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/7384832/torkell-ellingsen-publications-by-citations.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

108  
papers

2,796  
citations

25  
h-index

50  
g-index

124  
ext. papers

3,436  
ext. citations

4  
avg, IF

4.76  
L-index

#	Paper	IF	Citations
108	MRI bone oedema is the strongest predictor of subsequent radiographic progression in early rheumatoid arthritis. Results from a 2-year randomised controlled trial (CIMESTRA). <i>Annals of the Rheumatic Diseases</i> , <b>2009</b> , 68, 384-90	2.4	299
107	Treatment of lateral epicondylitis with platelet-rich plasma, glucocorticoid, or saline: a randomized, double-blind, placebo-controlled trial. <i>American Journal of Sports Medicine</i> , <b>2013</b> , 41, 625-35	6.8	241
106	Comparative effectiveness of injection therapies in lateral epicondylitis: a systematic review and network meta-analysis of randomized controlled trials. <i>American Journal of Sports Medicine</i> , <b>2013</b> , 41, 1435-46	6.8	170
105	Radiographic progression and remission rates in early rheumatoid arthritis - MRI bone oedema and anti-CCP predicted radiographic progression in the 5-year extension of the double-blind randomised CIMESTRA trial. <i>Annals of the Rheumatic Diseases</i> , <b>2010</b> , 69, 1789-95	2.4	136
104	A population-based cohort study of rheumatoid arthritis-associated interstitial lung disease: comorbidity and mortality. <i>Annals of the Rheumatic Diseases</i> , <b>2017</b> , 76, 1700-1706	2.4	131
103	Combination treatment with methotrexate, cyclosporine, and intraarticular betamethasone compared with methotrexate and intraarticular betamethasone in early active rheumatoid arthritis: an investigator-initiated, multicenter, randomized, double-blind, parallel-group, placebo-controlled study. <i>Arthritis and Rheumatism</i> , <b>2006</b> , 54, 1401-9		129
102	Ultrasound-Guided Injection Therapy of Achilles Tendinopathy With Platelet-Rich Plasma or Saline: A Randomized, Blinded, Placebo-Controlled Trial. <i>American Journal of Sports Medicine</i> , <b>2016</b> , 44, 1990-7	6.8	93
101	Neutrophil Extracellular Traps in Ulcerative Colitis: A Proteome Analysis of Intestinal Biopsies. <i>Inflammatory Bowel Diseases</i> , <b>2015</b> , 21, 2052-67	4.5	88
100	Plasma adiponectin in patients with active, early, and chronic rheumatoid arthritis who are steroid- and disease-modifying antirheumatic drug-naïve compared with patients with osteoarthritis and controls. <i>Journal of Rheumatology</i> , <b>2009</b> , 36, 1885-91	4.1	87
99	Adalimumab added to a treat-to-target strategy with methotrexate and intra-articular triamcinolone in early rheumatoid arthritis increased remission rates, function and quality of life. The OPERA Study: an investigator-initiated, randomised, double-blind, parallel-group, placebo-controlled trial. <i>Annals of the Rheumatic Diseases</i> , <b>2014</b> , 73, 654-61	2.4	84
98	Aggressive combination therapy with intra-articular glucocorticoid injections and conventional disease-modifying anti-rheumatic drugs in early rheumatoid arthritis: second-year clinical and radiographic results from the CIMESTRA study. <i>Annals of the Rheumatic Diseases</i> , <b>2008</b> , 67, 815-22	2.4	73
97	Short- and long-term efficacy of intra-articular injections with betamethasone as part of a treat-to-target strategy in early rheumatoid arthritis: impact of joint area, repeated injections, MRI findings, anti-CCP, IgM-RF and CRP. <i>Annals of the Rheumatic Diseases</i> , <b>2012</b> , 71, 851-6	2.4	55
96	Stereological quantification of immune-competent cells in baseline biopsy specimens from achilles tendons: results from patients with chronic tendinopathy followed for more than 4 years. <i>American Journal of Sports Medicine</i> , <b>2014</b> , 42, 2435-45	6.8	52
95	Prediction of treatment response to adalimumab: a double-blind placebo-controlled study of circulating microRNA in patients with early rheumatoid arthritis. <i>Pharmacogenomics Journal</i> , <b>2016</b> , 16, 141-6	3.5	44
94	A treat-to-target strategy with methotrexate and intra-articular triamcinolone with or without adalimumab effectively reduces MRI synovitis, osteitis and tenosynovitis and halts structural damage progression in early rheumatoid arthritis: results from the OPERA randomised controlled trial. <i>Annals of the Rheumatic Diseases</i> , <b>2017</b> , 76, 867-75	2.4	44
93	Effect of Magnetic Resonance Imaging vs Conventional Treat-to-Target Strategies on Disease Activity Remission and Radiographic Progression in Rheumatoid Arthritis: The IMAGINE-RA Randomized Clinical Trial. <i>JAMA - Journal of the American Medical Association</i> , <b>2019</b> , 321, 461-472	27.4	41
92	Discordance of Global Assessments by Patient and Physician Is Higher in Female than in Male Patients Regardless of the Physician's Sex: Data on Patients with Rheumatoid Arthritis, Axial Spondyloarthritis, and Psoriatic Arthritis from the DANBIO Registry. <i>Journal of Rheumatology</i> , <b>2015</b> , 42, 1781-5	4.1	41

91	Efficacy and safety of faecal microbiota transplantation in patients with psoriatic arthritis: protocol for a 6-month, double-blind, randomised, placebo-controlled trial. <i>BMJ Open</i> , <b>2018</b> , 8, e019231	3	41
90	Long-Term Prognosis of Plantar Fasciitis: A 5- to 15-Year Follow-up Study of 174 Patients With Ultrasound Examination. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2018</b> , 6, 2325967118757983	3.5	39
89	Plasma monocyte chemoattractant protein 1 is a marker for joint inflammation in rheumatoid arthritis. <i>Journal of Rheumatology</i> , <b>2001</b> , 28, 41-6	4.1	38
88	Folate, homocysteine, and cobalamin status in patients with rheumatoid arthritis treated with methotrexate, and the effect of low dose folic acid supplement. <i>Journal of Rheumatology</i> , <b>2004</b> , 31, 2374-81	4.1	36
87	Levels of lectin pathway proteins in plasma and synovial fluid of rheumatoid arthritis and osteoarthritis. <i>Rheumatology International</i> , <b>2012</b> , 32, 1457-63	3.6	34
86	Proteome Analysis of Rheumatoid Arthritis Gut Mucosa. <i>Journal of Proteome Research</i> , <b>2017</b> , 16, 346-354	3.6	33
85	Clinical and radiographic outcome of a treat-to-target strategy using methotrexate and intra-articular glucocorticoids with or without adalimumab induction: a 2-year investigator-initiated, double-blinded, randomised, controlled trial (OPERA). <i>Annals of the Rheumatic Diseases</i> , <b>2016</b> , 75, 1645-53	2.4	30
84	Differential effect of methotrexate on the increased CCR2 density on circulating CD4 T lymphocytes and monocytes in active chronic rheumatoid arthritis, with a down regulation only on monocytes in responders. <i>Annals of the Rheumatic Diseases</i> , <b>2007</b> , 66, 151-7	2.4	28
83	Ultrasonographic assessment of tendon thickness, Doppler activity and bony spurs of the elbow in patients with lateral epicondylitis and healthy subjects: a reliability and agreement study. <i>Ultraschall in Der Medizin</i> , <b>2013</b> , 34, 468-74	3.8	25
82	Localization of monocyte chemotactic and activating factor (MCAF/MCP-1) in psoriasis. <i>Journal of Dermatological Science</i> , <b>1996</b> , 13, 228-36	4.3	25
81	In active chronic rheumatoid arthritis, dipeptidyl peptidase IV density is increased on monocytes and CD4(+) T lymphocytes. <i>Scandinavian Journal of Immunology</i> , <b>2007</b> , 66, 451-7	3.4	24
80	Monocyte chemoattractant protein 1 (MCP-1) in temporal arteritis and polymyalgia rheumatica. <i>Annals of the Rheumatic Diseases</i> , <b>2000</b> , 59, 775-80	2.4	24
79	Periarticular and generalised bone loss in patients with early rheumatoid arthritis: influence of alendronate and intra-articular glucocorticoid treatment. Post hoc analyses from the CIMESTRA trial. <i>Annals of the Rheumatic Diseases</i> , <b>2014</b> , 73, 1123-9	2.4	23
78	Plasma MicroRNA Profiles in Patients with Early Rheumatoid Arthritis Responding to Adalimumab plus Methotrexate vs Methotrexate Alone: A Placebo-controlled Clinical Trial. <i>Journal of Rheumatology</i> , <b>2018</b> , 45, 53-61	4.1	23
77	Plasma level of CXC-chemokine CXCL12 is increased in rheumatoid arthritis and is independent of disease activity and methotrexate treatment. <i>Journal of Rheumatology</i> , <b>2006</b> , 33, 1754-9	4.1	23
76	Magnetic resonance imaging assessed inflammation in the wrist is associated with patient-reported physical impairment, global assessment of disease activity and pain in early rheumatoid arthritis: longitudinal results from two randomised controlled trials. <i>Annals of the Rheumatic Diseases</i> , <b>2017</b> , 76, 1707-14	2.4	21
75	Interactions between smoking, increased serum levels of anti-CCP antibodies, rheumatoid factors, and erosive joint disease in patients with early, untreated rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , <b>2015</b> , 44, 8-12	1.9	19
74	M-ficolin levels reflect disease activity and predict remission in early rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , <b>2013</b> , 65, 3045-50		18

73	Rheumatoid Arthritis-Associated Interstitial Lung Disease: Clinical Characteristics and Predictors of Mortality. <i>Respiration</i> , <b>2019</b> , 98, 455-460	3.7	17
72	A Proposal for a Study on Treatment Selection and Lifestyle Recommendations in Chronic Inflammatory Diseases: A Danish Multidisciplinary Collaboration on Prognostic Factors and Personalised Medicine. <i>Nutrients</i> , <b>2017</b> , 9,	6.7	16
71	Circulating surfactant protein -D is low and correlates negatively with systemic inflammation in early, untreated rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , <b>2010</b> , 12, R39	5.7	16
70	In vitro migration of mononuclear cells towards synovial fluid and plasma from rheumatoid arthritis patients correlates to RANTES synovial fluid levels and to clinical pain parameters. <i>Scandinavian Journal of Rheumatology</i> , <b>2000</b> , 29, 216-21	1.9	16
69	Left ventricular function at two-year follow-up in treatment-naive rheumatoid arthritis patients is associated with anti-cyclic citrullinated peptide antibody status: a cohort study. <i>Scandinavian Journal of Rheumatology</i> , <b>2017</b> , 46, 432-440	1.9	15
68	Development of heart failure in patients with rheumatoid arthritis: A Danish population-based study. <i>European Journal of Clinical Investigation</i> , <b>2018</b> , 48, e12915	4.6	15
67	Increased mortality among patients with rheumatoid arthritis and COPD: A population-based study. <i>Respiratory Medicine</i> , <b>2018</b> , 140, 101-107	4.6	15
66	Anti-cyclic citrullinated peptide antibodies, 28-joint Disease Activity Score, and magnetic resonance imaging bone oedema at baseline predict 11 years functional and radiographic outcome in early rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , <b>2019</b> , 48, 1-8	1.9	15
65	Urinary excretion of alpha-GST and albumin in rheumatoid arthritis patients treated with methotrexate or other DMARDs alone or in combination with NSAIDs. <i>Scandinavian Journal of Rheumatology</i> , <b>2005</b> , 34, 34-9	1.9	15
64	Galectin-3 is Persistently Increased in Early Rheumatoid Arthritis (RA) and Associates with Anti-CCP Seropositivity and MRI Bone Lesions, While Early Fibrosis Markers Correlate with Disease Activity. <i>Scandinavian Journal of Immunology</i> , <b>2017</b> , 86, 471-478	3.4	14
63	Predictive value of a multi-biomarker disease activity score for clinical remission and radiographic progression in patients with early rheumatoid arthritis: a post-hoc study of the OPERA trial. <i>Scandinavian Journal of Rheumatology</i> , <b>2019</b> , 48, 9-16	1.9	14
62	Cartilage oligomeric matrix protein associates differentially with erosions and synovitis and has a different temporal course in cyclic citrullinated peptide antibody (anti-CCP)-positive versus anti-CCP-negative early rheumatoid arthritis. <i>Journal of Rheumatology</i> , <b>2011</b> , 38, 1563-8	4.1	14
61	Use of Cerebrospinal Fluid Biomarkers in Diagnosis and Monitoring of Rheumatoid Meningitis. <i>Frontiers in Neurology</i> , <b>2019</b> , 10, 666	4.1	13
60	Upregulated baseline plasma CCL19 and CCR7 cell-surface expression on monocytes in early rheumatoid arthritis normalized during treatment and CCL19 correlated with radiographic progression. <i>Scandinavian Journal of Rheumatology</i> , <b>2014</b> , 43, 91-100	1.9	13
59	Left ventricular function in treatment-naive early rheumatoid arthritis. <i>American Journal of Cardiovascular Disease</i> , <b>2014</b> , 4, 79-86	0.9	13
58	Uncoupling of collagen II metabolism in newly diagnosed, untreated rheumatoid arthritis is linked to inflammation and antibodies against cyclic citrullinated peptides. <i>Journal of Rheumatology</i> , <b>2010</b> , 37, 1113-20	4.1	12
57	The effects of low-dose methotrexate on thymidylate synthetase activity in human peripheral blood mononuclear cells. <i>Clinical and Experimental Rheumatology</i> , <b>2000</b> , 18, 691-8	2.2	12
56	Ultrasonographic Characteristics of the Common Extensor Tendon of the Elbow in Asymptomatic Individuals: Thickness, Color Doppler Activity, and Bony Spurs. <i>Orthopaedic Journal of Sports Medicine</i> , <b>2017</b> , 5, 2325967117704186	3.5	11

55	Reproducibility of tender point examination in chronic low back pain patients as measured by intrarater and inter-rater reliability and agreement: a validation study. <i>BMJ Open</i> , <b>2013</b> , 3,	3	11
54	Impact of red and processed meat and fibre intake on treatment outcomes among patients with chronic inflammatory diseases: protocol for a prospective cohort study of prognostic factors and personalised medicine. <i>BMJ Open</i> , <b>2018</b> , 8, e018166	3	11
53	CRP genotype and haplotype associations with serum C-reactive protein level and DAS28 in untreated early rheumatoid arthritis patients. <i>Arthritis Research and Therapy</i> , <b>2014</b> , 16, 475	5.7	10
52	Active conventional treatment and three different biological treatments in early rheumatoid arthritis: phase IV investigator initiated, randomised, observer blinded clinical trial. <i>BMJ, The</i> , <b>2020</b> , 371, m4328	5.9	10
51	Multifactorial intervention to prevent cardiovascular disease in patients with early rheumatoid arthritis: protocol for a multicentre randomised controlled trial. <i>BMJ Open</i> , <b>2016</b> , 6, e009134	3	9
50	Calprotectin in patients with chronic rheumatoid arthritis correlates with disease activity and responsiveness to methotrexate. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , <b>2018</b> , 78, 62-67	2	9
49	Efficacy and safety of cannabidiol followed by an open label add-on of tetrahydrocannabinol for the treatment of chronic pain in patients with rheumatoid arthritis or ankylosing spondylitis: protocol for a multicentre, randomised, placebo-controlled study. <i>BMJ Open</i> , <b>2019</b> , 9, e028197	3	8
48	How do I establish a stool bank for fecal microbiota transplantation within the blood- and tissue transplant service?. <i>Transfusion</i> , <b>2020</b> , 60, 1135-1141	2.9	8
47	The D-vitamin metabolite 1,25(OH) D in serum is associated with disease activity and Anti-Citrullinated Protein Antibodies in active and treatment naïve, early Rheumatoid Arthritis Patients. <i>Scandinavian Journal of Immunology</i> , <b>2018</b> , 88, e12704	3.4	8
46	Safety and efficacy of faecal microbiota transplantation for active peripheral psoriatic arthritis: an exploratory randomised placebo-controlled trial. <i>Annals of the Rheumatic Diseases</i> , <b>2021</b> , 80, 1158-1167	2.4	8
45	Cardiovascular risk and mortality in rheumatoid arthritis compared with diabetes mellitus and the general population. <i>Rheumatology</i> , <b>2021</b> , 60, 1400-1409	3.9	8
44	Hand bone loss in early rheumatoid arthritis during a methotrexate-based treat-to-target strategy with or without adalimumab-a substudy of the optimized treatment algorithm in early RA (OPERA) trial. <i>Clinical Rheumatology</i> , <b>2017</b> , 36, 781-789	3.9	7
43	Effect of a treat-to-target strategy based on methotrexate and intra-articular betamethasone with or without additional cyclosporin on MRI-assessed synovitis, osteitis, tenosynovitis, bone erosion, and joint space narrowing in early rheumatoid arthritis: results from a 2-year randomized double-blind placebo-controlled trial (CIMESTRA). <i>Scandinavian Journal of Rheumatology</i> , <b>2017</b> , 46, 335-345	1.9	7
42	The Utility of 18F-FDG PET/CT in Patients With Clinical Suspicion of Polymyalgia Rheumatica and Giant Cell Arteritis: A Prospective, Observational, and Cross-sectional Study. <i>ACR Open Rheumatology</i> , <b>2020</b> , 2, 478-490	3.5	7
41	Impact of rheumatoid arthritis on major cardiovascular events in patients with and without coronary artery disease. <i>Annals of the Rheumatic Diseases</i> , <b>2020</b> , 79, 1182-1188	2.4	6
40	"To be lonely in your own loneliness": The interplay between self-perceived loneliness and rheumatoid arthritis in everyday life: A qualitative study. <i>Musculoskeletal Care</i> , <b>2020</b> , 18, 450-458	1.6	6
39	The impact of sex and disease classification on patient-reported outcome measures in axial spondyloarthritis: a descriptive prospective cross-sectional study. <i>Arthritis Research and Therapy</i> , <b>2019</b> , 21, 221	5.7	6
38	Insulin-like growth factor I receptor density on CD4+T-lymphocytes from active early steroid- and DMARD-naïve rheumatoid arthritis patients is up-regulated and not influenced by 1 year of clinically effective treatment. <i>Rheumatology International</i> , <b>2012</b> , 32, 501-4	3.6	6

37	Assessing the effect of interventions for axial spondyloarthritis according to the endorsed ASAS/OMERACT core outcome set: a meta-research study of trials included in Cochrane reviews. <i>Arthritis Research and Therapy</i> , <b>2020</b> , 22, 177	5.7	6
36	Outcome of COVID-19 in hospitalized patients with chronic inflammatory diseases. A population based national register study in Denmark. <i>Journal of Autoimmunity</i> , <b>2021</b> , 120, 102632	15.5	6
35	Sexual Health and Dysfunction in Patients With Rheumatoid Arthritis: A Cross-sectional Single-Center Study. <i>Sexual Medicine</i> , <b>2020</b> , 8, 615-630	2.7	5
34	Towards consensus in defining and handling contextual factors within rheumatology trials: an initial qualitative study from an OMERACT working group. <i>Annals of the Rheumatic Diseases</i> , <b>2021</b> , 80, 242-249 <sup>2,4</sup>	2.4	5
33	Prevalence of Newly Diagnosed Malignancies in Patients with Polymyalgia Rheumatica and Giant Cell Arteritis, Comparison of 18F-FDG PET/CT Scan with Chest X-ray and Abdominal Ultrasound: Data from a 40 Week Prospective, Exploratory, Single Centre Study. <i>Journal of Clinical Medicine</i> , <b>2020</b> , 9,	5.1	4
32	Clinical Value of Ultrasonographic Assessment in Lateral Epicondylitis Versus Asymptomatic Healthy Controls. <i>American Journal of Sports Medicine</i> , <b>2020</b> , 48, 1873-1883	6.8	4
31	Clinical characteristics of importance to outcome in patients with axial spondyloarthritis: protocol for a prospective descriptive and exploratory cohort study. <i>BMJ Open</i> , <b>2017</b> , 7, e015536	3	4
30	Heart Failure and Ischemic Heart Disease in Patients With Rheumatoid Arthritis. <i>Journal of the American College of Cardiology</i> , <b>2017</b> , 70, 3069-3071	15.1	4
29	Three Quality Improvement Initiatives and Performance of Rheumatoid Arthritis Disease Activity Measures in Electronic Health Records: Results From an Interrupted Time Series Study. <i>Arthritis Care and Research</i> , <b>2020</b> , 72, 283-291	4.7	4
28	Predictors of joint damage progression and stringent remission in patients with established rheumatoid arthritis in clinical remission. <i>Rheumatology</i> , <b>2021</b> , 60, 380-391	3.9	4
27	Elevated Neurofilament Light Chain in Cerebrospinal Fluid and Plasma Reflect Inflammatory MRI Activity in Neurosarcoidosis. <i>Brain Sciences</i> , <b>2021</b> , 11,	3.4	4
26	Association between baseline vitamin D metabolite levels and long-term cardiovascular events in patients with rheumatoid arthritis from the CIMESTRA trial: protocol for a cohort study with patient-record evaluated outcomes. <i>BMJ Open</i> , <b>2017</b> , 7, e014816	3	2
25	Proteomics dataset: The colon mucosa from inflammatory bowel disease patients, gastrointestinal asymptomatic rheumatoid arthritis patients, and controls. <i>Data in Brief</i> , <b>2017</b> , 15, 511-516	1.2	2
24	Population characteristics as important contextual factors in rheumatological trials: an exploratory meta-epidemiological study from an OMERACT Working Group. <i>Annals of the Rheumatic Diseases</i> , <b>2020</b> , 79, 1269-1276	2.4	2
23	Circulating serum interleukin-6, serum chitinase-3-like protein-1, and plasma vascular endothelial growth factor are not predictive for remission and radiographic progression in patients with early rheumatoid arthritis: post-hoc explorative and validation studies based on the CIMESTRA and OPERA trials. <i>Scandinavian Journal of Rheumatology</i> , <b>2018</b> , 47, 259-269	1.9	2
22	OMERACT consensus-based operational definition of contextual factors in rheumatology clinical trials: A mixed methods study. <i>Seminars in Arthritis and Rheumatism</i> , <b>2021</b> , 51, 601-606	5.3	2
21	Prognostic impacts of glucocorticoid treatment in patients with polymyalgia rheumatica and giant cell arteritis. <i>Scientific Reports</i> , <b>2021</b> , 11, 6220	4.9	2
20	Similar lipid level changes in early rheumatoid arthritis patients following 1-year treat-to-target strategy with adalimumab plus methotrexate versus placebo plus methotrexate: secondary analyses from the randomised controlled OPERA trial. <i>Rheumatology International</i> , <b>2021</b> , 41, 543-549	3.6	2

19	Impact of season on the association between vitamin D levels at diagnosis and one-year remission in early Rheumatoid Arthritis. <i>Scientific Reports</i> , <b>2020</b> , 10, 7371	4.9	1
18	[F]Sodium fluoride positron emission tomography/computed tomography: a predictor of early rheumatoid arthritis? A case report. <i>Scandinavian Journal of Rheumatology</i> , <b>2018</b> , 47, 425-426	1.9	1
17	Up-regulated CD26 density on synovial fluid monocytes from patients with juvenile idiopathic arthritis. <i>Scandinavian Journal of Rheumatology</i> , <b>2008</b> , 37, 76-8	1.9	1
16	Experiences and perceptions of patients with psoriatic arthritis participating in a trial of faecal microbiota transplantation: a nested qualitative study. <i>BMJ Open</i> , <b>2021</b> , 11, e039471	3	1
15	Response to: Correspondence on Safety and efficacy of faecal microbiota transplantation for active peripheral psoriatic arthritis: an exploratory randomised placebo-controlled trial by McGonagle. <i>Annals of the Rheumatic Diseases</i> , <b>2021</b> ,	2.4	1
14	How to increase value and reduce waste in research: initial experiences of applying Lean thinking and visual management in research leadership. <i>BMJ Open</i> , <b>2022</b> , 12, e058179	3	1
13	Clinical presentation and treatment response in patients with polymyalgia rheumatica and giant cell arteritis during a 40-week follow-up.. <i>Rheumatology Advances in Practice</i> , <b>2021</b> , 5, rkab091	1.1	0
12	Altered retinal oxygen metabolism in patients with combined ocular and central nervous system sarcoidosis. <i>Rheumatology</i> , <b>2021</b> , 60, 3301-3306	3.9	0
11	Methotrexate pharmacokinetic is influenced by co-administration of cyclosporin in rheumatoid arthritis patients. Results from a randomized clinical trial. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , <b>2020</b> , 80, 185-190	2	0
10	FRI0067 Changes in Multi-Biomarker Disease Activity (MBDA) Score Correlate with Changes in Established Disease Activity Measurements in Patients with Early Ra from The Opera Study. <i>Annals of the Rheumatic Diseases</i> , <b>2016</b> , 75, 450.1-450	2.4	0
9	Inflammatory profiles in plasma and cerebrospinal fluid of patients with neurosarcoidosis.. <i>Journal of Neuroimmunology</i> , <b>2022</b> , 367, 577849	3.5	0
8	The Use of Telehealth for Psychological Counselling of Vulnerable Adult Patients With Rheumatic Diseases or Diabetes: Explorative Study Inspired by Participatory Design.. <i>JMIR Human Factors</i> , <b>2022</b> , 9, e30829	2.5	
7	FRI0203 A SINGLE-CENTER INVESTIGATION ON THE PREVALENCE OF MALIGNANCIES IN PATIENTS WITH POLYMYALGIA RHEUMATICA AND GIANT CELL ARTERITIS BY WAY OF 18F-FDG PET/CT: A PROSPECTIVE COHORT STUDY. <i>Annals of the Rheumatic Diseases</i> , <b>2020</b> , 79, 685.1-685	2.4	
6	THU0614-HPR ASSESSING THE EFFECT OF INTERVENTIONS FOR AXIAL SPONDYLOARTHRITIS ACCORDING TO THE ENDORSED ASAS/OMERACT CORE OUTCOME SET: A META-RESEARCH STUDY OF TRIALS INCLUDED IN COCHRANE REVIEWS. <i>Annals of the Rheumatic Diseases</i> , <b>2020</b> , 79, 548.2-550	2.4	
5	Chronic Lateral Epicondylitis New Perspectives on Injection Strategies and Ultrasound Evaluation <b>2015</b> , 23-31		
4	Ultrasound-guided injection therapy with platelet-rich plasma in recreational athletes with patellar tendinopathy: A randomized, single blinded, placebo-controlled trial with 3 months follow-up. <i>Translational Sports Medicine</i> , <b>2021</b> , 4, 344-355	1.3	
3	Comment on: Cardiovascular risk and mortality in rheumatoid arthritis compared with diabetes mellitus and the general population: reply. <i>Rheumatology</i> , <b>2021</b> , 60, e419-e420	3.9	
2	Effect of initiating biologics compared to intensifying conventional DMARDs on clinical and MRI outcomes in established rheumatoid arthritis patients in clinical remission: Secondary analyses of the IMAGINE-RA trial. <i>Scandinavian Journal of Rheumatology</i> , <b>2021</b> , 1-11	1.9	

- 1 Risk of harm in synthetic and biological intervention trials in patients with inflammatory arthritis: protocol for a metaepidemiological study focusing on contextual factors. *BMJ Open*, **2021**, 11, e049850 <sup>3</sup>