Antoine Depaulis

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/738520/antoine-depaulis-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

 155
 10,296
 56
 98

 papers
 citations
 h-index
 g-index

 161
 11,266
 4.9
 5.64

 ext. papers
 ext. citations
 avg, IF
 L-index

#	Paper	IF	Citations
155	Neurovascular multiparametric MRI defines epileptogenic and seizure propagation regions in experimental mesiotemporal lobe epilepsy. <i>Epilepsia</i> , 2021 , 62, 1244-1255	6.4	2
154	In vivo Eminobutyric acid increase as a biomarker of the epileptogenic zone: An unbiased metabolomics approach. <i>Epilepsia</i> , 2021 , 62, 163-175	6.4	7
153	Reprogramming reactive glia into interneurons reduces chronic seizure activity in a mouse model of mesial temporal lobe epilepsy. <i>Cell Stem Cell</i> , 2021 , 28, 2104-2121.e10	18	9
152	Early reduced dopaminergic tone mediated by D3 receptor and dopamine transporter in absence epileptogenesis. <i>Epilepsia</i> , 2019 , 60, 2128-2140	6.4	4
151	Sensory coding is impaired in rat absence epilepsy. <i>Journal of Physiology</i> , 2019 , 597, 951-966	3.9	11
150	Pathophysiology of absence epilepsy: Insights from genetic models. <i>Neuroscience Letters</i> , 2018 , 667, 53-65	3.3	26
149	Glial responses during epileptogenesis in Mus musculus point to potential therapeutic targets. <i>PLoS ONE</i> , 2018 , 13, e0201742	3.7	10
148	WONOEP appraisal: Biomarkers of epilepsy-associated comorbidities. <i>Epilepsia</i> , 2017 , 58, 331-342	6.4	26
147	Building Up Absence Seizures in the Somatosensory Cortex: From Network to Cellular Epileptogenic Processes. <i>Cerebral Cortex</i> , 2017 , 27, 4607-4623	5.1	22
146	NADPH oxidases as drug targets and biomarkers in neurodegenerative diseases: What is the evidence?. <i>Free Radical Biology and Medicine</i> , 2017 , 112, 387-396	7.8	60
145	Identification and characterization of outcome measures reported in animal models of epilepsy: Protocol for a systematic review of the literature-A TASK2 report of the AES/ILAE Translational Task Force of the ILAE. <i>Epilepsia</i> , 2017 , 58 Suppl 4, 68-77	6.4	5
144	Genetic Models of Absence Epilepsy in Rats and Mice 2017 , 455-471		5
143	Experimental Treatment Options in Absence Epilepsy. Current Pharmaceutical Design, 2017, 23, 5577-5	5923	7
142	The genetic absence epilepsy rat from Strasbourg as a model to decipher the neuronal and network mechanisms of generalized idiopathic epilepsies. <i>Journal of Neuroscience Methods</i> , 2016 , 260, 159-74	3	70
141	Synchrotron X-ray microtransections: a non invasive approach for epileptic seizures arising from eloquent cortical areas. <i>Scientific Reports</i> , 2016 , 6, 27250	4.9	13
140	Differential Effects of Antiepileptic Drugs on Focal Seizures in the Intrahippocampal Kainate Mouse Model of Mesial Temporal Lobe Epilepsy. <i>CNS Neuroscience and Therapeutics</i> , 2016 , 22, 497-506	6.8	48
139	High-Throughput LC-MS/MS Proteomic Analysis of a Mouse Model of Mesiotemporal Lobe Epilepsy Predicts Microglial Activation Underlying Disease Development. <i>Journal of Proteome Research</i> , 2016 , 15, 1546-62	5.6	24

(2012-2016)

138	Activation of GABA receptors controls mesiotemporal lobe epilepsy despite changes in chloride transporters expression: In vivo and in silico approach. <i>Experimental Neurology</i> , 2016 , 284, 11-28	5.7	15	
137	Assessing Susceptibility to Epilepsy in Three Rat Strains Using Brain Metabolic Profiling Based on HRMAS NMR Spectroscopy and Chemometrics. <i>Journal of Proteome Research</i> , 2015 , 14, 2177-89	5.6	15	
136	Animal models for mesiotemporal lobe epilepsy: The end of a misunderstanding?. <i>Revue Neurologique</i> , 2015 , 171, 217-26	3	20	
135	Synchrotron X-ray microbeams: A promising tool for drug-resistant epilepsy treatment. <i>Physica Medica</i> , 2015 , 31, 607-14	2.7	14	
134	Microfabrication, characterization and in vivo MRI compatibility of diamond microelectrodes array for neural interfacing. <i>Materials Science and Engineering C</i> , 2015 , 46, 25-31	8.3	21	
133	Revisiting hippocampal sclerosis in mesial temporal lobe epilepsy according to the "two-hit" hypothesis. <i>Revue Neurologique</i> , 2015 , 171, 227-35	3	11	
132	Activation of mTOR signaling pathway is secondary to neuronal excitability in a mouse model of mesio-temporal lobe epilepsy. <i>European Journal of Neuroscience</i> , 2015 , 41, 976-88	3.5	41	
131	Seizure expression, behavior, and brain morphology differences in colonies of Genetic Absence Epilepsy Rats from Strasbourg. <i>Epilepsia</i> , 2014 , 55, 1959-68	6.4	47	
130	Epilepsy: Animal Models to Reproduce Human Etiopathology. <i>Methods and Principles in Medicinal Chemistry</i> , 2014 , 415-430	0.4		
129	Long-term modifications of epileptogenesis and hippocampal rhythms after prolonged hyperthermic seizures in the mouse. <i>Neurobiology of Disease</i> , 2014 , 69, 156-68	7.5	8	
128	Neural adaptation to responsive stimulation: a comparison of auditory and deep brain stimulation in a rat model of absence epilepsy. <i>Brain Stimulation</i> , 2013 , 6, 241-7	5.1	22	
127	Occurrence of the synthetic analgesic tramadol in an African medicinal plant. <i>Angewandte Chemie - International Edition</i> , 2013 , 52, 11780-4	16.4	27	
126	Do seizures and epileptic activity worsen epilepsy and deteriorate cognitive function?. <i>Epilepsia</i> , 2013 , 54 Suppl 8, 14-21	6.4	46	
125	Synchrotron X-ray interlaced microbeams suppress paroxysmal oscillations in neuronal networks initiating generalized epilepsy. <i>Neurobiology of Disease</i> , 2013 , 51, 152-60	7.5	22	
124	Occurrence of the Synthetic Analgesic Tramadol in an African Medicinal Plant. <i>Angewandte Chemie</i> , 2013 , 125, 11996-12000	3.6	5	
123	Is ictal dystonia associated with an inhibitory effect on seizure propagation in focal epilepsies?. <i>Epilepsy Research</i> , 2012 , 99, 274-80	3	10	
122	Animal models to study aetiopathology of epilepsy: what are the features to model?. <i>Epileptic Disorders</i> , 2012 , 14, 217-25	1.9	34	
121	Specific in vivo staining of astrocytes in the whole brain after intravenous injection of sulforhodamine dyes. <i>PLoS ONE</i> , 2012 , 7, e35169	3.7	52	

120	Inflammatory changes during epileptogenesis and spontaneous seizures in a mouse model of mesiotemporal lobe epilepsy. <i>Epilepsia</i> , 2011 , 52, 2315-25	6.4	95
119	Increase in BDNF-mediated TrkB signaling promotes epileptogenesis in a mouse model of mesial temporal lobe epilepsy. <i>Neurobiology of Disease</i> , 2011 , 42, 35-47	7.5	142
118	Dentate gyrus and hilus transection blocks seizure propagation and granule cell dispersion in a mouse model for mesial temporal lobe epilepsy. <i>Hippocampus</i> , 2011 , 21, 334-43	3.5	37
117	Radiation Therapy Using Synchrotron Radiation: Preclinical Studies Toward Clinical Trials. <i>Synchrotron Radiation News</i> , 2011 , 24, 8-12	0.6	1
116	Deep brain stimulation in epilepsy: what is next?. Current Opinion in Neurology, 2010, 23, 177-82	7.1	59
115	Involvement of the thalamic parafascicular nucleus in mesial temporal lobe epilepsy. <i>Journal of Neuroscience</i> , 2010 , 30, 16523-35	6.6	43
114	In silico dynamic molecular interaction networks for the discovery of new therapeutic targets. <i>Current Pharmaceutical Design</i> , 2010 , 16, 2241-51	3.3	2
113	Comparative study of five antiepileptic drugs on a translational cognitive measure in the rat: relationship to antiepileptic property. <i>Psychopharmacology</i> , 2010 , 207, 513-27	4.7	20
112	High-precision radiosurgical dose delivery by interlaced microbeam arrays of high-flux low-energy synchrotron X-rays. <i>PLoS ONE</i> , 2010 , 5, e9028	3.7	69
111	Manipulating the epileptic brain using stimulation: a review of experimental and clinical studies. <i>Epileptic Disorders</i> , 2009 , 11, 100-12	1.9	40
110	La souris MTLE: un´modle validlpour´l¶aluation de´mollules anti-Bileptiques pour le´traitement de l¶ilepsie mBiotemporale. <i>Epilepsies</i> , 2009 , 21, 184-192		O
109	Long-term effects of febrile status epilepticus: What animal models can tell us?. <i>Epilepsia</i> , 2009 , 50 Suppl 12, 27-8	6.4	3
108	Identifying neural drivers with functional MRI: an electrophysiological validation. <i>PLoS Biology</i> , 2008 , 6, 2683-97	9.7	392
107	Epilepsy in Dcx knockout mice associated with discrete lamination defects and enhanced excitability in the hippocampus. <i>PLoS ONE</i> , 2008 , 3, e2473	3.7	56
106	Fetal exposure to GABA-acting antiepileptic drugs generates hippocampal and cortical dysplasias. <i>Epilepsia</i> , 2007 , 48, 684-93	6.4	100
105	Controlling seizures is not controlling epilepsy: a parametric study of deep brain stimulation for epilepsy. <i>Neurobiology of Disease</i> , 2007 , 27, 292-300	7.5	57
104	Right temporal cerebral dysfunction heralds symptoms of acute mountain sickness. <i>Journal of Neurology</i> , 2007 , 254, 359-63	5.5	13
103	Deep layer somatosensory cortical neurons initiate spike-and-wave discharges in a genetic model of absence seizures. <i>Journal of Neuroscience</i> , 2007 , 27, 6590-9	6.6	307

(2001-2007)

102	Short-term changes in bilateral hippocampal coherence precede epileptiform events. <i>NeuroImage</i> , 2007 , 38, 138-49	7.9	33
101	Genetic Models of Absence Epilepsy in the Rat 2006 , 233-248		45
100	Reelin deficiency and displacement of mature neurons, but not neurogenesis, underlie the formation of granule cell dispersion in the epileptic hippocampus. <i>Journal of Neuroscience</i> , 2006 , 26, 4701-13	6.6	235
99	Evidence for a role of the parafascicular nucleus of the thalamus in the control of epileptic seizures by the superior colliculus. <i>Epilepsia</i> , 2005 , 46, 141-5	6.4	28
98	Glutamate receptor antagonists and benzodiazepine inhibit the progression of granule cell dispersion in a mouse model of mesial temporal lobe epilepsy. <i>Epilepsia</i> , 2005 , 46, 193-202	6.4	49
97	Hypothalamic response to experimental allergic encephalomyelitis: role of substance P. <i>NeuroImmunoModulation</i> , 2004 , 11, 28-35	2.5	4
96	High temporal resolution for in vivo monitoring of neurotransmitters in awake epileptic rats using brain microdialysis and capillary electrophoresis with laser-induced fluorescence detection. <i>Journal of Neuroscience Methods</i> , 2004 , 140, 29-38	3	77
95	A genetic switch for epilepsy in adult mice. <i>Journal of Neuroscience</i> , 2004 , 24, 10568-78	6.6	63
94	Modifications of local cerebral glucose utilization in thalamic structures following injection of a dopaminergic agonist in the nucleus accumbensinvolvement in antiepileptic effects?. <i>Experimental Neurology</i> , 2004 , 188, 452-60	5.7	9
93	Physiopathologie des crises d\$ilepsie. <i>EMC - Neurologie</i> , 2004 , 1, 1-10		
92	PET evidence for a role of the basal ganglia in patients with ring chromosome 20 epilepsy. <i>Neurology</i> , 2004 , 63, 73-7	6.5	74
91	Neuropeptide Y delays hippocampal kindling in the rat. <i>Hippocampus</i> , 2003 , 13, 557-60	3.5	34
90	Induced down-regulation of neuropeptide Y-Y1 receptors delays initiation of kindling. <i>European Journal of Neuroscience</i> , 2003 , 18, 768-74	3.5	28
89	Suppression of absence seizures by electrical and pharmacological activation of the caudal superior colliculus in a genetic model of absence epilepsy in the rat. <i>Experimental Neurology</i> , 2002 , 177, 503-14	5.7	19
88	Evolution of hippocampal epileptic activity during the development of hippocampal sclerosis in a mouse model of temporal lobe epilepsy. <i>Neuroscience</i> , 2002 , 112, 101-11	3.9	295
87	Control of Epileptic Seizures. <i>Advances in Behavioral Biology</i> , 2002 , 169-178		
86	The control of seizures by the basal ganglia? A review of experimental data. <i>Epileptic Disorders</i> , 2002 , 4 Suppl 3, S61-72	1.9	28
85	Neuropeptide Y and epilepsy: varying effects according to seizure type and receptor activation. <i>Peptides</i> , 2001 , 22, 529-39	3.8	44

84	Inhibition of the substantia nigra suppresses absences and clonic seizures in audiogenic rats, but not tonic seizures: evidence for seizure specificity of the nigral control. <i>Neuroscience</i> , 2001 , 105, 203-11	3.9	90
83	Different representations of inescapable noxious stimuli in the periaqueductal gray and upper cervical spinal cord of freely moving rats. <i>Neuroscience Letters</i> , 2001 , 313, 17-20	3.3	45
82	BDNF and epilepsythe bad could turn out to be good. <i>Trends in Neurosciences</i> , 2001 , 24, 318-9	13.3	26
81	Overexpression of neuropeptide Y induced by brain-derived neurotrophic factor in the rat hippocampus is long lasting. <i>European Journal of Neuroscience</i> , 2000 , 12, 595-605	3.5	67
8o	Endogenous control of hippocampal epileptogenesis: a molecular cascade involving brain-derived neurotrophic factor and neuropeptide Y. <i>Epilepsia</i> , 2000 , 41 Suppl 6, S127-33	6.4	43
79	Dopamine in the striatum modulates seizures in a genetic model of absence epilepsy in the rat. <i>Neuroscience</i> , 2000 , 100, 335-44	3.9	97
78	Brain-derived neurotrophic factor delays hippocampal kindling in the rat. Neuroscience, 2000, 100, 777-8	8§ .9	74
77	Low-voltage-activated calcium channel subunit expression in a genetic model of absence epilepsy in the rat. <i>Molecular Brain Research</i> , 2000 , 75, 159-65		119
76	Neuroprotective effects of chronic estradiol benzoate treatment on hippocampal cell loss induced by status epilepticus in the female rat. <i>Neuroscience Letters</i> , 2000 , 281, 79-82	3.3	65
75	Evidence for the involvement of the pallidum in the modulation of seizures in a genetic model of absence epilepsy in the rat. <i>Neuroscience Letters</i> , 1999 , 265, 131-4	3.3	29
74	Recurrent seizures and hippocampal sclerosis following intrahippocampal kainate injection in adult mice: electroencephalography, histopathology and synaptic reorganization similar to mesial temporal lobe epilepsy. <i>Neuroscience</i> , 1999 , 89, 717-29	3.9	314
73	High-frequency stimulation of the subthalamic nucleus suppresses absence seizures in the rat: comparison with neurotoxic lesions. <i>Epilepsy Research</i> , 1998 , 31, 39-46	3	165
72	The role of basal ganglia in the control of generalized absence seizures. <i>Epilepsy Research</i> , 1998 , 32, 213	3-323	131
71	Role of the subthalamo-nigral input in the control of amygdala-kindled seizures in the rat. <i>Brain Research</i> , 1998 , 807, 78-83	3.7	55
70	Pathophysiological mechanisms of genetic absence epilepsy in the rat. <i>Progress in Neurobiology</i> , 1998 , 55, 27-57	10.9	483
69	Protective Effects of Brain-Derived Neurotrophic Factor in Hippocampal Kindling. <i>Advances in Behavioral Biology</i> , 1998 , 409-420		1
68	Anxiogenic-like consequences in animal models of complex partial seizures. <i>Neuroscience and Biobehavioral Reviews</i> , 1997 , 21, 767-74	9	56
67	Involvement of nigral glutamatergic inputs in the control of seizures in a genetic model of absence epilepsy in the rat. <i>Neuroscience</i> . 1996 . 71. 721-8	3.9	68

66	Amygdala kindling in the rat: anxiogenic-like consequences. <i>Neuroscience</i> , 1996 , 73, 971-8	3.9	79
65	Ultrasonic vocalization (22-28 kHz) in a model of chronic pain, the arthritic rat: effects of analgesic drugs. <i>NeuroReport</i> , 1996 , 7, 581-4	1.7	72
64	Parkinsonian-like locomotor impairment in mice lacking dopamine D2 receptors. <i>Nature</i> , 1995 , 377, 424	l-§ 0.4	568
63	Mesopontine cholinergic control over generalized non-convulsive seizures in a genetic model of absence epilepsy in the rat. <i>Neuroscience</i> , 1995 , 69, 1183-93	3.9	28
62	Protective effects of brain-derived neurotrophic factor on the development of hippocampal kindling in the rat. <i>NeuroReport</i> , 1995 , 6, 1937-41	1.7	101
61	Quiescence and hyporeactivity evoked by activation of cell bodies in the ventrolateral midbrain periaqueductal gray of the rat. <i>Experimental Brain Research</i> , 1994 , 99, 75-83	2.3	121
60	Endogenous control of epilepsy: the nigral inhibitory system. <i>Progress in Neurobiology</i> , 1994 , 42, 33-52	10.9	200
59	Convergence of deep somatic and visceral nociceptive information onto a discrete ventrolateral midbrain periaqueductal gray region. <i>Neuroscience</i> , 1994 , 61, 727-32	3.9	163
58	Nucleus basalis lesions suppress spike and wave discharges in rats with spontaneous absence-epilepsy. <i>Neuroscience</i> , 1994 , 59, 531-9	3.9	20
57	Effects of cholinergic drugs on genetic absence seizures in rats. <i>European Journal of Pharmacology</i> , 1993 , 234, 263-8	5.3	33
56	Reciprocal positive transfer between kindling of audiogenic seizures and electrical kindling of inferior colliculus. <i>Epilepsy Research</i> , 1993 , 15, 133-9	3	19
55	22-28 kHz ultrasonic vocalizations associated with defensive reactions in male rats do not result from fear or aversion. <i>Psychopharmacology</i> , 1993 , 111, 190-4	4.7	24
54	Involvement of intrathalamic GABAB neurotransmission in the control of absence seizures in the rat. <i>Neuroscience</i> , 1992 , 48, 87-93	3.9	217
53	The GABAA receptor complex in experimental absence seizures in rat: an autoradiographic study. <i>Neuroscience Letters</i> , 1992 , 140, 9-12	3.3	25
52	Dorsal tegmentum kindling in rats. <i>Neuroscience Letters</i> , 1992 , 134, 284-7	3.3	14
51	Longitudinal neuronal organization of defensive reactions in the midbrain periaqueductal gray region of the rat. <i>Experimental Brain Research</i> , 1992 , 90, 307-18	2.3	158
50	Opposite effects of pentylenetetrazol on self-defensive and submissive postures in the rat. <i>Psychopharmacology</i> , 1992 , 107, 457-60	4.7	2
49	Positive transfer of audiogenic kindling to electrical hippocampal kindling in rats. <i>Epilepsy Research</i> , 1992 , 11, 159-66	3	32

48	Are rats with genetic absence epilepsy behaviorally impaired?. Epilepsy Research, 1991, 9, 97-104	3	45
47	Lesions of noradrenergic neurons in rats with spontaneous generalized non-convulsive epilepsy. <i>Epilepsy Research</i> , 1991 , 9, 79-85	3	15
46	Opposite effects of agonist and inverse agonist ligands of benzodiazepine receptor on self-defensive and submissive postures in the rat. <i>Psychopharmacology</i> , 1991 , 103, 56-61	4.7	17
45	Quantitative analysis and computer simulation of oxytocin-neurophysin processing in the rat neurohypophysis. <i>Neurochemistry International</i> , 1991 , 19, 297-312	4.4	2
44	Intrathalamic injections of gamma-hydroxybutyric acid increase genetic absence seizures in rats. <i>Neuroscience Letters</i> , 1991 , 125, 19-21	3.3	21
43	Evidence for a critical role of GABAergic transmission within the thalamus in the genesis and control of absence seizures in the rat. <i>Brain Research</i> , 1991 , 545, 1-7	3.7	142
42	Emerging Principles of Organization of the Midbrain Periaqueductal Gray Matter 1991, 1-8		51
41	Midbrain Periaqueductal Gray Control of Defensive Behavior in the Cat and the Rat 1991 , 175-198		75
40	Suppression of spontaneous generalized non-convulsive seizures in the rat by microinjection of GABA antagonists into the superior colliculus. <i>Epilepsy Research</i> , 1990 , 5, 192-8	3	36
39	The GABAergic nigro-collicular pathway is not involved in the inhibitory control of audiogenic seizures in the rat. <i>Neuroscience Letters</i> , 1990 , 111, 269-74	3.3	23
38	Immediate effects of 14 non MAOI antidepressants in rats with spontaneous petit mal-like seizures. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1990 , 14, 261-70	5.5	5
37	Involvement of the nigral output pathways in the inhibitory control of the substantia nigra over generalized non-convulsive seizures in the rat. <i>Neuroscience</i> , 1990 , 39, 339-49	3.9	51
36	Potentiation of gamma-vinyl GABA (vigabatrin) effects by glycine. <i>European Journal of Pharmacology</i> , 1990 , 182, 109-15	5.3	32
35	Mapping of spontaneous spike and wave discharges in Wistar rats with genetic generalized non-convulsive epilepsy. <i>Brain Research</i> , 1990 , 523, 87-91	3.7	128
34	Interhemispheric desynchronization of spontaneous spike-wave discharges by corpus callosum transection in rats with petit mal-like epilepsy. <i>Epilepsy Research</i> , 1989 , 4, 8-13	3	39
33	Suppressive effects of intranigral injection of muscimol in three models of generalized non-convulsive epilepsy induced by chemical agents. <i>Brain Research</i> , 1989 , 498, 64-72	3.7	96
32	Anticonvulsant effect of muscimol injected into the thalamus of spontaneously epileptic Mongolian gerbils. <i>Brain Research</i> , 1989 , 487, 363-7	3.7	13
31	Characterization of pretentorial periaqueductal gray matter neurons mediating intraspecific defensive behaviors in the rat by microinjections of kainic acid. <i>Brain Research</i> , 1989 , 486, 121-32	3.7	89

(1986-1988)

30	Audiogenic seizures in Wistar rats before and after repeated auditory stimuli: clinical, pharmacological, and electroencephalographic studies. <i>Journal of Neural Transmission</i> , 1988 , 72, 235-4	4 4.3	47	
29	Relationship between analgesia and cardiovascular changes induced by electrical stimulation of the mesencephalic periaqueductal gray matter in the rat. <i>Brain Research</i> , 1988 , 451, 326-32	3.7	16	
28	Evidence that activation of GABA receptors in the substantia nigra suppresses spontaneous spike-and-wave discharges in the rat. <i>Brain Research</i> , 1988 , 448, 20-9	3.7	83	
27	Elicitation of intraspecific defence reactions in the rat from midbrain periaqueductal grey by microinjection of kainic acid, without neurotoxic effects. <i>Neuroscience Letters</i> , 1988 , 88, 291-6	3.3	77	
26	Effects of drugs affecting dopaminergic neurotransmission in rats with spontaneous petit mal-like seizures. <i>Neuropharmacology</i> , 1988 , 27, 269-74	5.5	70	
25	Effects of gamma-hydroxybutyrate and gamma-butyrolactone derivates on spontaneous generalized non-convulsive seizures in the rat. <i>Neuropharmacology</i> , 1988 , 27, 683-9	5.5	35	
24	Relationship between spike-wave discharges and vigilance levels in rats with spontaneous petit mal-like epilepsy. <i>Neuroscience Letters</i> , 1988 , 94, 187-91	3.3	69	
23	Selective increase of offensive behavior in the rat following intrahypothalamic 5,7-DHT-induced serotonin depletion. <i>Behavioural Brain Research</i> , 1988 , 29, 85-91	3.4	74	
22	Bidirectional effects of beta-carbolines in rats with spontaneous petit mal-like seizures. <i>Brain Research Bulletin</i> , 1987 , 19, 327-35	3.9	18	
21	Effects of drugs affecting noradrenergic neurotransmission in rats with spontaneous petit mal-like seizures. <i>European Journal of Pharmacology</i> , 1987 , 135, 397-402	5.3	57	
20	Diazepam dissociates the analgesic and aversive effects of periaqueductal gray stimulation in the rat. <i>Brain Research</i> , 1987 , 423, 395-8	3.7	33	
19	GABAergic modulation of the analgesic effects of morphine microinjected in the ventral periaqueductal gray matter of the rat. <i>Brain Research</i> , 1987 , 436, 223-8	3.7	140	
18	Spontaneous spike and wave discharges in thalamus and cortex in a rat model of genetic petit mal-like seizures. <i>Experimental Neurology</i> , 1987 , 96, 127-36	5.7	145	
17	Kindling of audiogenic seizures in Wistar rats: an EEG study. <i>Experimental Neurology</i> , 1987 , 97, 160-8	5.7	132	
16	Kindling of audiogenic seizures in the rat. International Journal of Neuroscience, 1987, 36, 167-76	2	31	
15	Involvement of brain opiate receptors in the immune-suppressive effect of morphine. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1986 , 83, 7114-7	11.5	172	
14	Parachlorophenylalanine-induced serotonin depletion increases offensive but not defensive aggression in male rats. <i>Physiology and Behavior</i> , 1986 , 36, 653-8	3.5	107	
13	Ontogeny of spontaneous petit mal-like seizures in Wistar rats. <i>Developmental Brain Research</i> , 1986 , 30, 85-87		44	

12	gamma-aminobutyric acid antagonist, into the midbrain periaqueductal gray matter. <i>Brain Research</i> , 1986 , 367, 87-95	3.7	87
11	Elicitation of conspecific attack or defense in the male rat by intraventricular injection of a GABA agonist or antagonist. <i>Physiology and Behavior</i> , 1985 , 35, 447-53	3.5	28
10	Identification of midbrain neurones mediating defensive behaviour in the rat by microinjections of excitatory amino acids. <i>Behavioural Brain Research</i> , 1985 , 15, 107-19	3.4	150
9	Gabaergic modulation of mouse-killing in the rat. <i>Psychopharmacology</i> , 1984 , 83, 367-72	4.7	13
8	Enhancement of spike and wave discharges by GABAmimetic drugs in rats with spontaneous petit-mal-like epilepsy. <i>Neuroscience Letters</i> , 1984 , 44, 91-4	3.3	163
7	Biphasic effects of Ro 15-1788 on spontaneous petit mal-like seizures in rats. <i>European Journal of Pharmacology</i> , 1984 , 102, 355-9	5.3	35
6	A model of chronic spontaneous petit mal-like seizures in the rat: comparison with pentylenetetrazol-induced seizures. <i>Epilepsia</i> , 1984 , 25, 326-31	6.4	170
5	Relationship between mousekilling and conspecific aggression in the male rat. <i>Aggressive Behavior</i> , 1983 , 9, 259-268	2.8	16
4	Induction of mouse-killing in the rat by intraventricular injection of a GABA-agonist. <i>Physiology and Behavior</i> , 1983 , 30, 383-8	3.5	26
3	A microcomputer method for behavioral data acquisition and subsequent analysis. <i>Pharmacology Biochemistry and Behavior</i> , 1983 , 19, 729-32	3.9	21
2	Spontaneous paroxysmal electroclinical patterns in rat: a model of generalized non-convulsive epilepsy. <i>Neuroscience Letters</i> , 1982 , 33, 97-101	3.3	242
1	GABAergic influences on defensive fighting in rats. <i>Pharmacology Biochemistry and Behavior</i> , 1982 , 17, 451-6	3.9	42