Alan A Montgomery

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7396114/publications.pdf

Version: 2024-02-01

205 papers 19,130 citations

³⁸⁷⁴² 50 h-index

127 g-index

218 all docs

218 docs citations

times ranked

218

34680 citing authors

#	Article	IF	CITATIONS
1	Lactic acid gel versus metronidazole for recurrent bacterial vaginosis in women aged 16 years and over: the VITA RCT. Health Technology Assessment, 2022, 26, 1-170.	2.8	11
2	The FEED1 trial: protocol for a randomised controlled trial of full milk feeds versus intravenous fluids with gradual feeding for preterm infants (30–33 weeks gestational age). Trials, 2022, 23, 64.	1.6	4
3	The COVID-19 pandemic has highlighted the need to invest in care home research infrastructure. Age and Ageing, 2022, 51, .	1.6	6
4	Should we adjudicate outcomes in stroke trials? A systematic review. International Journal of Stroke, 2022, , 174749302210946.	5.9	1
5	STAndardised Diagnostic Assessment for children and young people with emotional difficulties (STADIA): protocol for a multicentre randomised controlled trial. BMJ Open, 2022, 12, e053043.	1.9	2
6	Using the Vitiligo Noticeability Scale in clinical trials: construct validity, interpretability, reliability and acceptability. British Journal of Dermatology, 2022, 187, 548-556.	1.5	4
7	Dexamethasone in Hospitalized Patients with Covid-19. New England Journal of Medicine, 2021, 384, 693-704.	27.0	8,063
8	Randomized controlled trial of topical corticosteroid and homeâ€based narrowband ultraviolet B for active and limited vitiligo: results of the HIâ€Light Vitiligo Trial*. British Journal of Dermatology, 2021, 184, 828-839.	1.5	25
9	An economic evaluation of the randomized controlled trial of topical corticosteroid and homeâ€based narrowband ultraviolet B for active and limited vitiligo (the Hlâ€Light Vitiligo Trial)*. British Journal of Dermatology, 2021, 184, 840-848.	1.5	6
10	Promoting Independence in Dementia (PRIDE): A Feasibility Randomized Controlled Trial. Clinical Interventions in Aging, 2021, Volume 16, 363-378.	2.9	7
11	Regulatory Approach for Transitioning from Gamma Ray to X-ray Radiation Sterilization. Biomedical Instrumentation and Technology, 2021, 55, 58-66.	0.4	5
12	Protocol for surgical and non-surgical treatment for metacarpal shaft fractures in adults: an observational feasibility study. BMJ Open, 2021, 11, e046913.	1.9	2
13	Association Between Administration of IL-6 Antagonists and Mortality Among Patients Hospitalized for COVID-19. JAMA - Journal of the American Medical Association, 2021, 326, 499.	7.4	498
14	Developing a core outcome set for hand fractures and joint injuries in adults: a systematic review. Journal of Hand Surgery: European Volume, 2021, 46, 488-495.	1.0	6
15	Statistical reanalysis of vascular event outcomes in primary and secondary vascular prevention trials. BMC Medical Research Methodology, 2021, 21, 218.	3.1	4
16	POSNOCâ€"POsitive Sentinel NOde: adjuvant therapy alone versus adjuvant therapy plus Clearance or axillary radiotherapy: a randomised controlled trial of axillary treatment in women with early-stage breast cancer who have metastases in one or two sentinel nodes. BMJ Open, 2021, 11, e054365.	1.9	29
17	Cognitive rehabilitation for attention and memory in people with multiple sclerosis: a randomized controlled trial (CRAMMS). Clinical Rehabilitation, 2020, 34, 229-241.	2.2	10
18	Effect of Hydroxychloroquine in Hospitalized Patients with Covid-19. New England Journal of Medicine, 2020, 383, 2030-2040.	27.0	1,013

#	Article	IF	CITATIONS
19	Lopinavir–ritonavir in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. Lancet, The, 2020, 396, 1345-1352.	13.7	569
20	An algorithm for diagnosing IgEâ€mediated food allergy in study participants who do not undergo food challenge. Clinical and Experimental Allergy, 2020, 50, 334-342.	2.9	14
21	Two-by-two factorial randomised study within a trial (SWAT) to evaluate strategies for follow-up in a randomised prevention trial. Trials, 2020, 21, 529.	1.6	8
22	A Systematic Review of Treatment Interventions for Metacarpal Shaft Fractures in Adults. Hand, 2020, , 155894472097436.	1.2	3
23	Optimising recruitment to the HAND-1 RCT feasibility study: integration of the QuinteT Recruitment Intervention (QRI). Pilot and Feasibility Studies, 2020, 6, 173.	1.2	2
24	Outcome assessment by central adjudicators in randomised stroke trials: Simulation of differential and non-differential misclassification. European Stroke Journal, 2020, 5, 174-183.	5.5	6
25	Protocol: The Lacunar Intervention Trial 2 (LACI-2). A trial of two repurposed licenced drugs to prevent progression of cerebral small vessel disease. European Stroke Journal, 2020, 5, 297-308.	5.5	22
26	Cost-benefit of outcome adjudication in nine randomised stroke trials. Clinical Trials, 2020, 17, 576-580.	1.6	5
27	Data sharing: experience of accessing individual patient data from completed randomised controlled trials in vascular and cognitive medicine. BMJ Open, 2020, 10, e038765.	1.9	4
28	The feasibility and acceptability of an early intervention in primary care to prevent chronic fatigue syndrome (CFS) in adults: randomised controlled trial. Pilot and Feasibility Studies, 2020, 6, 65.	1.2	3
29	Adding evidence of the effects of treatments into relevant Wikipedia pages: a randomised trial. BMJ Open, 2020, 10, e033655.	1.9	8
30	Daily emollient during infancy for prevention of eczema: the BEEP randomised controlled trial. Lancet, The, 2020, 395, 962-972.	13.7	178
31	Systematic review and network meta-analysis with individual participant data on cord management at preterm birth (iCOMP): study protocol. BMJ Open, 2020, 10, e034595.	1.9	16
32	Group cognitive rehabilitation to reduce the psychological impact of multiple sclerosis on quality of life: the CRAMMS RCT. Health Technology Assessment, 2020, 24, 1-182.	2.8	14
33	Home-based narrowband UVB, topical corticosteroid or combination for children and adults with vitiligo: HI-Light Vitiligo three-arm RCT. Health Technology Assessment, 2020, 24, 1-128.	2.8	18
34	TANDEM TRIAL: a factorial randomised controlled trial of dose and review schedule of bevacizumab (Avastin) for neovascular macular degeneration in the East Midlands. BMJ Open Ophthalmology, 2020, 5, e000588.	1.6	0
35	Outcome Assessment by Central Adjudicators Versus Site Investigators in Stroke Trials. Stroke, 2019, 50, 2187-2196.	2.0	13
36	Gentamicin, azithromycin and ceftriaxone in the treatment of gonorrhoea: the relationship between antibiotic MIC and clinical outcome. Journal of Antimicrobial Chemotherapy, 2019, 75, 449-457.	3.0	14

#	Article	IF	CITATIONS
37	Prehospital Transdermal Glyceryl Trinitrate for Ultra-Acute Intracerebral Hemorrhage. Stroke, 2019, 50, 3064-3071.	2.0	36
38	Short email with attachment versus long email without attachment when contacting authors to request unpublished data for a systematic review: a nested randomised trial. BMJ Open, 2019, 9, e025273.	1.9	7
39	Clinical and cost effectiveness of memory rehabilitation following traumatic brain injury: a pragmatic cluster randomized controlled trial. Clinical Rehabilitation, 2019, 33, 1171-1184.	2.2	9
40	Effect of a stepped-care intervention delivered by lay health workers on major depressive disorder among primary care patients in Nigeria (STEPCARE): a cluster-randomised controlled trial. The Lancet Global Health, 2019, 7, e951-e960.	6.3	47
41	Gentamicin compared with ceftriaxone for the treatment of gonorrhoea (G-ToG): a randomised non-inferiority trial. Lancet, The, 2019, 393, 2511-2520.	13.7	84
42	Full Milk Feeds from Day 1. Neonatology, 2019, 116, 67-67.	2.0	0
43	Prehospital transdermal glyceryl trinitrate in patients with ultra-acute presumed stroke (RIGHT-2): an ambulance-based, randomised, sham-controlled, blinded, phase 3 trial. Lancet, The, 2019, 393, 1009-1020.	13.7	119
44	High- versus low-intensity interventions for perinatal depression delivered by non-specialist primary maternal care providers in Nigeria: cluster randomised controlled trial (the EXPONATE trial). British Journal of Psychiatry, 2019, 215, 528-535.	2.8	38
45	Day of the week to tweet: a randomised controlled trial. BMJ Open, 2019, 9, e025380.	1.9	2
46	Integrating quantitative and qualitative data and findings when undertaking randomised controlled trials. BMJ Open, 2019, 9, e032081.	1.9	42
47	Metronidazole versus lactic acid for treating bacterial vaginosis (VITA): protocol for a randomised controlled trial to assess the clinical and cost effectiveness of topical lactic acid gel for treating second and subsequent episodes of bacterial vaginosis. Trials, 2019, 20, 648.	1.6	8
48	Promoting Independence in Dementia (PRIDE): protocol for a feasibility randomised controlled trial. Trials, 2019, 20, 709.	1.6	8
49	The QuinteT Recruitment Intervention supported five randomized trials to recruit to target: a mixed-methods evaluation. Journal of Clinical Epidemiology, 2019, 106, 108-120.	5.0	49
50	A Validation and Calibration Process for Self-reported Tobacco Use With Participants' Cotinine Levels: An Example From the Building Blocks Trial. Nicotine and Tobacco Research, 2019, 21, 1660-1664.	2.6	0
51	Ambulance-delivered transdermal glyceryl trinitrate versus sham for ultra-acute stroke: Rationale, design and protocol for the Rapid Intervention with Glyceryl trinitrate in Hypertensive stroke Trial-2 (RIGHT-2) trial (ISRCTN26986053). International Journal of Stroke, 2019, 14, 191-206.	5.9	20
52	Follow-on RifAximin for the Prevention of recurrence following standard treatment of Infection with <i>Clostridium Difficile</i> (RAPID): a randomised placebo controlled trial. Gut, 2019, 68, 1224-1231.	12.1	23
53	A group memory rehabilitation programme for people with traumatic brain injuries: the ReMemBrIn RCT. Health Technology Assessment, 2019, 23, 1-194.	2.8	11
54	Gentamicin as an alternative to ceftriaxone in the treatment of gonorrhoea: the G-TOG non-inferiority RCT. Health Technology Assessment, 2019, 23, 1-104.	2.8	4

#	Article	IF	CITATIONS
55	Eicosapentaenoic acid and/or aspirin for preventing colorectal adenomas during colonoscopic surveillance in the NHS Bowel Cancer Screening Programme: the seAFOod RCT. Efficacy and Mechanism Evaluation, 2019, 6, 1-154.	0.7	7
56	The Clinical Effectiveness and Cost-Effectiveness of Lamotrigine in Borderline Personality Disorder: A Randomized Placebo-Controlled Trial. American Journal of Psychiatry, 2018, 175, 756-764.	7.2	78
57	Antiplatelet therapy with aspirin, clopidogrel, and dipyridamole versus clopidogrel alone or aspirin and dipyridamole in patients with acute cerebral ischaemia (TARDIS): a randomised, open-label, phase 3 superiority trial. Lancet, The, 2018, 391, 850-859.	13.7	125
58	Clinical and cost-effectiveness of the Lightning Process in addition to specialist medical care for paediatric chronic fatigue syndrome: randomised controlled trial. Archives of Disease in Childhood, 2018, 103, 155-164.	1.9	29
59	Statistical analysis plan for the â€~Rapid Intervention with Glyceryl trinitrate in Hypertensive stroke Trial-2 (RIGHT-2)'. European Stroke Journal, 2018, 3, 193-196.	5.5	8
60	Eicosapentaenoic acid and aspirin, alone and in combination, for the prevention of colorectal adenomas (seAFOod Polyp Prevention trial): a multicentre, randomised, double-blind, placebo-controlled, 2â€^×â€^2 factorial trial. Lancet, The, 2018, 392, 2583-2594.	13.7	102
61	Home interventions and light therapy for the treatment of vitiligo (HI-Light Vitiligo Trial): study protocol for a randomised controlled trial. BMJ Open, 2018, 8, e018649.	1.9	8
62	Central adjudication of serious adverse events did not affect trial's safety results: Data from the Efficacy of Nitric Oxide in Stroke (ENOS) trial. PLoS ONE, 2018, 13, e0208142.	2.5	6
63	Monitoring performance of sites within multicentre randomised trials: a systematic review of performance metrics. Trials, 2018, 19, 562.	1.6	9
64	Central masked adjudication of stroke diagnosis at trial entry offered no advantage over diagnosis by local clinicians: Secondary analysis and simulation. Contemporary Clinical Trials Communications, 2018, 12, 176-181.	1.1	5
65	Doing trials within trials: a qualitative study of stakeholder views on barriers and facilitators to the routine adoption of methodology research in clinical trials. Trials, 2018, 19, 481.	1.6	6
66	Development of a standardised set of metrics for monitoring site performance in multicentre randomised trials: a Delphi study. Trials, 2018, 19, 557.	1.6	19
67	Lamotrigine for people with borderline personality disorder: a RCT. Health Technology Assessment, 2018, 22, 1-68.	2.8	24
68	Triple versus guideline antiplatelet therapy to prevent recurrence after acute ischaemic stroke or transient ischaemic attack: the TARDIS RCT. Health Technology Assessment, 2018, 22, 1-76.	2.8	8
69	Large multi-centre pilot randomized controlled trial testing a low-cost, tailored, self-help smoking cessation text message intervention for pregnant smokers (MiQuit). Addiction, 2017, 112, 1238-1249.	3.3	66
70	Psychoeducation and Problem Solving (PEPS) Therapy for Adults With Personality Disorder: A Pragmatic Randomized-Controlled Trial. Journal of Personality Disorders, 2017, 31, 810-826.	1.4	14
71	Investigating the effect of independent, blinded digital image assessment on the STOP GAP trial. Trials, 2017, 18, 53.	1.6	2
72	Guidelines for the Content of Statistical Analysis Plans in Clinical Trials. JAMA - Journal of the American Medical Association, 2017, 318, 2337.	7.4	290

#	Article	IF	CITATIONS
73	Silk garments plus standard care compared with standard care for treating eczema in children: A randomised, controlled, observer-blind, pragmatic trial (CLOTHES Trial). PLoS Medicine, 2017, 14, e1002280.	8.4	41
74	Effectiveness and cost-effectiveness of daily all-over-body application of emollient during the first year of life for preventing atopic eczema in high-risk children (The BEEP trial): protocol for a randomised controlled trial. Trials, 2017, 18, 343.	1.6	56
75	A feasibility study investigating the acceptability and design of a multicentre randomised controlled trial of needle fasciotomy versus limited fasciectomy for the treatment of Dupuytren's contractures of the fingers (HAND-1): study protocol for a randomised controlled trial. Trials, 2017, 18, 392.	1.6	9
76	Randomised controlled trial of silk therapeutic garments for the management of atopic eczema in children: the CLOTHES trial. Health Technology Assessment, 2017, 21, 1-260.	2.8	24
77	Evaluation of the effectiveness and cost-effectiveness of lightweight fibreglass heel casts in the management of ulcers of the heel in diabetes: a randomised controlled trial. Health Technology Assessment, 2017, 21, 1-92.	2.8	15
78	An evidence-based approach to the use of telehealth in long-term health conditions: development of an intervention and evaluation through pragmatic randomised controlled trials in patients with depression or raised cardiovascular risk. Programme Grants for Applied Research, 2017, 5, 1-468.	1.0	4
79	The Community In-reach Rehabilitation and Care Transition (CIRACT) clinical and cost-effectiveness randomisation controlled trial in older people admitted to hospital as an acute medical emergency. Age and Ageing, 2016, 46, 26-32.	1.6	15
80	Effectiveness of a Nurse-Led Intensive Home-Visitation Program for First-Time Teenage Mothers (Building Blocks). Obstetrical and Gynecological Survey, 2016, 71, 263-265.	0.4	0
81	Gentamicin versus ceftriaxone for the treatment of gonorrhoea (G-TOG trial): study protocol for a randomised trial. Trials, 2016, 17, 558.	1.6	10
82	Cost-effectiveness of telehealth for patients with depression: evidence from the Healthlines randomised controlled trial. BJPsych Open, 2016, 2, 262-269.	0.7	19
83	Effectiveness of an integrated telehealth service for patients with depression: a pragmatic randomised controlled trial of a complex intervention. Lancet Psychiatry,the, 2016, 3, 515-525.	7.4	73
84	Influence of school-related factors on smoking among Chilean adolescents: a cross-sectional multilevel study. BMC Pediatrics, 2016, 16, 79.	1.7	8
85	Anxiety, depression and risk of cannabis use: Examining the internalising pathway to use among Chilean adolescents. Drug and Alcohol Dependence, 2016, 166, 109-115.	3.2	33
86	Validation of the Vitiligo Noticeability Scale: a patient-reported outcome measure of vitiligo treatment success. British Journal of Dermatology, 2016, 174, 386-394.	1.5	34
87	Indicated school-based intervention to improve depressive symptoms among at risk Chilean adolescents: a randomized controlled trial. BMC Psychiatry, 2016, 16, 276.	2.6	24
88	Cost-effectiveness of telehealth for patients with raised cardiovascular disease risk: evidence from the Healthlines randomised controlled trial. BMJ Open, 2016, 6, e012352.	1.9	25
89	Telehealth for patients at high risk of cardiovascular disease: pragmatic randomised controlled trial. BMJ, The, 2016, 353, i2647.	6.0	50
90	RAPID INTERVENTION WITH GLYCERYL TRINITRATE IN HYPERTENSIVE STROKE TRIAL-2 (RIGHT-2): SAFETY AND EFFICACY OF TRANSDERMAL GLYCERYL TRINITRATE, A NITRIC OXIDE DONOR. Emergency Medicine Journal, 2016, 33, e13.2-e14.	1.0	2

#	Article	IF	CITATIONS
91	Increasing follow-up questionnaire response rates in a randomized controlled trial of telehealth for depression: three embedded controlled studies. Trials, 2016, 17, 107.	1.6	10
92	Effectiveness of a nurse-led intensive home-visitation programme for first-time teenage mothers (Building Blocks): a pragmatic randomised controlled trial. Lancet, The, 2016, 387, 146-155.	13.7	217
93	A mechanistic multicentre, parallel group, randomised placebo-controlled trial of mesalazine for the treatment of IBS with diarrhoea (IBS-D). Gut, 2016, 65, 91-99.	12.1	85
94	Comparing the cost-effectiveness and clinical effectiveness of a new community in-reach rehabilitation service with the cost-effectiveness and clinical effectiveness of an established hospital-based rehabilitation service for older people: a pragmatic randomised controlled trial with microcost and qualitative analysis – the Community In-reach Rehabilitation And Care Transition	1.4	4
95	CIRACT) study. Health Services and Delivers Research, 2016, 4, 1-90. Psychoeducation with problem-solving (PEPS) therapy for adults with personality disorder: a pragmatic randomised controlled trial to determine the clinical effectiveness and cost-effectiveness of a manualised intervention to improve social functioning. Health Technology Assessment, 2016, 20, 1-250.	2.8	11
96	Relationship between Expectation Management and Client Retention in Online Cognitive Behavioural Therapy. Behavioural and Cognitive Psychotherapy, 2015, 43, 732-743.	1.2	10
97	Patient–doctor continuity and diagnosis of cancer: electronic medical records study in general practice. British Journal of General Practice, 2015, 65, e305-e311.	1.4	24
98	To tweet or not to tweet about schizophrenia systematic reviews (TweetSz): study protocol for a randomised controlled trial. BMJ Open, 2015, 5, e007695.	1.9	2
99	Cognitive Rehabilitation for Attention and Memory in people with Multiple Sclerosis: study protocol for a randomised controlled trial (CRAMMS). Trials, 2015, 16, 556.	1.6	13
100	Lamotrigine versus inert placebo in the treatment of borderline personality disorder: study protocol for a randomized controlled trial and economic evaluation. Trials, 2015, 16, 308.	1.6	9
101	The provision of NHS health checks in a community setting: an ethnographic account. BMC Health Services Research, 2015, 15, 546.	2.2	17
102	Who does not participate in telehealth trials and why? A cross-sectional survey. Trials, 2015, 16, 258.	1.6	44
103	A multi-centre, parallel group superiority trial of silk therapeutic clothing compared to standard care for the management of eczema in children (CLOTHES Trial): study protocol for a randomised controlled trial. Trials, 2015, 16, 390.	1.6	10
104	TElehealth in CHronic disease: mixed-methods study to develop the TECH conceptual model for intervention design and evaluation. BMJ Open, 2015, 5, e006448-e006448.	1.9	66
105	Decisions, choice and shared decision making in antenatal clinics: An observational study. Patient Education and Counseling, 2015, 98, 1106-1111.	2.2	8
106	Randomised trial of the fascia iliaca block versus the  3-in-1' block for femoral neck fractures in the emergency department. Emergency Medicine Journal, 2015, 32, 685-689.	1.0	40
107	Rehabilitation of Memory following Brain Injury (ReMemBrIn): study protocol for a randomised controlled trial. Trials, 2015, 16, 6.	1.6	9
108	Increasing children's physical activity through a teaching-assistant led extracurricular intervention: process evaluation of the action 3:30 randomised feasibility trial. BMC Public Health, 2015, 15, 156.	2.9	18

#	Article	IF	CITATIONS
109	Pilot study to evaluate a tailored text message intervention for pregnant smokers (MiQuit): study protocol for a randomised controlled trial. Trials, 2015, 16, 29.	1.6	17
110	The Community In-Reach and Care Transition (CIRACT) clinical and cost-effectiveness study: study protocol for a randomised controlled trial. Trials, 2015, 16, 41.	1.6	3
111	Comparing different dosing regimens of bevacizumab in the treatment of neovascular macular degeneration: study protocol for a randomised controlled trial. Trials, 2015, 16, 85.	1.6	7
112	A pilot randomized controlled trial of a stepped care intervention package for depression in primary care in Nigeria. BMC Psychiatry, 2015, 15, 96.	2.6	24
113	The Association Between School Bonding and Smoking Amongst Chilean Adolescents. Substance Abuse, 2015, 36, 515-523.	2.3	5
114	A cluster randomized clinical trial of a stepped care intervention for depression in primary care (STEPCARE)- study protocol. BMC Psychiatry, 2015, 15, 148.	2.6	14
115	Expanding care for perinatal women with depression (EXPONATE): study protocol for a randomized controlled trial of an intervention package for perinatal depression in primary care. BMC Psychiatry, 2015, 136.	2.6	28
116	Peer victimization during adolescence: concurrent and prospective impact on symptoms of depression and anxiety. Anxiety, Stress and Coping, 2015, 28, 105-120.	2.9	121
117	Blinded randomised controlled trial of low-dose Adjuvant Steroids in Adults admitted to hospital with Pandemic influenza (ASAP): a trial â€`in hibernation', ready for rapid activation. Health Technology Assessment, 2015, 19, 1-78.	2.8	15
118	Antibiotic Prescribing in Primary Care and Antimicrobial Resistance in Patients Admitted to Hospital with Urinary Tract Infection: A Controlled Observational Pilot Study. Antibiotics, 2014, 3, 29-38.	3.7	11
119	Randomised feasibility trial of a teaching assistant led extracurricular physical activity intervention for 9 to 11 year olds: Action 3:30. International Journal of Behavioral Nutrition and Physical Activity, 2014, 11, 114.	4.6	29
120	Changing eating behaviours to treat childhood obesity in the community using Mandolean: the Community Mandolean randomised controlled trial (ComMando) – a pilot study. Health Technology Assessment, 2014, 18, 1-75.	2.8	37
121	Effectiveness of PhysioDirect telephone assessment and advice services for patients with musculoskeletal problems:. British Journal of Sports Medicine, 2014, 48, 1391-1391.	6.7	3
122	Costâ€effectiveness of classroomâ€based cognitive behaviour therapy in reducing symptoms of depression in adolescents: a trialâ€based analysis. Journal of Child Psychology and Psychiatry and Allied Disciplines, 2014, 55, 1390-1397.	5 . 2	23
123	Health services changes: is a run-in period necessary before evaluation in randomised clinical trials?. Trials, 2014, 15, 41.	1.6	4
124	Effectiveness and cost-effectiveness of a telehealth intervention to support the management of long-term conditions: study protocol for two linked randomized controlled trials. Trials, 2014, 15, 36.	1.6	33
125	Factors associated with the development of self-harm amongst a socio-economically deprived cohort of adolescents in Santiago, Chile. Social Psychiatry and Psychiatric Epidemiology, 2014, 49, 629-637.	3.1	20
126	Instrumental delivery and ultrasound: a multicentre randomised controlled trial of ultrasound assessment of the fetal head position versus standard care as an approach to prevent morbidity at instrumental delivery. BJOG: an International Journal of Obstetrics and Gynaecology, 2014, 121, 1029-1038.	2.3	115

#	Article	IF	CITATIONS
127	School travel mode, parenting practices and physical activity among UK Year 5 and 6 children. BMC Public Health, 2014, 14, 370.	2.9	8
128	Sedentary time and markers of inflammation in people with newly diagnosed type 2 diabetes. Nutrition, Metabolism and Cardiovascular Diseases, 2014, 24, 956-962.	2.6	42
129	Associations Between Physical Activity Parenting Practices and Adolescent Girls' Self-Perceptions and Physical Activity Intentions. Journal of Physical Activity and Health, 2014, 11, 734-740.	2.0	10
130	Are People With Chronic Diseases Interested in Using Telehealth? A Cross-Sectional Postal Survey. Journal of Medical Internet Research, 2014, 16, e123.	4.3	103
131	Action 3:30: protocol for a randomized feasibility trial of a teaching assistant led extracurricular physical activity intervention. Trials, 2013, 14, 122.	1.6	9
132	Detecting depression among adolescents in Santiago, Chile: sex differences. BMC Psychiatry, 2013, 13, 122.	2.6	27
133	The effectiveness and cost-effectiveness of the Family Nurse Partnership home visiting programme for first time teenage mothers in England: a protocol for the Building Blocks randomised controlled trial. BMC Pediatrics, 2013, 13, 114.	1.7	36
134	Could a brief assessment of negative emotions and self-esteem identify adolescents at current and future risk of self-harm in the community? A prospective cohort analysis. BMC Public Health, 2013, 13, 604.	2.9	10
135	A randomized controlled trial of eicosapentaenoic acid and/or aspirin for colorectal adenoma prevention during colonoscopic surveillance in the NHS Bowel Cancer Screening Programme (The) Tj ETQq1 1 0.78	84314 rgB 1.6	ST ₃ Overlo <mark>ck</mark>
136	Comparing specialist medical care with specialist medical care plus the Lightning Process® for chronic fatigue syndrome or myalgic encephalomyelitis (CFS/ME): study protocol for a randomised controlled trial (SMILE Trial). Trials, 2013, 14, 444.	1.6	12
137	The feasibility and acceptability of conducting a trial of specialist medical care and the Lightning Process in children with chronic fatigue syndrome: feasibility randomized controlled trial (SMILE) Tj ETQq1 1 0.784	13.164 rgBT	/ © verlock
138	Self-harm in young adolescents (12–16Âyears): onset and short-term continuation in a community sample. BMC Psychiatry, 2013, 13, 328.	2.6	75
139	A pragmatic randomised controlled trial of â€PhysioDirect' telephone assessment and advice services for patients with musculoskeletal problems: economic evaluation. BMJ Open, 2013, 3, e003406.	1.9	14
140	Effectiveness of PhysioDirect telephone assessment and advice services for patients with musculoskeletal problems: pragmatic randomised controlled trial. BMJ, The, 2013, 346, f43-f43.	6.0	56
141	An evaluation of the impact and costs of three strategies used to recruit acutely unwell young children to a randomised controlled trial in primary care. Clinical Trials, 2013, 10, 593-603.	1.6	3
142	Women and their partners' preferences for Down's syndrome screening tests: a discrete choice experiment. Prenatal Diagnosis, 2013, 33, 449-456.	2.3	35
143	Republished research: Facilitated physical activity as a treatment for depressed adults: randomised controlled trial:. British Journal of Sports Medicine, 2013, 47, 629-629.	6.7	1
144	School Intervention to Improve Mental Health of Students in Santiago, Chile. JAMA Pediatrics, 2013, 167, 1004.	6.2	58

#	Article	IF	CITATIONS
145	Depression Symptom Trajectories and Associated Risk Factors among Adolescents in Chile. PLoS ONE, 2013, 8, e78323.	2.5	20
146	A pragmatic randomised controlled trial of the effectiveness and cost-effectiveness of ‬PhysioDirect' telephone assessment and advice services for physiotherapy. Health Technology Assessment, 2013, 17, 1-157.	2.8	52
147	A cluster randomised controlled trial to determine the clinical effectiveness and cost-effectiveness of classroom-based cognitive–behavioural therapy (CBT) in reducing symptoms of depression in high-risk adolescents. Health Technology Assessment, 2013, 17, vii-xvii, 1-109.	2.8	51
148	Classroom based cognitive behavioural therapy in reducing symptoms of depression in high risk adolescents: pragmatic cluster randomised controlled trial. BMJ, The, 2012, 345, e6058-e6058.	6.0	92
149	Facilitated physical activity as a treatment for depressed adults: randomised controlled trial. BMJ, The, 2012, 344, e2758-e2758.	6.0	163
150	Study Protocol. IDUS – Instrumental delivery & Ditrasound. A multi-centre randomised controlled trial of ultrasound assessment of the fetal head position versus standard care as an approach to prevent morbidity at instrumental delivery. BMC Pregnancy and Childbirth, 2012, 12, 95.	2.4	9
151	Bristol Girls Dance Project Feasibility Trial: outcome and process evaluation results. International Journal of Behavioral Nutrition and Physical Activity, 2012, 9, 83.	4.6	39
152	School connectedness, peer attachment, and selfâ€esteem as predictors ofÂadolescent depression. Journal of Adolescence, 2012, 35, 1061-1067.	2.4	138
153	Effect of antibiotic prescribing in primary care on meticillin-resistant Staphylococcus aureus carriage in community-resident adults: a controlled observational study. International Journal of Antimicrobial Agents, 2012, 39, 135-141.	2. 5	21
154	A qualitative investigation of the decisionâ€making process of couples considering prenatal screening for Down syndrome. Prenatal Diagnosis, 2012, 32, 57-63.	2.3	24
155	A pragmatic randomised controlled trial to evaluate the cost-effectiveness of a physical activity intervention as a treatment for depression: the treating depression with physical activity (TREAD) trial Health Technology Assessment, 2012, 16, 1-164, iii-iv.	2.8	53
156	Diet or diet plus physical activity versus usual care in patients with newly diagnosed type 2 diabetes: the Early ACTID randomised controlled trial. Lancet, The, 2011, 378, 129-139.	13.7	249
157	Diet or diet plus physical activity in patients with early type 2 diabetes – Authors' reply. Lancet, The, 2011, 378, 2067-2068.	13.7	0
158	Preferences for mode of delivery after previous caesarean section: what do women want, what do they get and how do they value outcomes?. Health Expectations, 2011, 14, 397-404.	2.6	13
159	Reporting of factorial trials of complex interventions in community settings: a systematic review. Trials, 2011, 12, 179.	1.6	23
160	School-based intervention to improve the mental health of low-income, secondary school students in Santiago, Chile (YPSA): study protocol for a randomized controlled trial. Trials, 2011, 12, 49.	1.6	15
161	The frequency distribution of presenting symptoms in children aged six months to six years to primary care. Primary Health Care Research and Development, 2011, 12, 123-134.	1.2	36
162	Oxytocin bolus versus oxytocin bolus and infusion for control of blood loss at elective caesarean section: double blind, placebo controlled, randomised trial. BMJ: British Medical Journal, 2011, 343, d4661-d4661.	2.3	74

#	Article	IF	Citations
163	Medicine dosing by weight in the home: can parents accurately weigh preschool children? A method comparison study. Archives of Disease in Childhood, 2011, 96, 1187-1190.	1.9	4
164	Epidemiology and impact of multimorbidity in primary care: a retrospective cohort study. British Journal of General Practice, 2011, 61, e12-e21.	1.4	647
165	Physical activity as a treatment for depression: the TREAD randomised trial protocol. Trials, 2010, 11, 105.	1.6	16
166	Protocol for a randomised controlled trial of a school based cognitive behaviour therapy (CBT) intervention to prevent depression in high risk adolescents (PROMISE). Trials, 2010, 11, 114.	1.6	16
167	Patients' experience and satisfaction in primary care: secondary analysis using multilevel modelling. BMJ: British Medical Journal, 2010, 341, c5004-c5004.	2.3	125
168	Economic Evaluation of the DiAMOND Randomized Trial: Cost and Outcomes of 2 Decision Aids for Mode of Delivery among Women with a Previous Cesarean Section. Medical Decision Making, 2010, 30, 453-463.	2.4	14
169	Study Protocol. ECSSIT – Elective Caesarean Section Syntocinon® Infusion Trial. A multi-centre randomised controlled trial of oxytocin (Syntocinon®) 5 IU bolus and placebo infusion versus oxytocin 5 IU bolus and 40 IU infusion for the control of blood loss at elective caesarean section. BMC Pregnancy and Childbirth, 2009, 9, 36.	2.4	10
170	'PhysioDirect' telephone assessment and advice services for physiotherapy: protocol for a pragmatic randomised controlled trial. BMC Health Services Research, 2009, 9, 136.	2.2	17
171	The external validity of published randomized controlled trials in primary care. BMC Family Practice, 2009, 10, 5.	2.9	33
172	Healthcare professionals' views on two computerâ€based decision aids for women choosing mode of delivery after previous caesarean section: a qualitative study. BJOG: an International Journal of Obstetrics and Gynaecology, 2009, 116, 906-914.	2.3	15
173	How should continuity of care in primary health care be assessed? British Journal of General Practice, 2009, 59, e134-e141.	1.4	85
174	Paracetamol plus ibuprofen for the treatment of fever in children (PITCH): randomised controlled trial. BMJ: British Medical Journal, 2008, 337, a1302-a1302.	2.3	124
175	Paracetamol plus ibuprofen for the treatment of fever in children (PITCH): economic evaluation of a randomised controlled trial. BMJ: British Medical Journal, 2008, 337, a1490-a1490.	2.3	23
176	Two decision aids for mode of delivery among women with previous caesarean section: randomised controlled trial. BMJ: British Medical Journal, 2007, 334, 1305.	2.3	140
177	The impact of co-located NHS walk-in centres on emergency departments. Emergency Medicine Journal, 2007, 24, 265-269.	1.0	56
178	Comparing care at walk-in centres and at accident and emergency departments: an exploration of patient choice, preference and satisfaction. Emergency Medicine Journal, 2007, 24, 260-264.	1.0	28
179	Common mental disorders and the built environment in Santiago, Chile. British Journal of Psychiatry, 2007, 190, 394-401.	2.8	67
180	Decision-making about mode of delivery after previous caesarean section: development and piloting of two computer-based decision aids. Health Expectations, 2007, 10, 161-172.	2.6	33

#	Article	IF	CITATIONS
181	Impact of Advanced Access on access, workload, and continuity: controlled before-and-after and simulated-patient study. British Journal of General Practice, 2007, 57, 608-14.	1.4	36
182	Does Advanced Access improve access to primary health care? Questionnaire survey of patients. British Journal of General Practice, 2007, 57, 615-21.	1.4	40
183	Adherence to antihypertensive medication assessed by self-report was associated with electronic monitoring compliance. Journal of Clinical Epidemiology, 2006, 59, 650-651.	5.0	54
184	Women's experience of decision making about mode of delivery after a previous caesarean section: the role of health professionals and information about health risks. BJOG: an International Journal of Obstetrics and Gynaecology, 2006, 113, 1438-1445.	2.3	83
185	Preference-Based Antithrombotic Therapy in Atrial Fibrillation: Implications for Clinical Decision Making. Medical Decision Making, 2005, 25, 548-559.	2.4	67
186	The relationship between primary care antibiotic prescribing and bacterial resistance in adults in the community: a controlled observational study using individual patient data. Journal of Antimicrobial Chemotherapy, 2005, 56, 146-153.	3.0	60
187	A Pilot, Randomized, Double-Blinded, Placebo-Controlled Trial of Individualized Homeopathy for Symptoms of Estrogen Withdrawal in Breast-Cancer Survivors. Journal of Alternative and Complementary Medicine, 2005, 11, 13-20.	2.1	66
188	Organisational factors in relation to control of blood pressure: an observational study. British Journal of General Practice, 2005, 55, 931-7.	1.4	19
189	Treating nausea and vomiting during pregnancy: case outcome. BMJ: British Medical Journal, 2004, 328, 503.1.	2.3	6
190	The DiAMOND trial protocol: a randomised controlled trial of two decision aids for mode of delivery among women with a previous caesarean section [ISRCTN84367722]. BMC Pregnancy and Childbirth, 2004, 4, 25.	2.4	19
191	Decision analysis for newly diagnosed hypertensive patients: a qualitative investigation. Patient Education and Counseling, 2004, 53, 197-203.	2.2	12
192	Coronary risk scores. Lancet, The, 2004, 363, 572-573.	13.7	2
193	Design, analysis and presentation of factorial randomised controlled trials. BMC Medical Research Methodology, 2003, 3, 26.	3.1	237
194	Clinical management of urinary tract infection in women: a prospective cohort study. Family Practice, 2003, 20, 1-6.	1.9	33
195	Quality of care for elderly residents in nursing homes and elderly people living at home: controlled observational study. BMJ: British Medical Journal, 2003, 326, 580-580.	2.3	169
196	The influence of absolute cardiovascular risk, patient utilities, and costs on the decision to treat hypertension. Journal of Hypertension, 2003, 21, 1753-1759.	0.5	70
197	A factorial randomised controlled trial of decision analysis and an information video plus leaflet for newly diagnosed hypertensive patients. British Journal of General Practice, 2003, 53, 446-53.	1.4	74
198	Inter-rater agreement in the scoring of abstracts submitted to a primary care research conference. BMC Health Services Research, 2002, 2, 8.	2.2	127

#	Article	IF	Citations
199	Daily regimen and compliance with treatment. BMJ: British Medical Journal, 2002, 324, 425-425.	2.3	4
200	Rectal bleeding and colorectal cancer. BMJ: British Medical Journal, 2001, 322, 610-610.	2.3	0
201	Shared decision making in hypertension: the impact of patient preferences on treatment choice. Family Practice, 2001, 18, 309-313.	1.9	70
202	The impact of patients' preferences on the treatment of atrial fibrillation: observational study of patient based decision analysis Commentary: patients, preferences, and evidence. BMJ: British Medical Journal, 2000, 320, 1380-1384.	2.3	217
203	Evaluation of computer based clinical decision support system and risk chart for management of hypertension in primary care: randomised controlled trial. BMJ: British Medical Journal, 2000, 320, 686-690.	2.3	225
204	Birth size and arterial compliance in young adults. Lancet, The, 2000, 355, 2136-2137.	13.7	45
205	Staff training to improve participant recruitment into surgical randomised controlled trials: A feasibility study within a trial (SWAT) across four host trials simultaneously. Research Methods in Medicine & Health Sciences, 0, , 263208432211069.	1.2	3