

Heidi K Swanson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7399810/publications.pdf>

Version: 2024-02-01

50
papers

1,223
citations

516710

16
h-index

395702

33
g-index

51
all docs

51
docs citations

51
times ranked

1612
citing authors

#	ARTICLE	IF	CITATIONS
1	Circumpolar patterns of Arctic freshwater fish biodiversity: A baseline for monitoring. <i>Freshwater Biology</i> , 2022, 67, 176-193.	2.4	17
2	Diversity of diatoms, benthic macroinvertebrates, and fish varies in response to different environmental correlates in Arctic rivers across North America. <i>Freshwater Biology</i> , 2022, 67, 95-115.	2.4	15
3	Hair to blood mercury concentration ratios and a retrospective hair segmental mercury analysis in the Northwest Territories, Canada. <i>Environmental Research</i> , 2022, 203, 111800.	7.5	11
4	Habitat area and environmental filters determine avian richness along an elevation gradient in mountain peatlands. <i>Journal of Avian Biology</i> , 2022, 2022, .	1.2	3
5	Understanding among-lake variability of mercury concentrations in Northern Pike (<i>Esox lucius</i>): A whole-ecosystem study in subarctic lakes. <i>Science of the Total Environment</i> , 2022, 822, 153430.	8.0	10
6	Overwintering ecology and movement of anadromous Arctic char (<i>Salvelinus alpinus</i>) in a large, ice-covered river in the Canadian Arctic. <i>Journal of Fish Biology</i> , 2022, 100, 1432-1446.	1.6	7
7	Contributions and perspectives of Indigenous Peoples to the study of mercury in the Arctic. <i>Science of the Total Environment</i> , 2022, 841, 156566.	8.0	10
8	<i>Ecological Diversity</i> , 2021, , 69-117.		15
9	Dietary and non-dietary contributions to among-individual variation in carbon and nitrogen isotopic composition of lake trout. <i>Ecological Indicators</i> , 2021, 123, 107349.	6.3	3
10	David W. Schindler (1940–2021): Trailblazing scientist and advocate for the environment. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, e2106365118.	7.1	0
11	A Bayesian mixing model framework for quantifying temporal variation in source of sediment to lakes across broad hydrological gradients of floodplains. <i>Limnology and Oceanography: Methods</i> , 2021, 19, 540-551.	2.0	2
12	Catchments affect growth rate of Northern Pike, <i>Esox lucius</i> , in subarctic lakes. <i>Aquatic Sciences</i> , 2021, 83, 1.	1.5	4
13	David W. Schindler (1940–2021). <i>Trends in Ecology and Evolution</i> , 2021, 36, 665-667.	8.7	0
14	A meta-collection of nitrogen stable isotope data measured in Arctic marine organisms from the Canadian Beaufort Sea, 1983–2013. <i>BMC Research Notes</i> , 2021, 14, 347.	1.4	2
15	The physical and chemical limnology of Yukon's largest lake, Lhã™Än MÃcñâ™ (Kluane Lake), prior to the 2016 â€™Ä™ÿ ChÃ™ diversion. <i>Arctic Science</i> , 2021, 7, 655-678.	2.3	4
16	<i>Trophic Ecology</i> , 2021, , 287-314.		7
17	Mercury accumulation in sediments of Lhã™Än MÃcñâ™ (Kluane Lake, YT): Response to past hydrological change. <i>Arctic, Antarctic, and Alpine Research</i> , 2021, 53, 179-195.	1.1	1
18	Fish growth rates and lake sulphate explain variation in mercury levels in ninespine stickleback (<i>Pungitius pungitius</i>) on the Arctic Coastal Plain of Alaska. <i>Science of the Total Environment</i> , 2020, 743, 140564.	8.0	13

#	ARTICLE	IF	CITATIONS
19	Dietary versus nondietary fatty acid profiles of lake trout ecotypes from Lake Superior and Great Bear Lake: Are fish really what they eat?. Canadian Journal of Fisheries and Aquatic Sciences, 2020, 77, 1209-1220.	1.4	5
20	Gut contents from multiple morphs of lake trout (<i>Salvelinus namaycush</i>) at two offshore shoals in Lake Superior. Journal of Great Lakes Research, 2020, 46, 1382-1390.	1.9	7
21	Anadromy and marine habitat use of Lake trout (<i>Salvelinus namaycush</i>) from the central Canadian Arctic. Journal of Fish Biology, 2020, 96, 1489-1494.	1.6	3
22	Spatiotemporal patterns in trophic niche overlap among five salmonines in Lake Michigan, USA. Canadian Journal of Fisheries and Aquatic Sciences, 2020, 77, 1059-1075.	1.4	10
23	Towards reconciliation: 10 Calls to Action to natural scientists working in Canada. Facets, 2020, 5, 769-783.	2.4	85
24	Screening-level risk assessment of methylmercury for non-anadromous Arctic char (<i>Salvelinus</i>)	4.8	11
25	Seasonal variation in resource overlap of invasive and native fishes revealed by stable isotopes. Biological Invasions, 2019, 21, 315-321.	2.4	13
26	Feeding of Greenland halibut (<i>Reinhardtius hippoglossoides</i>) in the Canadian Beaufort Sea. Journal of Marine Systems, 2018, 183, 32-41.	2.1	17
27	Relationships between depth and $\delta^{15}N$ of Arctic benthos vary among regions and trophic functional groups. Deep-Sea Research Part I: Oceanographic Research Papers, 2018, 135, 56-64.	1.4	9
28	Design of a human biomonitoring community-based project in the Northwest Territories Mackenzie Valley, Canada, to investigate the links between nutrition, contaminants and country foods. International Journal of Circumpolar Health, 2018, 77, 1510714.	1.2	13
29	Mercury and omega-3 fatty acid profiles in freshwater fish of the Dehcho Region, Northwest Territories: Informing risk benefit assessments. Science of the Total Environment, 2018, 637-638, 1508-1517.	8.0	25
30	Optimal sampling methods for modelling the occupancy of Arctic grayling (<i>Thymallus arcticus</i>) in the Canadian Barrenlands. Canadian Journal of Fisheries and Aquatic Sciences, 2017, 74, 1564-1574.	1.4	9
31	Associations between omega-3 fatty acids, selenium content, and mercury levels in wild-harvested fish from the Dehcho Region, Northwest Territories, Canada. Journal of Toxicology and Environmental Health - Part A: Current Issues, 2017, 80, 18-31.	2.3	22
32	Trophic variability of Arctic fishes in the Canadian Beaufort Sea: a fatty acids and stable isotopes approach. Polar Biology, 2016, 39, 1267-1282.	1.2	24
33	Low Annual Fidelity and Early Upstream Migration of Anadromous Arctic Char in a Variable Environment. Transactions of the American Fisheries Society, 2016, 145, 931-942.	1.4	26
34	A new probabilistic method for quantifying n -dimensional ecological niches and niche overlap. Ecology, 2015, 96, 318-324.	3.2	306
35	Mercury in freshwater ecosystems of the Canadian Arctic: Recent advances on its cycling and fate. Science of the Total Environment, 2015, 509-510, 41-66.	8.0	64
36	Long-distance anadromous migration in a fresh water specialist: the Lake Trout (— <i>Salvelinus</i>)	0.1	10

#	ARTICLE	IF	CITATIONS
37	Species and Life History Affect the Utility of Otolith Chemical Composition for Determining Natal Stream of Origin for Pacific Salmon. Transactions of the American Fisheries Society, 2013, 142, 1370-1380.	1.4	12
38	SCIENTISTS, ON SAVING SCIENCE. Limnology and Oceanography Bulletin, 2013, 22, 76-78.	0.4	2
39	Biomagnification of mercury through lake trout (<i>Salvelinus namaycush</i>) food webs of lakes with different physical, chemical and biological characteristics. Science of the Total Environment, 2012, 438, 135-143.	8.0	96
40	Quantifying importance of marine prey in the diets of two partially anadromous fishes. Canadian Journal of Fisheries and Aquatic Sciences, 2011, 68, 2020-2028.	1.4	27
41	Differences in Mercury Bioaccumulation between Polar Bears (<i>Ursus maritimus</i>) from the Canadian high- and sub-Arctic. Environmental Science & Technology, 2011, 45, 5922-5928.	10.0	49
42	Comparison of mercury concentrations in landlocked, resident, and sea-run fish (<i>Salvelinus</i>) in the Arctic. Environmental Science & Technology, 2010, 44, 3286-3292.	4.3	35
43	Effects of Partially Anadromous Arctic Charr (<i>Salvelinus alpinus</i>) Populations on Ecology of Coastal Arctic Lakes. Ecosystems, 2010, 13, 261-274.	3.4	25
44	Mercury Concentrations in Arctic Food Fishes Reflect the Presence of Anadromous Arctic Charr (<i>Salvelinus alpinus</i>), Species, and Life History. Environmental Science & Technology, 2010, 44, 3286-3292.	10.0	61
45	Anadromy in Arctic populations of lake trout (<i>Salvelinus namaycush</i>): otolith microchemistry, stable isotopes, and comparisons with Arctic char (<i>Salvelinus alpinus</i>). Canadian Journal of Fisheries and Aquatic Sciences, 2010, 67, 842-853.	1.4	61
46	The Effect of Anadromous Arctic Charr (<i>Salvelinus alpinus</i>) on Food Web Structure and Contaminant Concentrations in Coastal Arctic Lakes. Arctic, 2009, 60, .	0.4	1
47	Mercury Bioaccumulation in Forage Fish Communities Invaded by Rainbow Smelt (<i>Osmerus mordax</i>). Environmental Science & Technology, 2006, 40, 1439-1446.	10.0	35
48	TEMPORAL CHANGES IN MERCURY BIOACCUMULATION BY PREDATORY FISHES OF BOREAL LAKES FOLLOWING THE INVASION OF AN EXOTIC FORAGE FISH. Environmental Toxicology and Chemistry, 2003, 22, 2057.	4.3	29
49	The Canadian Beaufort Shelf trophic structure: evaluating an ecosystem modelling approach by comparison with observed stable isotopic structure. Arctic Science, 0, , .	2.3	5
50	Occupancy of young-of-year Arctic grayling (<i>Thymallus arcticus</i>) in Barrenland streams. Hydrobiologia, 0, , 1.	2.0	1