

Ashley R Pedler

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7407901/publications.pdf>

Version: 2024-02-01

25
papers

884
citations

516710

16
h-index

610901

24
g-index

25
all docs

25
docs citations

25
times ranked

855
citing authors

#	ARTICLE	IF	CITATIONS
1	Magnetic Resonance Spectroscopy Assessment of Brain Metabolite Concentrations in Individuals With Chronic Whiplash-associated Disorder. <i>Clinical Journal of Pain</i> , 2021, 37, 28-37.	1.9	2
2	Comparative Strength and Endurance Parameters of the Craniocervical and Cervicothoracic Extensors and Flexors in Females With and Without Idiopathic Neck Pain. <i>Journal of Applied Biomechanics</i> , 2019, 35, 209-215.	0.8	9
3	Investigating the Fear Avoidance Model in People With Whiplash. <i>Clinical Journal of Pain</i> , 2018, 34, 130-137.	1.9	11
4	Intramuscular fat is present in cervical multifidus but not soleus in patients with chronic whiplash associated disorders. <i>PLoS ONE</i> , 2018, 13, e0197438.	2.5	14
5	Exercise induced hypoalgesia is elicited by isometric, but not aerobic exercise in individuals with chronic whiplash associated disorders. <i>Scandinavian Journal of Pain</i> , 2017, 15, 14-21.	1.3	52
6	A Web Based Version of the Cervical Joint Position Error Test: Reliability of Measurements from Face Tracking Software. <i>Lecture Notes of the Institute for Computer Sciences, Social-Informatics and Telecommunications Engineering</i> , 2017, , 297-301.	0.3	0
7	Addition of posttraumatic stress and sensory hypersensitivity more accurately estimates disability and pain than fear avoidance measures alone after whiplash injury. <i>Pain</i> , 2016, 157, 1645-1654.	4.2	31
8	Evaluating the neck joint position sense error with a standard computer and a webcam. <i>Manual Therapy</i> , 2016, 26, 231-234.	1.6	11
9	Physiotherapists' Beliefs About Whiplash-associated Disorder: A Comparison Between Singapore and Queensland, Australia. <i>Physiotherapy Research International</i> , 2015, 20, 77-86.	1.5	9
10	Morphological changes in the cervical muscles of women with chronic whiplash can be modified with exercise—a pilot study. <i>Muscle and Nerve</i> , 2015, 52, 772-779.	2.2	37
11	The Geography of Fatty Infiltrates Within the Cervical Multifidus and Semispinalis Cervicis in Individuals With Chronic Whiplash-Associated Disorders. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2015, 45, 281-288.	3.5	43
12	Differential Changes in Muscle Composition Exist in Traumatic and Nontraumatic Neck Pain. <i>Spine</i> , 2014, 39, 39-47.	2.0	87
13	Less Efficacious Conditioned Pain Modulation and Sensory Hypersensitivity in Chronic Whiplash-associated Disorders in Singapore. <i>Clinical Journal of Pain</i> , 2014, 30, 436-442.	1.9	30
14	Patients with chronic whiplash can be subgrouped on the basis of symptoms of sensory hypersensitivity and posttraumatic stress. <i>Pain</i> , 2013, 154, 1640-1648.	4.2	37
15	Laterality judgments are not impaired in patients with chronic whiplash associated disorders. <i>Manual Therapy</i> , 2013, 18, 72-76.	1.6	19
16	Magnetic Resonance Imaging Changes in the Size and Shape of the Oropharynx Following Acute Whiplash Injury. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2012, 42, 912-918.	3.5	13
17	Evidence of Spinal Cord Hyperexcitability as Measured With Nociceptive Flexion Reflex (NFR) Threshold in Chronic Lateral Epicondylalgia With or Without a Positive Neurodynamic Test. <i>Journal of Pain</i> , 2012, 13, 676-684.	1.4	37
18	Assessing Fear-Avoidance Beliefs in Patients With Whiplash-associated Disorders. <i>Clinical Journal of Pain</i> , 2011, 27, 502-507.	1.9	26

#	ARTICLE	IF	CITATIONS
19	Relationship Between Pressure Pain Thresholds and Pain Ratings in Patients With Whiplash-associated Disorders. <i>Clinical Journal of Pain</i> , 2011, 27, 495-501.	1.9	57
20	The Temporal Development of Fatty Infiltrates in the Neck Muscles Following Whiplash Injury: An Association with Pain and Posttraumatic Stress. <i>PLoS ONE</i> , 2011, 6, e21194.	2.5	91
21	Magnetic Resonance Imaging Findings of Fatty Infiltrate in the Cervical Flexors in Chronic Whiplash. <i>Spine</i> , 2010, 35, 948-954.	2.0	105
22	Cervical lateral glide increases nociceptive flexion reflex threshold but not pressure or thermal pain thresholds in chronic whiplash associated disorders: A pilot randomised controlled trial. <i>Manual Therapy</i> , 2010, 15, 149-153.	1.6	76
23	Diffusion-Weighted MRI for the Healthy Cervical Multifidus: A Potential Method for Studying Neck Muscle Physiology Following Spinal Trauma. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2010, 40, 722-728.	3.5	16
24	The Pain Catastrophising Scale. <i>Journal of Physiotherapy</i> , 2010, 56, 137.	1.7	14
25	A neuropathic pain component is common in acute whiplash and associated with a more complex clinical presentation. <i>Manual Therapy</i> , 2009, 14, 173-179.	1.6	57