
Penny Rapaport

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7410313/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	START (STrAtegies for RelaTives) study: a pragmatic randomised controlled trial to determine the clinical effectiveness and cost-effectiveness of a manual-based coping strategy programme in promoting the mental health of carers of people with dementia. Health Technology Assessment, 2014, 18, 1-242.	1.3	156
2	Clinical effectiveness of a manual based coping strategy programme (START, STrAtegies for RelaTives) in promoting the mental health of carers of family members with dementia: pragmatic randomised controlled trial. BMJ, The, 2013, 347, f6276-f6276.	3.0	154
3	Prevalence of and associations with agitation in residents with dementia living in care homes: MARQUE cross-sectional study. BJPsych Open, 2017, 3, 171-178.	0.3	98

Do care homes deliver person-centred care? A cross-sectional survey of staff-reported abusive and positive behaviours towards residents from the MARQUE (Managing Agitation and Raising Quality of) Tj ETQq0 0 O1r.gBT /Overstock 10 Tf 4

5	Long-term clinical and cost-effectiveness of psychological intervention for family carers of people with dementia: a single-blind, randomised, controlled trial. Lancet Psychiatry,the, 2014, 1, 539-548.	3.7	68
6	Cost effectiveness of a manual based coping strategy programme in promoting the mental health of family carers of people with dementia (the START (STrAtegies for RelaTives) study): a pragmatic randomised controlled trial. BMJ, The, 2013, 347, f6342-f6342.	3.0	67
7	Systematic review of the effective components of psychosocial interventions delivered by care home staff to people with dementia. BMJ Open, 2017, 7, e014177.	0.8	52
8	Developing the New Interventions for independence in Dementia Study (NIDUS) theoretical model for supporting people to live well with dementia at home for longer: a systematic review of theoretical models and Randomised Controlled Trial evidence. Social Psychiatry and Psychiatric Epidemiology, 2020, 55, 1-14.	1.6	48
9	Clinical and cost-effectiveness of the Managing Agitation and Raising Quality of Life (MARQUE) intervention for agitation in people with dementia in care homes: a single-blind, cluster-randomised controlled trial. Lancet Psychiatry,the, 2019, 6, 293-304.	3.7	41
10	Augmenting the theory of planned behaviour: Motivation to provide practical assistance and emotional support to parents. Psychology and Health, 2000, 15, 309-324.	1.2	38
11	"l just keep thinking that I don't want to rely on people.―a qualitative study of how people living with dementia achieve and maintain independence at home: stakeholder perspectives. BMC Geriatrics, 2020, 20, 5.	1.1	36
12	Effectiveness of START psychological intervention in reducing abuse by dementia family carers: randomized controlled trial. International Psychogeriatrics, 2016, 28, 881-887.	0.6	33
13	Care workers, the unacknowledged persons in person-centred care: A secondary qualitative analysis of UK care home staff interviews. PLoS ONE, 2018, 13, e0200031.	1.1	33
14	DREAMS-START (Dementia RElAted Manual for Sleep; STrAtegies for RelaTives) for people with dementia and sleep disturbances: a single-blind feasibility and acceptability randomized controlled trial. International Psychogeriatrics, 2019, 31, 251-265.	0.6	32
15	How do care home staff understand, manage and respond to agitation in people with dementia? A qualitative study. BMJ Open, 2018, 8, e022260.	0.8	30
16	Coping strategies as mediators of the effect of the START (strategies for RelaTives) intervention on psychological morbidity for family carers of people with dementia in a randomised controlled trial. Journal of Affective Disorders, 2014, 168, 298-305.	2.0	29
17	Clinical effectiveness of the START (STrAtegies for RelaTives) psychological intervention for family carers and the effects on the cost of care for people with dementia: 6-year follow-up of a randomised controlled trial. British Journal of Psychiatry, 2020, 216, 35-42.	1.7	27
18	"In the Bengali Vocabulary, There Is No Such Word as Care Home―Caring Experiences of UK Bangladeshi and Indian Family Carers of People Living With Dementia at Home. Gerontologist, The, 2020, 60, 331-339.	2.3	22

Penny Rapaport

#	Article	IF	CITATIONS
19	A systematic review evaluating the impact of paid home carer training, supervision, and other interventions on the health and well-being of older home care clients. International Psychogeriatrics, 2017, 29, 595-604.	0.6	21
20	START (STrAtegies for RelaTives) coping strategy for family carers of adults with dementia: qualitative study of participants' views about the intervention. BMJ Open, 2014, 4, e005273-e005273.	0.8	19
21	Social connectedness and dementia prevention: Pilot of the APPLE-Tree video-call intervention during the Covid-19 pandemic. Dementia, 2021, 20, 2779-2801.	1.0	17
22	Effects of nonpharmacological interventions on functioning of people living with dementia at home: A systematic review of randomised controlled trials. International Journal of Geriatric Psychiatry, 2019, 34, 1386-1402.	1.3	14
23	†You can't just put somebody in a situation with no armour'. An ethnographic exploration of the training and support needs of homecare workers caring for people living with dementia. Dementia, 2021, 20, 2982-3005.	1.0	14
24	An intervention to improve sleep for people living with dementia: Reflections on the development and co-production of DREAMS:START (Dementia RElAted Manual for Sleep: STrAtegies for RelaTives). Dementia, 2018, 17, 976-989.	1.0	13
25	A systematic review of observational studies of adult home care. Health and Social Care in the Community, 2019, 27, 1388-1400.	0.7	13
26	APPLEâ€Tree (Active Prevention in People at risk of dementia: Lifestyle, bEhaviour change and Technology) Tj ETC Psychiatry, 2020, 35, 811-819.	Qq0 0 0 rg 1.3	gBT /Overlock 13
27	" <i>Falling through the cracks</i> â€; Stakeholders' views around the concept and diagnosis of mild cognitive impairment and their understanding of dementia prevention. International Journal of Geriatric Psychiatry, 2020, 35, 1349-1357.	1.3	12
28	A mixed-methods feasibility study of a goal-focused manualised intervention to support people with dementia to stay living independently at home with support from family carers: NIDUS (New) Tj ETQq0 0 0 rgBT /	Overlock	101 1 f 50 377
29	Clinical and cost-effectiveness of a New psychosocial intervention to support Independence in Dementia (NIDUS-family) for family carers and people living with dementia in their own homes: a randomised controlled trial. Trials, 2021, 22, 865.	0.7	11
30	Carer coping and resident agitation as predictors of quality of life in care home residents living with dementia: Managing Agitation and Raising Quality of Life (MARQUE) English national care home prospective cohort study. International Journal of Geriatric Psychiatry, 2019, 34, 106-113.	1.3	10
31	Supporting independence at home for people living with dementia: a qualitative ethnographic study of homecare. Social Psychiatry and Psychiatric Epidemiology, 2021, 56, 2323-2336.	1.6	10
32	Co-designing complex interventions with people living with dementia and their supporters. Dementia, 2022, 21, 426-441.	1.0	10
33	Time to reflect is a rare and valued opportunity; a pilot of the NIDUSâ€professional dementia training intervention for homecare workers during the Covidâ€19 pandemic. Health and Social Care in the Community, 2022, 30, .	0.7	8
34	A manual-based intervention for carers of people with dementia and sleep disturbances: an acceptability and feasibility RCT. Health Technology Assessment, 2018, 22, 1-408.	1.3	7
35	Prevalence of Dementia in Older Adults in Central and Eastern Europe: A Systematic Review and Meta-Analysis. Psychiatry International, 2021, 2, 191-210.	0.5	6
36	Estimation of the minimal clinically important difference on the Aberrant Behaviour Checklist–Irritability (ABC-I) for people with intellectual disabilities who display aggressive challenging behaviour: A triangulated approach. Research in Developmental Disabilities, 2022, 124, 104202.	1.2	6

Penny Rapaport

#	Article	IF	CITATIONS
37	Why do staff and family think differently about quality of life in dementia? A qualitative study exploring perspectives in care homes. International Journal of Geriatric Psychiatry, 2019, 34, 1784-1791.	1.3	5
38	Exploring how people with dementia can be best supported to manage long-term conditions: a qualitative study of stakeholder perspectives. BMJ Open, 2020, 10, e041873.	0.8	4
39	Longâ€ŧerm implementation of the Managing Agitation and Raising QUality of lifE intervention in care homes: AÂqualitative study. International Journal of Geriatric Psychiatry, 2021, 36, 1252-1263.	1.3	4
40	Implementation of START (STrAtegies for RelaTives) for dementia carers in the third sector: Widening access to evidence-based interventions. PLoS ONE, 2021, 16, e0250410.	1.1	4
41	Disseminating START: training clinical psychologists and admiral nurses as trainers in a psychosocial intervention for carers of people with dementia's depressive and anxiety symptoms. BMJ Open, 2017, 7, e017759.	0.8	3
42	Implementation should now be considered in parallel with designing clinical research. International Psychogeriatrics, 2020, 32, 1393-1395.	0.6	3
43	Experience of UK Latin Americans caring for a relative living with dementia: A qualitative study of family carers. Dementia, 2022, 21, 1574-1595.	1.0	2
44	Promoting Functional Independence in Dementia. , 2021, , 315-339.		1
45	Acceptability and feasibility of a Japanese version of STrAtegies for RelaTives (START-J): A manualized coping strategy program for family caregivers of relatives living with dementia. Dementia, 2021, 20, 985-1004.	1.0	1
46	[P4–317]: HOW DO CARE HOME STAFF EXPERIENCE, UNDERSTAND AND MANAGE AGITATION IN RESIDENTS WITH DEMENTIA? A QUALITATIVE STUDY. Alzheimer's and Dementia, 2017, 13, P1411.	0.4	0
47	[P1–298]: RESEARCHING DEMENTIAâ€RELATED SLEEP PROBLEMS USING ACTIâ€WATCHES: PRELIMINARY INSIG FROM THE DREAMS START STUDY. Alzheimer's and Dementia, 2017, 13, P367.	CHTS 0.4	0
48	411 - Implementing the STrAtegies for RelaTives (START) intervention in Portugal – preliminary findings. International Psychogeriatrics, 2020, 32, 127-127.	0.6	0