

# Tomonori Okamura

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/7414334/tomonori-okamura-publications-by-citations.pdf>

**Version:** 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

439  
papers

14,022  
citations

54  
h-index

100  
g-index

505  
ext. papers

16,217  
ext. citations

4.4  
avg, IF

5.75  
L-index

#	Paper	IF	Citations
439	Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. <i>Nature</i> , <b>2011</b> , 478, 103-9	50.4	1564
438	Cardiovascular disease and risk factors in Asia: a selected review. <i>Circulation</i> , <b>2008</b> , 118, 2702-9	16.7	500
437	A simple method to estimate populational 24-h urinary sodium and potassium excretion using a casual urine specimen. <i>Journal of Human Hypertension</i> , <b>2002</b> , 16, 97-103	2.6	419
436	Executive summary of the Japan Atherosclerosis Society (JAS) guidelines for the diagnosis and prevention of atherosclerotic cardiovascular diseases in Japan -2012 version. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2013</b> , 20, 517-23	4	313
435	Vegetarian diets and blood pressure: a meta-analysis. <i>JAMA Internal Medicine</i> , <b>2014</b> , 174, 577-87	11.5	310
434	Japan Atherosclerosis Society (JAS) Guidelines for Prevention of Atherosclerotic Cardiovascular Diseases 2017. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2018</b> , 25, 846-984	4	295
433	Serum triglycerides and risk of coronary heart disease among Japanese men and women. <i>American Journal of Epidemiology</i> , <b>2001</b> , 153, 490-9	3.8	246
432	Marine-derived n-3 fatty acids and atherosclerosis in Japanese, Japanese-American, and white men: a cross-sectional study. <i>Journal of the American College of Cardiology</i> , <b>2008</b> , 52, 417-24	15.1	181
431	Linoleic acid, other fatty acids, and the risk of stroke. <i>Stroke</i> , <b>2002</b> , 33, 2086-93	6.7	169
430	Cigarette smoking as a risk factor for stroke death in Japan: NIPPON DATA80. <i>Stroke</i> , <b>2004</b> , 35, 1836-41	6.7	150
429	Japanese men have larger areas of visceral adipose tissue than Caucasian men in the same levels of waist circumference in a population-based study. <i>International Journal of Obesity</i> , <b>2006</b> , 30, 1163-5	5.5	148
428	A low prevalence of coronary heart disease among subjects with increased high-density lipoprotein cholesterol levels, including those with plasma cholesteryl ester transfer protein deficiency. <i>Preventive Medicine</i> , <b>1998</b> , 27, 659-67	4.3	148
427	The relationship between serum total cholesterol and all-cause or cause-specific mortality in a 17.3-year study of a Japanese cohort. <i>Atherosclerosis</i> , <b>2007</b> , 190, 216-23	3.1	136
426	Impact of high-normal blood pressure on the risk of cardiovascular disease in a Japanese urban cohort: the Suita study. <i>Hypertension</i> , <b>2008</b> , 52, 652-9	8.5	127
425	Guidelines for the management of familial hypercholesterolemia. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2012</b> , 19, 1043-60	4	126
424	Less subclinical atherosclerosis in Japanese men in Japan than in White men in the United States in the post-World War II birth cohort. <i>American Journal of Epidemiology</i> , <b>2007</b> , 165, 617-24	3.8	115
423	What cause of mortality can we predict by cholesterol screening in the Japanese general population?. <i>Journal of Internal Medicine</i> , <b>2003</b> , 253, 169-80	10.8	115

422	Serum total homocysteine concentrations and risk of stroke and its subtypes in Japanese. <i>Circulation</i> , <b>2004</b> , 109, 2766-72	16.7	111
421	Predicting coronary heart disease using risk factor categories for a Japanese urban population, and comparison with the framingham risk score: the suita study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 784-98	4	110
420	Relation of blood pressure and all-cause mortality in 180,000 Japanese participants: pooled analysis of 13 cohort studies. <i>Hypertension</i> , <b>2008</b> , 51, 1483-91	8.5	109
419	Relationship between metabolic risk factor clustering and cardiovascular mortality stratified by high blood glucose and obesity: NIPPON DATA90, 1990-2000. <i>Diabetes Care</i> , <b>2007</b> , 30, 1533-8	14.6	106
418	A nutrient-wide association study on blood pressure. <i>Circulation</i> , <b>2012</b> , 126, 2456-64	16.7	104
417	The inverse relationship between serum high-density lipoprotein cholesterol level and all-cause mortality in a 9.6-year follow-up study in the Japanese general population. <i>Atherosclerosis</i> , <b>2006</b> , 184, 143-50	3.1	102
416	Resting heart rate and cause-specific death in a 16.5-year cohort study of the Japanese general population. <i>American Heart Journal</i> , <b>2004</b> , 147, 1024-32	4.9	100
415	Guidelines for Diagnosis and Treatment of Familial Hypercholesterolemia 2017. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2018</b> , 25, 751-770	4	97
414	Relation between sleep quality and quantity, quality of life, and risk of developing diabetes in healthy workers in Japan: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) Study. <i>BMC Public Health</i> , <b>2007</b> , 7, 129	4.1	96
413	Low-density lipoprotein cholesterol and non-high-density lipoprotein cholesterol and the incidence of cardiovascular disease in an urban Japanese cohort study: The Suita study. <i>Atherosclerosis</i> , <b>2009</b> , 203, 587-92	3.1	94
412	Effects of habitual alcohol intake on ambulatory blood pressure, heart rate, and its variability among Japanese men. <i>Hypertension</i> , <b>2009</b> , 53, 13-9	8.5	90
411	Relationship between blood pressure category and incidence of stroke and myocardial infarction in an urban Japanese population with and without chronic kidney disease: the Suita Study. <i>Stroke</i> , <b>2009</b> , 40, 2674-9	6.7	89
410	Blood pressure categories and long-term risk of cardiovascular disease according to age group in Japanese men and women. <i>Hypertension Research</i> , <b>2012</b> , 35, 947-53	4.7	87
409	Diagnostic criteria for dyslipidemia. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2013</b> , 20, 655-60	4	83
408	Common variants in the ATP2B1 gene are associated with susceptibility to hypertension: the Japanese Millennium Genome Project. <i>Hypertension</i> , <b>2010</b> , 56, 973-80	8.5	83
407	LOX index, a novel predictive biochemical marker for coronary heart disease and stroke. <i>Clinical Chemistry</i> , <b>2010</b> , 56, 550-8	5.5	82
406	Alcohol intake and premature coronary heart disease in urban Japanese men. <i>American Journal of Epidemiology</i> , <b>1998</b> , 147, 59-65	3.8	80
405	J Epidemiol 2010;20(Suppl 3):S5068514 Integration of Data from NIPPON DATA80/90 and National Nutrition Survey in Japan: For Cohort Studies of Representative Japanese on Nutrition. <i>Journal of Epidemiology</i> , <b>2010</b> , 20, 346-346	3.4	78

404	Chronic kidney disease is a risk factor for cardiovascular death in a community-based population in Japan: NIPPON DATA90. <i>Circulation Journal</i> , <b>2006</b> , 70, 954-9	2.9	78
403	Association between fish consumption and all-cause and cause-specific mortality in Japan: NIPPON DATA80, 1980-99. <i>American Journal of Medicine</i> , <b>2005</b> , 118, 239-45	2.4	76
402	Impact of metabolic syndrome components on the incidence of cardiovascular disease in a general urban Japanese population: the Suita study. <i>Hypertension Research</i> , <b>2008</b> , 31, 2027-35	4.7	74
401	Small dense low-density lipoproteins cholesterol can predict incident cardiovascular disease in an urban Japanese cohort: the Suita study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2013</b> , 20, 195-203	4	72
400	Age-specific effects of systolic and diastolic blood pressures on mortality due to cardiovascular diseases among Japanese men (NIPPON DATA80). <i>Journal of Hypertension</i> , <b>2006</b> , 24, 459-62	1.9	69
399	Genetic link between obesity and MMP14-dependent adipogenic collagen turnover. <i>Diabetes</i> , <b>2010</b> , 59, 2484-94	0.9	68
398	Epidemiologic study of the association of low-Km mitochondrial acetaldehyde dehydrogenase genotypes with blood pressure level and the prevalence of hypertension in a general population. <i>Hypertension Research</i> , <b>2002</b> , 25, 857-64	4.7	68
397	Lower levels of serum albumin and total cholesterol associated with decline in activities of daily living and excess mortality in a 12-year cohort study of elderly Japanese. <i>Journal of the American Geriatrics Society</i> , <b>2008</b> , 56, 529-35	5.6	67
396	Combined cardiovascular risk factors and outcome: NIPPON DATA80, 1980-1994. <i>Circulation Journal</i> , <b>2006</b> , 70, 960-4	2.9	67
395	Serum Uric Acid and Mortality Form Cardiovascular Disease: EPOCH-JAPAN Study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2016</b> , 23, 692-703	4	66
394	Egg consumption, serum cholesterol, and cause-specific and all-cause mortality: the National Integrated Project for Prospective Observation of Non-communicable Disease and Its Trends in the Aged, 1980 (NIPPON DATA80). <i>American Journal of Clinical Nutrition</i> , <b>2004</b> , 80, 58-63	7	66
393	Metabolomic profiling reveals novel biomarkers of alcohol intake and alcohol-induced liver injury in community-dwelling men. <i>Environmental Health and Preventive Medicine</i> , <b>2016</b> , 21, 18-26	4.2	64
392	Relation between serum total cholesterol level and cardiovascular disease stratified by sex and age group: a pooled analysis of 65 594 individuals from 10 cohort studies in Japan. <i>Journal of the American Heart Association</i> , <b>2012</b> , 1, e001974	6	64
391	Validation of the association of genetic variants on chromosome 9p21 and 1q41 with myocardial infarction in a Japanese population. <i>Circulation Journal</i> , <b>2008</b> , 72, 1213-7	2.9	63
390	Association of extremely high levels of high-density lipoprotein cholesterol with cardiovascular mortality in a pooled analysis of 9 cohort studies including 43,407 individuals: The EPOCH-JAPAN study. <i>Journal of Clinical Lipidology</i> , <b>2018</b> , 12, 674-684.e5	4.9	61
389	Impact of metabolic syndrome on the risk of cardiovascular disease mortality in the United States and in Japan. <i>American Journal of Cardiology</i> , <b>2014</b> , 113, 84-9	3	60
388	Associations of socioeconomic status with prevalence, awareness, treatment, and control of hypertension in a general Japanese population: NIPPON DATA2010. <i>Journal of Hypertension</i> , <b>2017</b> , 35, 401-408	1.9	59
387	Long-term risk of BP values above normal for cardiovascular mortality: a 24-year observation of Japanese aged 30 to 92 years. <i>Journal of Hypertension</i> , <b>2012</b> , 30, 2299-306	1.9	59

386	CDH13 gene coding T-cadherin influences variations in plasma adiponectin levels in the Japanese population. <i>Human Mutation</i> , <b>2012</b> , 33, 402-10	4.7	55
385	Dietary sodium-to-potassium ratio as a risk factor for stroke, cardiovascular disease and all-cause mortality in Japan: the NIPPON DATA80 cohort study. <i>BMJ Open</i> , <b>2016</b> , 6, e011632	3	53
384	Serum 1,5-anhydro-D-glucitol levels predict first-ever cardiovascular disease: an 11-year population-based cohort study in Japan, the Suita study. <i>Atherosclerosis</i> , <b>2011</b> , 216, 477-83	3.1	53
383	Effect of a 4-year workplace-based physical activity intervention program on the blood lipid profiles of participating employees: the high-risk and population strategy for occupational health promotion (HIPOP-OHP) study. <i>Atherosclerosis</i> , <b>2008</b> , 197, 784-90	3.1	53
382	A prospective study of passive smoking and risk of diabetes in a cohort of workers: the High-Risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) study. <i>Diabetes Care</i> , <b>2008</b> , 31, 732-4	14.6	53
381	Mortality risk attributable to atrial fibrillation in middle-aged and elderly people in the Japanese general population: nineteen-year follow-up in NIPPON DATA80. <i>Circulation Journal</i> , <b>2007</b> , 71, 814-9	2.9	52
380	Influence of smoking combined with another risk factor on the risk of mortality from coronary heart disease and stroke: pooled analysis of 10 Japanese cohort studies. <i>Cerebrovascular Diseases</i> , <b>2012</b> , 33, 480-91	3.2	50
379	Higher liver fat content among Japanese in Japan compared with non-Hispanic whites in the United States. <i>Metabolism: Clinical and Experimental</i> , <b>2009</b> , 58, 1200-7	12.7	50
378	Hypertension and lifetime risk of stroke. <i>Journal of Hypertension</i> , <b>2016</b> , 34, 116-22	1.9	48
377	Fruit and vegetable intake and mortality from cardiovascular disease in Japan: a 24-year follow-up of the NIPPON DATA80 Study. <i>European Journal of Clinical Nutrition</i> , <b>2015</b> , 69, 482-8	5.2	47
376	A genome-wide association study of hypertension-related phenotypes in a Japanese population. <i>Circulation Journal</i> , <b>2010</b> , 74, 2353-9	2.9	47
375	Relationship of elevated casual blood glucose level with coronary heart disease, cardiovascular disease and all-cause mortality in a representative sample of the Japanese population. NIPPON DATA80. <i>Diabetologia</i> , <b>2008</b> , 51, 575-82	10.3	47
374	White blood cell count and risk of all-cause and cardiovascular mortality in nationwide sample of Japanese--results from the NIPPON DATA90. <i>Circulation Journal</i> , <b>2007</b> , 71, 479-85	2.9	47
373	Effects of public education by television on knowledge of early stroke symptoms among a Japanese population aged 40 to 74 years: a controlled study. <i>Stroke</i> , <b>2012</b> , 43, 545-9	6.7	46
372	Specific alcoholic beverage and blood pressure in a middle-aged Japanese population: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) Study. <i>Journal of Human Hypertension</i> , <b>2004</b> , 18, 9-16	2.6	46
371	Prediction of cardiovascular disease mortality by proteinuria and reduced kidney function: pooled analysis of 39,000 individuals from 7 cohort studies in Japan. <i>American Journal of Epidemiology</i> , <b>2013</b> , 178, 1-11	3.8	45
370	Population attributable numbers and fractions of deaths due to smoking: a pooled analysis of 180,000 Japanese. <i>Preventive Medicine</i> , <b>2011</b> , 52, 60-5	4.3	45
369	Genetic variation in aldehyde dehydrogenase 2 and the effect of alcohol consumption on cholesterol levels. <i>Atherosclerosis</i> , <b>2002</b> , 164, 171-7	3.1	45

368	Changes in Weight, cardiovascular risk factors and estimated risk of coronary heart disease following smoking cessation in Japanese male workers: HIPOP-OHP study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2010</b> , 17, 12-20	4	44
367	gamma-Glutamyltransferase predicts cardiovascular death among Japanese women. <i>Atherosclerosis</i> , <b>2007</b> , 194, 498-504	3.1	44
366	Identification of genetic markers associated with high-density lipoprotein-cholesterol by genome-wide screening in a Japanese population: the Suita study. <i>Circulation Journal</i> , <b>2009</b> , 73, 1119-26 <sup>2.9</sup>	2.9	43
365	Effects of serum B vitamins on elevated plasma homocysteine levels associated with the mutation of methylenetetrahydrofolate reductase gene in Japanese. <i>Atherosclerosis</i> , <b>2002</b> , 164, 321-8	3.1	43
364	Long-chain n-3 polyunsaturated fatty acids intake and cardiovascular disease mortality risk in Japanese: a 24-year follow-up of NIPPON DATA80. <i>Atherosclerosis</i> , <b>2014</b> , 232, 384-9	3.1	42
363	A multicenter study on the precision and accuracy of homogeneous assays for LDL-cholesterol: comparison with a beta-quantification method using fresh serum obtained from non-diseased and diseased subjects. <i>Atherosclerosis</i> , <b>2012</b> , 225, 208-15	3.1	42
362	HbA1c and the risks for all-cause and cardiovascular mortality in the general Japanese population: NIPPON DATA90. <i>Diabetes Care</i> , <b>2013</b> , 36, 3759-65	14.6	42
361	The high-risk and population strategy for occupational health promotion (HIPOP-OHP) study: study design and cardiovascular risk factors at the baseline survey. <i>Journal of Human Hypertension</i> , <b>2004</b> , 18, 475-85	2.6	42
360	The combined impact of blood pressure category and glucose abnormality on the incidence of cardiovascular diseases in a Japanese urban cohort: the Suita Study. <i>Hypertension Research</i> , <b>2010</b> , 33, 1238-43	4.7	41
359	Much lower prevalence of coronary calcium detected by electron-beam computed tomography among men aged 40-49 in Japan than in the US, despite a less favorable profile of major risk factors. <i>International Journal of Epidemiology</i> , <b>2005</b> , 34, 173-9	7.8	41
358	Haplotype analyses of cholesteryl ester transfer protein gene promoter: a clue to an unsolved mystery of TaqIB polymorphism. <i>Journal of Molecular Medicine</i> , <b>2003</b> , 81, 246-55	5.5	41
357	Reliability of plasma polar metabolite concentrations in a large-scale cohort study using capillary electrophoresis-mass spectrometry. <i>PLoS ONE</i> , <b>2018</b> , 13, e0191230	3.7	40
356	Cardiovascular risk with and without antihypertensive drug treatment in the Japanese general population: participant-level meta-analysis. <i>Hypertension</i> , <b>2014</b> , 63, 1189-97	8.5	40
355	Risk of smoking and metabolic syndrome for incidence of cardiovascular disease--comparison of relative contribution in urban Japanese population: the Suita study. <i>Circulation Journal</i> , <b>2009</b> , 73, 2258-63 <sup>2.9</sup>	2.9	40
354	Coronary artery calcification in Japanese men in Japan and Hawaii. <i>American Journal of Epidemiology</i> , <b>2007</b> , 166, 1280-7	3.8	40
353	Long chain n-3 polyunsaturated fatty acids and incidence rate of coronary artery calcification in Japanese men in Japan and white men in the USA: population based prospective cohort study. <i>Heart</i> , <b>2014</b> , 100, 569-73	5.1	39
352	What factors are associated with high plasma B-type natriuretic peptide levels in a general Japanese population?. <i>Journal of Human Hypertension</i> , <b>2005</b> , 19, 165-72	2.6	39
351	LOX-1 in atherosclerotic disease. <i>Clinica Chimica Acta</i> , <b>2015</b> , 440, 157-63	6.2	38

350	Prognostic value of cardiac magnetic resonance imaging for idiopathic pulmonary arterial hypertension before initiating intravenous prostacyclin therapy. <i>Circulation Journal</i> , <b>2012</b> , 76, 1737-43	2.9	38
349	Multicenter study to determine the diagnosis criteria of heterozygous familial hypercholesterolemia in Japan. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2012</b> , 19, 1019-26	4	38
348	Comparison of low-density lipoprotein cholesterol concentrations measured by a direct homogeneous assay and by the Friedewald formula in a large community population. <i>Clinica Chimica Acta</i> , <b>2010</b> , 411, 1774-80	6.2	37
347	Differential association of docosahexaenoic and eicosapentaenoic acids with carotid intima-media thickness. <i>Stroke</i> , <b>2011</b> , 42, 2538-43	6.7	37
346	Lipoprotein-associated phospholipase A2 is related to risk of subclinical atherosclerosis but is not supported by Mendelian randomization analysis in a general Japanese population. <i>Atherosclerosis</i> , <b>2016</b> , 246, 141-7	3.1	36
345	Associations of obesity measures with metabolic risk factors in a community-based population in Japan. <i>Circulation Journal</i> , <b>2007</b> , 71, 776-81	2.9	36
344	Comprehensive risk management for the prevention of cardiovascular disease: executive summary of the Japan Atherosclerosis Society (JAS) guidelines for the diagnosis and prevention of atherosclerotic cardiovascular diseases in Japan -- 2012. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2013</b> , 20, 603-15	4	36
343	Having few remaining teeth is associated with a low nutrient intake and low serum albumin levels in middle-aged and older Japanese individuals: findings from the NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , <b>2019</b> , 24, 1	4.2	36
342	Low-carbohydrate diets and cardiovascular and total mortality in Japanese: a 29-year follow-up of NIPPON DATA80. <i>British Journal of Nutrition</i> , <b>2014</b> , 112, 916-24	3.6	35
341	Carotid intima-media thickness and plaque in apparently healthy Japanese individuals with an estimated 10-year absolute risk of CAD death according to the Japan Atherosclerosis Society (JAS) guidelines 2012: the Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2013</b> , 20, 755-66	4	35
340	Serum n-6 fatty acids and lipoprotein subclasses in middle-aged men: the population-based cross-sectional ERA-JUMP study. <i>American Journal of Clinical Nutrition</i> , <b>2010</b> , 91, 1195-203	7	35
339	The relationship between waist circumference and the risk of stroke and myocardial infarction in a Japanese urban cohort: the Suita study. <i>Stroke</i> , <b>2010</b> , 41, 550-3	6.7	35
338	Integration of data from NIPPON DATA80/90 and National Nutrition Survey in Japan: for cohort studies of representative Japanese on nutrition. <i>Journal of Epidemiology</i> , <b>2010</b> , 20 Suppl 3, S506-14	3.4	35
337	Body mass index and risk of stroke mortality among a random sample of Japanese adults: 19-year follow-up of NIPPON DATA80. <i>Cerebrovascular Diseases</i> , <b>2006</b> , 22, 409-15	3.2	35
336	Relation of adult height with stroke mortality in Japan: NIPPON DATA80. <i>Stroke</i> , <b>2007</b> , 38, 22-6	6.7	35
335	The proportion of individuals with alcohol-induced hypertension among total hypertensives in a general Japanese population: NIPPON DATA90. <i>Hypertension Research</i> , <b>2007</b> , 30, 663-8	4.7	35
334	Polymorphism of the angiotensin converting enzyme gene and blood pressure in a Japanese general population (the Shigaraki Study). <i>Hypertension Research</i> , <b>2002</b> , 25, 843-8	4.7	35
333	Lifetime risk of stroke in Japan. <i>Stroke</i> , <b>2010</b> , 41, 1552-4	6.7	34

332	Effectiveness of a low-intensity intra-worksite intervention on smoking cessation in Japanese employees: a three-year intervention trial. <i>Journal of Occupational Health</i> , <b>2006</b> , 48, 175-82	2.3	34
331	Metabolic Profiling of Total Physical Activity and Sedentary Behavior in Community-Dwelling Men. <i>PLoS ONE</i> , <b>2016</b> , 11, e0164877	3.7	34
330	The Relationship between Very High Levels of Serum High-Density Lipoprotein Cholesterol and Cause-Specific Mortality in a 20-Year Follow-Up Study of Japanese General Population. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2016</b> , 23, 800-9	4	33
329	Secular trends of the impact of overweight and obesity on hypertension in Japan, 1980-2010. <i>Hypertension Research</i> , <b>2015</b> , 38, 790-5	4.7	32
328	Association of serum n-6 and n-3 polyunsaturated fatty acids with lipids in 3 populations of middle-aged men. <i>American Journal of Clinical Nutrition</i> , <b>2009</b> , 90, 49-55	7	32
327	Non-invasive measurement of brachial-ankle pulse wave velocity is associated with serum C-reactive protein but not with alpha-tocopherol in Japanese middle-aged male workers. <i>Hypertension Research</i> , <b>2004</b> , 27, 173-80	4.7	32
326	Relationship among physical activity, smoking, drinking and clustering of the metabolic syndrome diagnostic components. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2010</b> , 17, 644-50	4	32
325	Combined effect of blood pressure and total cholesterol levels on long-term risks of subtypes of cardiovascular death: Evidence for Cardiovascular Prevention from Observational Cohorts in Japan. <i>Hypertension</i> , <b>2015</b> , 65, 517-24	8.5	31
324	Association of four genetic loci with uric acid levels and reduced renal function: the J-SHIP Suita study. <i>American Journal of Nephrology</i> , <b>2010</b> , 32, 279-86	4.6	31
323	The taste of salt measured by a simple test and blood pressure in Japanese women and men. <i>Hypertension Research</i> , <b>2009</b> , 32, 399-403	4.7	31
322	Relationship between BMI and all-cause mortality in Japan: NIPPON DATA80. <i>Obesity</i> , <b>2008</b> , 16, 1714-7	8	31
321	A "natural experiment" in cardiovascular epidemiology in the early 21st century. <i>British Heart Journal</i> , <b>2003</b> , 89, 255-7		31
320	Familial hypercholesterolemia. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 6-10	4	30
319	High oolong tea consumption predicts future risk of diabetes among Japanese male workers: a prospective cohort study. <i>Diabetic Medicine</i> , <b>2011</b> , 28, 805-10	3.5	30
318	Triglycerides and non-high-density lipoprotein cholesterol and the incidence of cardiovascular disease in an urban Japanese cohort: the Suita study. <i>Atherosclerosis</i> , <b>2010</b> , 209, 290-4	3.1	30
317	Life expectancy among Japanese of different smoking status in Japan: NIPPON DATA80. <i>Journal of Epidemiology</i> , <b>2007</b> , 17, 31-7	3.4	30
316	Stages of change for salt intake and urinary salt excretion: baseline results from the High-Risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) study. <i>Hypertension Research</i> , <b>2004</b> , 27, 157-66	4.7	30
315	Prospective Study on Alcohol Intake and Risk of Subarachnoid Hemorrhage Among Japanese Men and Women. <i>Alcoholism: Clinical and Experimental Research</i> , <b>2000</b> , 24, 386-389	3.7	30



314	A revised definition of the metabolic syndrome predicts coronary artery disease and ischemic stroke after adjusting for low density lipoprotein cholesterol in a 13-year cohort study of Japanese: the Suita study. <i>Atherosclerosis</i> , <b>2011</b> , 217, 201-6	3.1	29
313	The association between high-density lipoprotein cholesterol level and cholesteryl ester transfer protein TaqIB gene polymorphism is influenced by alcohol drinking in a population-based sample. <i>Atherosclerosis</i> , <b>2007</b> , 191, 199-205	3.1	29
312	Comparison of HOMA-IR, HOMA-% and disposition index between US white men and Japanese men in Japan: the ERA JUMP study. <i>Diabetologia</i> , <b>2015</b> , 58, 265-71	10.3	28
311	Alcohol consumption and cardiovascular disease incidence in men with and without hypertension: the Suita study. <i>Hypertension Research</i> , <b>2013</b> , 36, 58-64	4.7	28
310	Association between J-point elevation and death from coronary artery disease--15-year follow up of the NIPPON DATA90. <i>Circulation Journal</i> , <b>2013</b> , 77, 1260-6	2.9	28
309	A Japanese diet and 19-year mortality: national integrated project for prospective observation of non-communicable diseases and its trends in the aged, 1980. <i>British Journal of Nutrition</i> , <b>2009</b> , 101, 1696-705	3.6	28
308	Electrocardiogram screening for left high R-wave predicts cardiovascular death in a Japanese community-based population: NIPPON DATA90. <i>Hypertension Research</i> , <b>2006</b> , 29, 353-60	4.7	28
307	Profiling of plasma metabolites in postmenopausal women with metabolic syndrome. <i>Menopause</i> , <b>2016</b> , 23, 749-58	2.5	28
306	Effects of stroke education of junior high school students on stroke knowledge of their parents: Tochigi project. <i>Stroke</i> , <b>2015</b> , 46, 572-4	6.7	27
305	Does high-sensitivity C-reactive protein or low-density lipoprotein cholesterol show a stronger relationship with the cardio-ankle vascular index in healthy community dwellers?: the KOBE study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2012</b> , 19, 1027-34	4	27
304	Dyslipidemia and cardiovascular disease: a series of epidemiologic studies in Japanese populations. <i>Journal of Epidemiology</i> , <b>2010</b> , 20, 259-65	3.4	27
303	Joint impact of smoking and hypertension on cardiovascular disease and all-cause mortality in Japan: NIPPON DATA80, a 19-year follow-up. <i>Hypertension Research</i> , <b>2007</b> , 30, 1169-75	4.7	27
302	Prognostic significance of endogenous erythropoietin in long-term outcome of patients with acute decompensated heart failure. <i>European Journal of Heart Failure</i> , <b>2016</b> , 18, 803-13	12.3	26
301	Validation of homogeneous assays for HDL-cholesterol using fresh samples from healthy and diseased subjects. <i>Atherosclerosis</i> , <b>2014</b> , 233, 253-9	3.1	26
300	Relationship between shift work and clustering of the metabolic syndrome diagnostic components. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 703-11	4	26
299	A combination of serum low albumin and above-average cholesterol level was associated with excess mortality. <i>Journal of Clinical Epidemiology</i> , <b>2004</b> , 57, 1188-95	5.7	26
298	A cross-sectional study of alcohol drinking and health-related quality of life among male workers in Japan. <i>Journal of Occupational Health</i> , <b>2005</b> , 47, 496-503	2.3	26
297	Smoking, Smoking Cessation, and Measures of Subclinical Atherosclerosis in Multiple Vascular Beds in Japanese Men. <i>Journal of the American Heart Association</i> , <b>2016</b> , 5,	6	26

296	Significant inverse association of equol-producer status with coronary artery calcification but not dietary isoflavones in healthy Japanese men. <i>British Journal of Nutrition</i> , <b>2017</b> , 117, 260-266	3.6	25
295	Risk of hypercholesterolemia for cardiovascular disease and the population attributable fraction in a 24-year Japanese cohort study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2015</b> , 22, 95-107	4	25
294	Relation of dietary and other lifestyle traits to difference in serum adiponectin concentration of Japanese in Japan and Hawaii: the INTERLIPID Study. <i>American Journal of Clinical Nutrition</i> , <b>2008</b> , 88, 424-30	7	25
293	Plasma level of homocysteine is correlated to extracranial carotid-artery atherosclerosis in non-hypertensive Japanese. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , <b>1999</b> , 6, 371-7		25
292	Relationship of Estimated GFR and Albuminuria to Concurrent Laboratory Abnormalities: An Individual Participant Data Meta-analysis in a Global Consortium. <i>American Journal of Kidney Diseases</i> , <b>2019</b> , 73, 206-217	7.4	25
291	Smoking increases the risk of all-cause and cardiovascular mortality in patients with chronic kidney disease. <i>Kidney International</i> , <b>2015</b> , 88, 1144-52	9.9	24
290	Diagnosis of atherosclerosis. Executive Summary of the Japan Atherosclerosis Society (JAS) Guidelines for the Diagnosis and Prevention of Atherosclerotic Cardiovascular Diseases in Japan 2012 Version. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 296-8	4	24
289	Public awareness of early symptoms of stroke and information sources about stroke among the general Japanese population: the Acquisition of Stroke Knowledge Study. <i>Cerebrovascular Diseases</i> , <b>2013</b> , 35, 241-9	3.2	24
288	Association of carotid atherosclerosis with genetic polymorphisms of the klotho gene in patients with hypertension. <i>Geriatrics and Gerontology International</i> , <b>2010</b> , 10, 311-8	2.9	24
287	Risk for metabolic diseases in normal weight individuals with visceral fat accumulation: a cross-sectional study in Japan. <i>BMJ Open</i> , <b>2017</b> , 7, e013831	3	23
286	Coronary artery calcification by computed tomography in epidemiologic research and cardiovascular disease prevention. <i>Journal of Epidemiology</i> , <b>2012</b> , 22, 188-98	3.4	23
285	Comparison of predictability of future cardiovascular events between chronic kidney disease (CKD) stage based on CKD epidemiology collaboration equation and that based on modification of diet in renal disease equation in the Japanese general population--Iwate KENCO Study. <i>Circulation Journal</i> , <b>2013</b> , 77, 1815-25	2.9	23
284	Higher levels of adiponectin in American than in Japanese men despite obesity. <i>Metabolism: Clinical and Experimental</i> , <b>2006</b> , 55, 1561-3	12.7	23
283	Long-term outcome of healthy participants with atrial premature complex: a 15-year follow-up of the NIPPON DATA 90 cohort. <i>PLoS ONE</i> , <b>2013</b> , 8, e80853	3.7	23
282	Age-specific impact of diabetes mellitus on the risk of cardiovascular mortality: An overview from the evidence for Cardiovascular Prevention from Observational Cohorts in the Japan Research Group (EPOCH-JAPAN). <i>Journal of Epidemiology</i> , <b>2017</b> , 27, 123-129	3.4	22
281	Medical costs of obese Japanese: a 10-year follow-up study of National Health Insurance in Shiga, Japan. <i>European Journal of Public Health</i> , <b>2007</b> , 17, 424-9	2.1	22
280	Worksite wellness for the primary and secondary prevention of cardiovascular disease in Japan: the current delivery system and future directions. <i>Progress in Cardiovascular Diseases</i> , <b>2014</b> , 56, 515-21	8.5	21
279	Interaction of Blood Pressure and Body Mass Index With Risk of Incident Atrial Fibrillation in a Japanese Urban Cohort: The Suita Study. <i>American Journal of Hypertension</i> , <b>2015</b> , 28, 1355-61	2.3	21

278	Lifetime risk of acute myocardial infarction in Japan. <i>Circulation: Cardiovascular Quality and Outcomes</i> , <b>2010</b> , 3, 701-3	5.8	21
277	Prognostic values of clockwise and counterclockwise rotation for cardiovascular mortality in Japanese subjects: a 24-year follow-up of the National Integrated Project for Prospective Observation of Noncommunicable Disease and Its Trends in the Aged, 1980-2004 (NIPPON DATA80). <i>Circulation</i> , <b>2012</b> , 125, 1226-33	16.7	21
276	Fish consumption and early atherosclerosis in middle-aged men. <i>Metabolism: Clinical and Experimental</i> , <b>2007</b> , 56, 1060-4	12.7	21
275	The value of combining serum alanine aminotransferase levels and body mass index to predict mortality and medical costs: a 10-year follow-up study of National Health Insurance in Shiga, Japan. <i>Journal of Epidemiology</i> , <b>2006</b> , 16, 15-20	3.4	21
274	Relationship between Non-High-Density Lipoprotein Cholesterol and Low-Density Lipoprotein Cholesterol in the General Population. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2016</b> , 23, 477-90	4	21
273	Continuous decline in mortality from coronary heart disease in Japan despite a continuous and marked rise in total cholesterol: Japanese experience after the Seven Countries Study. <i>International Journal of Epidemiology</i> , <b>2015</b> , 44, 1614-24	7.8	20
272	Effect of chronic kidney disease on individual and population medical expenditures in the Japanese population. <i>Hypertension Research</i> , <b>2009</b> , 32, 450-4	4.7	20
271	Awareness and treatment of hypertension and hypercholesterolemia in Japanese workers: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) study. <i>Hypertension Research</i> , <b>2007</b> , 30, 921-8	4.7	20
270	Effect of combined cardiovascular risk factors on individual and population medical expenditures: a 10-year cohort study of national health insurance in a Japanese population. <i>Circulation Journal</i> , <b>2007</b> , 71, 807-13	2.9	20
269	Relationship between metabolic syndrome and Trp64arg polymorphism of the beta-adrenergic receptor gene in a general sample: the Shigaraki study. <i>Hypertension Research</i> , <b>2006</b> , 29, 891-6	4.7	20
268	The National Integrated Project for Prospective Observation of Non-communicable Disease and its Trends in the Aged 2010 (NIPPON DATA2010): Objectives, Design, and Population Characteristics. <i>Journal of Epidemiology</i> , <b>2018</b> , 28 Suppl 3, S2-S9	3.4	20
267	Cholesteryl ester transfer protein, coronary calcium, and intima-media thickness of the carotid artery in middle-age Japanese men. <i>American Journal of Cardiology</i> , <b>2009</b> , 104, 818-22	3	19
266	Alcohol consumption and coronary artery calcium in middle-aged Japanese men. <i>American Journal of Cardiology</i> , <b>2006</b> , 98, 141-4	3	19
265	Different effects of blood pressure on mortality from stroke subtypes depending on BMI levels: a 19-year cohort study in the Japanese general population--NIPPON DATA80. <i>Journal of Human Hypertension</i> , <b>2005</b> , 19, 285-91	2.6	19
264	Lifetime Risk of Stroke and Coronary Heart Disease Deaths According to Blood Pressure Level: EPOCH-JAPAN (Evidence for Cardiovascular Prevention From Observational Cohorts in Japan). <i>Hypertension</i> , <b>2019</b> , 73, 52-59	8.5	19
263	Relationship Between Socioeconomic Status and the Prevalence of Underweight, Overweight or Obesity in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , <b>2018</b> , 28 Suppl 3, S10-S16	3.4	19
262	Influence of cigarette smoking on coronary artery and aortic calcium among random samples from populations of middle-aged Japanese and Korean men. <i>Journal of Epidemiology and Community Health</i> , <b>2013</b> , 67, 119-24	5.1	18
261	Effect of age on the association between waist-to-height ratio and incidence of cardiovascular disease: the Suita study. <i>Journal of Epidemiology</i> , <b>2013</b> , 23, 351-9	3.4	18

260	Association study of 11beta-hydroxysteroid dehydrogenase type 1 gene polymorphisms and metabolic syndrome in urban Japanese cohort. <i>Diabetes Research and Clinical Practice</i> , <b>2009</b> , 85, 132-8	7.4	18
259	Association of the functional variant in the 3-hydroxy-3-methylglutaryl-coenzyme a reductase gene with low-density lipoprotein-cholesterol in Japanese. <i>Circulation Journal</i> , <b>2010</b> , 74, 518-22	2.9	18
258	Impact of hypertension on medical economics: A 10-year follow-up study of national health insurance in Shiga, Japan. <i>Hypertension Research</i> , <b>2005</b> , 28, 859-64	4.7	18
257	Lipoprotein particle profiles compared with standard lipids in association with coronary artery calcification in the general Japanese population. <i>Atherosclerosis</i> , <b>2014</b> , 236, 237-43	3.1	17
256	LOX-1 ligands containing apolipoprotein B and carotid intima-media thickness in middle-aged community-dwelling US Caucasian and Japanese men. <i>Atherosclerosis</i> , <b>2013</b> , 229, 240-5	3.1	17
255	Associations between Inflammatory Markers and Subclinical Atherosclerosis in Middle-aged White, Japanese-American and Japanese Men: The ERA-JUMP Study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2015</b> , 22, 590-8	4	17
254	Relationship between 5-year decline in instrumental activity of daily living and accumulation of cardiovascular risk factors: NIPPON DATA90. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2010</b> , 17, 64-72	4	17
253	Association with serum gamma-glutamyltransferase levels and alcohol consumption on stroke and coronary artery disease: the Suita study. <i>Stroke</i> , <b>2011</b> , 42, 1764-7	6.7	17
252	Intima-media thickness of the carotid artery and the distribution of lipoprotein subclasses in men aged 40 to 49 years between whites in the United States and the Japanese in Japan for the ERA JUMP study. <i>Metabolism: Clinical and Experimental</i> , <b>2008</b> , 57, 177-82	12.7	17
251	Plasma fibrinogen and coronary heart disease in urban Japanese. <i>American Journal of Epidemiology</i> , <b>2000</b> , 152, 420-3	3.8	17
250	Explaining the decline in coronary heart disease mortality rates in Japan: Contributions of changes in risk factors and evidence-based treatments between 1980 and 2012. <i>International Journal of Cardiology</i> , <b>2019</b> , 291, 183-188	3.2	16
249	Relationship between non-high-density lipoprotein cholesterol and the long-term mortality of cardiovascular diseases: NIPPON DATA 90. <i>International Journal of Cardiology</i> , <b>2016</b> , 220, 262-7	3.2	16
248	Impact of hypertension on the lifetime risk of coronary heart disease. <i>Hypertension Research</i> , <b>2016</b> , 39, 548-51	4.7	16
247	Blood pressure, low-density lipoprotein cholesterol, and incidences of coronary artery disease and ischemic stroke in Japanese: the Suita study. <i>American Journal of Hypertension</i> , <b>2014</b> , 27, 1362-9	2.3	16
246	High-density lipoprotein particle concentration and subclinical atherosclerosis of the carotid arteries in Japanese men. <i>Atherosclerosis</i> , <b>2015</b> , 239, 444-50	3.1	16
245	Treatment A) lifestyle modification: executive summary of the Japan Atherosclerosis Society(JAS) guidelines for the diagnosis and prevention of atherosclerotic cardiovascular diseases in Japan--2012 version. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2013</b> , 20, 835-49	4	16
244	Steeper increases in body mass index during childhood correlate with blood pressure elevation in adolescence: a long-term follow-up study in a Japanese community. <i>Hypertension Research</i> , <b>2014</b> , 37, 179-84	4.7	16
243	Defining patients at extremely high risk for coronary artery disease in heterozygous familial hypercholesterolemia. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2012</b> , 19, 369-75	4	16

242	Blood cholesterol level and risk of stroke in community-based or worksite cohort studies: a review of Japanese cohort studies in the past 20 years. <i>Keio Journal of Medicine</i> , <b>2012</b> , 61, 79-88	1.6	16
241	A cross-sectional association of obesity with coronary calcium among Japanese, Koreans, Japanese Americans, and U.S. whites. <i>European Heart Journal Cardiovascular Imaging</i> , <b>2013</b> , 14, 921-7	4.1	16
240	Gender and age effects on ventricular repolarization abnormality in Japanese general carriers of a G643S common single nucleotide polymorphism for the KCNQ1 gene. <i>Circulation Journal</i> , <b>2006</b> , 70, 645-50	3.0	16
239	Combined analysis of polymorphisms in angiotensinogen and adducin genes and their effects on hypertension in a Japanese sample: The Shigaraki Study. <i>Hypertension Research</i> , <b>2005</b> , 28, 645-50	4.7	16
238	Association of blood pressure with estimates of 24-h urinary sodium and potassium excretion from repeated single-spot urine samples. <i>Hypertension Research</i> , <b>2019</b> , 42, 411-418	4.7	15
237	Nationwide distribution of cardiovascular practice in Japan - results of Japanese circulation society 2010 annual survey. <i>Circulation Journal</i> , <b>2015</b> , 79, 1058-67	2.9	15
236	Relationship between health-related quality of life and clustering of metabolic syndrome diagnostic components. <i>Quality of Life Research</i> , <b>2012</b> , 21, 1165-70	3.7	15
235	Prognostic value of ST-T abnormalities and left high R waves with cardiovascular mortality in Japanese (24-year follow-up of NIPPON DATA80). <i>American Journal of Cardiology</i> , <b>2011</b> , 107, 1718-24	3	15
234	Relationship of the polyunsaturated to saturated fatty acid ratio to cardiovascular risk factors and metabolic syndrome in Japanese: the INTERLIPID study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2010</b> , 17, 777-84	4	15
233	Cigarette smoking in middle age and a long-term risk of impaired activities of daily living: NIPPON DATA80. <i>Nicotine and Tobacco Research</i> , <b>2010</b> , 12, 944-9	4.9	15
232	Circulating levels of 8 cytokines and marine n-3 fatty acids and indices of obesity in Japanese, white, and Japanese American middle-aged men. <i>Journal of Interferon and Cytokine Research</i> , <b>2010</b> , 30, 541-8	3.5	15
231	High blood pressure in middle age is associated with a future decline in activities of daily living. NIPPON DATA80. <i>Journal of Human Hypertension</i> , <b>2009</b> , 23, 546-52	2.6	15
230	Association of lipoprotein-associated phospholipase A2 with coronary calcification among American and Japanese men. <i>Journal of Epidemiology</i> , <b>2007</b> , 17, 179-85	3.4	15
229	Applicability of the stages of change model for analyzing fruit and vegetable intake in relation to urinary potassium excretion: baseline results from the High-Risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) Study. <i>Hypertension Research</i> , <b>2004</b> , 27, 843-50	4.7	15
228	Risk Factors Associated With Ischemic Stroke in Japanese Patients With Nonvalvular Atrial Fibrillation. <i>JAMA Network Open</i> , <b>2020</b> , 3, e202881	10.4	15
227	Serum Polyunsaturated Fatty Acid Composition and Serum High-Sensitivity C-Reactive Protein Levels in Healthy Japanese Residents: The KOBE Study. <i>Journal of Nutrition, Health and Aging</i> , <b>2015</b> , 19, 719-28	5.2	14
226	Effects of intensive and moderate public education on knowledge of early stroke symptoms among a Japanese population: the Acquisition of Stroke Knowledge study. <i>Stroke</i> , <b>2013</b> , 44, 2829-34	6.7	14
225	Increased childhood BMI is associated with young adult serum uric acid levels: a linkage study from Japan. <i>Pediatric Research</i> , <b>2017</b> , 81, 293-298	3.2	14

224	A pooled analysis of the association of isolated low levels of high-density lipoprotein cholesterol with cardiovascular mortality in Japan. <i>European Journal of Epidemiology</i> , <b>2017</b> , 32, 547-557	12.1	14
223	HOMA-IR Values are Associated With Glycemic Control in Japanese Subjects Without Diabetes or Obesity: The KOBE Study. <i>Journal of Epidemiology</i> , <b>2015</b> , 25, 407-14	3.4	14
222	The Relationship between Lectin-Like Oxidized Low-Density Lipoprotein Receptor-1 Ligands Containing Apolipoprotein B and the Cardio-Ankle Vascular Index in Healthy Community Inhabitants: The KOBE Study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2015</b> , 22, 499-508	4	14
221	The reasonable reliability of a self-administered food frequency questionnaire for an urban, Japanese, middle-aged population: the Suita study. <i>Nutrition Research</i> , <b>2015</b> , 35, 14-22	4	14
220	Treated and untreated hypertension, hospitalization, and medical expenditure: an epidemiological study in 314622 beneficiaries of the medical insurance system in Japan. <i>Journal of Hypertension</i> , <b>2013</b> , 31, 1032-42	1.9	14
219	Lifestyle and work predictors of fatigue in Japanese manufacturing workers. <i>Occupational Medicine</i> , <b>2007</b> , 57, 262-9	2.1	14
218	Marital status and cardiovascular risk factors among middle-aged Japanese male workers: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) study. <i>Journal of Occupational Health</i> , <b>2008</b> , 50, 348-56	2.3	14
217	Sex differences in impact of coronary artery calcification to predict coronary artery disease. <i>Heart</i> , <b>2018</b> , 104, 1118-1124	5.1	14
216	Fasting but not casual blood glucose is associated with pancreatic cancer mortality in Japanese: EPOCH-JAPAN. <i>Cancer Causes and Control</i> , <b>2017</b> , 28, 625-633	2.8	13
215	High risks of all-cause and cardiovascular deaths in apparently healthy middle-aged people with preserved glomerular filtration rate and albuminuria: A prospective cohort study. <i>International Journal of Cardiology</i> , <b>2013</b> , 170, 167-72	3.2	13
214	The prevalence of aortic calcification in Japanese compared to white and Japanese-American middle-aged men is confounded by the amount of cigarette smoking. <i>International Journal of Cardiology</i> , <b>2013</b> , 167, 134-9	3.2	13
213	Diabetes and life expectancy among Japanese - NIPPON DATA80. <i>Diabetes Research and Clinical Practice</i> , <b>2012</b> , 96, e18-22	7.4	13
212	Treatment B) drug therapy: executive summary of the Japan Atherosclerosis Society(JAS) guidelines for the diagnosis and prevention of atherosclerotic cardiovascular diseases in Japan--2012 version. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2013</b> , 20, 850-60	4	13
211	Elevated serum creatine kinase predicts first-ever myocardial infarction: a 12-year population-based cohort study in Japan, the Suita study. <i>International Journal of Epidemiology</i> , <b>2009</b> , 38, 1571-9	7.8	13
210	Hypertension and life expectancy among Japanese: NIPPON DATA80. <i>Hypertension Research</i> , <b>2012</b> , 35, 954-8	4.7	13
209	Fatty acids intakes and serum lipid profiles: NIPPON DATA90 and the national nutrition monitoring. <i>Journal of Epidemiology</i> , <b>2010</b> , 20 Suppl 3, S544-8	3.4	13
208	Population attributable fraction of smoking and metabolic syndrome on cardiovascular disease mortality in Japan: a 15-year follow up of NIPPON DATA90. <i>BMC Public Health</i> , <b>2010</b> , 10, 306	4.1	13
207	Difference in carotid intima-media thickness between Korean and Japanese men. <i>Annals of Epidemiology</i> , <b>2008</b> , 18, 310-5	6.4	13

206	Low-risk profile for cardiovascular disease and mortality in Japanese. <i>Circulation Journal</i> , <b>2008</b> , 72, 545-509		13
205	Continuation of smoking cessation and following weight change after intervention in a healthy population with high smoking prevalence. <i>Journal of Occupational Health</i> , <b>2006</b> , 48, 402-6	2.3	13
204	Diabetes mellitus. Executive summary of the Japan Atherosclerosis Society (JAS) guidelines for the diagnosis and prevention of atherosclerotic cardiovascular diseases in Japan--2012 version. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 93-8	4	13
203	Underweight Young Women Without Later Weight Gain Are at High Risk for Osteopenia After Midlife: The KOBE Study. <i>Journal of Epidemiology</i> , <b>2016</b> , 26, 572-578	3.4	13
202	Cross-sectional association between exposure to particulate matter and inflammatory markers in the Japanese general population: NIPPON DATA2010. <i>Environmental Pollution</i> , <b>2016</b> , 213, 460-467	9.3	13
201	Association between dyslipidemia and plasma levels of branched-chain amino acids in the Japanese population without diabetes mellitus. <i>Journal of Clinical Lipidology</i> , <b>2019</b> , 13, 932-939.e2	4.9	12
200	Re-evaluation of the associations of egg intake with serum total cholesterol and cause-specific and total mortality in Japanese women. <i>European Journal of Clinical Nutrition</i> , <b>2018</b> , 72, 841-847	5.2	12
199	Effects of a Public Education Campaign on the Association Between Knowledge of Early Stroke Symptoms and Intention to Call an Ambulance at Stroke Onset: The Acquisition of Stroke Knowledge (ASK) Study. <i>Journal of Epidemiology</i> , <b>2016</b> , 26, 115-22	3.4	12
198	Cumulative impact of axial, structural, and repolarization ECG findings on long-term cardiovascular mortality among healthy individuals in Japan: National Integrated Project for Prospective Observation of Non-Communicable Disease and its Trends in the Aged, 1980 and 1990. <i>European Journal of Preventive Cardiology</i> , <b>2014</b> , 21, 1501-8	3.9	12
197	Bundle branch block and other cardiovascular disease risk factors: US-Japan comparison. <i>International Journal of Cardiology</i> , <b>2010</b> , 143, 432-40	3.2	12
196	Prognostic value of q wave for cardiovascular death in a 19-year prospective study of the Japanese general population. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2009</b> , 16, 40-50	4	12
195	Association of Total Energy Intake with 29-Year Mortality in the Japanese: NIPPON DATA80. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2016</b> , 23, 339-54	4	12
194	Vegetable Protein Intake was Inversely Associated with Cardiovascular Mortality in a 15-Year Follow-Up Study of the General Japanese Population. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2019</b> , 26, 198-206	4	12
193	Variation in process quality measures of diabetes care by region and institution in Japan during 2015-2016: An observational study of nationwide claims data. <i>Diabetes Research and Clinical Practice</i> , <b>2019</b> , 155, 107750	7.4	11
192	Coronary artery disease. Executive summary of the Japan Atherosclerosis Society (JAS) guidelines for the diagnosis and prevention of atherosclerotic cardiovascular diseases in Japan--2012 version. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 86-92	4	11
191	Metabolic syndrome. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 1-5	4	11
190	The clustering of cardiovascular disease risk factors and their impacts on annual medical expenditure in Japan: community-based cost analysis using Gamma regression models. <i>BMJ Open</i> , <b>2013</b> , 3,	3	11
189	Cardiovascular disease risk factors other than dyslipidemia. Executive summary of the Japan Atherosclerosis Society (JAS) guidelines for the diagnosis and prevention of atherosclerotic cardiovascular diseases in Japan - 2012 version. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2013</b> , 20, 733-42	4	11

188	Relationship between sleep duration and clustering of metabolic syndrome diagnostic components. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , <b>2011</b> , 4, 119-25	3.4	11
187	New diagnosis criteria for diabetes with hemoglobin A1c and risks of macro-vascular complications in an urban Japanese cohort: the Suita study. <i>Diabetes Research and Clinical Practice</i> , <b>2010</b> , 88, e20-3	7.4	11
186	The proportion of individuals with obesity-induced hypertension among total hypertensives in a general Japanese population: NIPPON DATA80, 90. <i>European Journal of Epidemiology</i> , <b>2007</b> , 22, 691-8	12.1	11
185	Reference Intervals of Serum Non-Cholesterol Sterols by Gender in Healthy Japanese Individuals. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2020</b> , 27, 409-417	4	11
184	Macronutrient Intake and Socioeconomic Status: NIPPON DATA2010. <i>Journal of Epidemiology</i> , <b>2018</b> , 28 Suppl 3, S17-S22	3.4	11
183	Socioeconomic Status Associated With Urinary Sodium and Potassium Excretion in Japan: NIPPON DATA2010. <i>Journal of Epidemiology</i> , <b>2018</b> , 28 Suppl 3, S29-S34	3.4	11
182	Risk of stroke and heart failure attributable to atrial fibrillation in middle-aged and elderly people: Results from a five-year prospective cohort study of Japanese community dwellers. <i>Journal of Epidemiology</i> , <b>2017</b> , 27, 360-367	3.4	10
181	HDL cholesterol performance using an ultracentrifugation reference measurement procedure and the designated comparison method. <i>Clinica Chimica Acta</i> , <b>2015</b> , 439, 185-90	6.2	10
180	Serum level of LOX-1 ligand containing ApoB is associated with increased carotid intima-media thickness in Japanese community-dwelling men, especially those with hypercholesterolemia LOX-1 ligand and IMT in Japanese. <i>Journal of Clinical Lipidology</i> , <b>2016</b> , 10, 172-80.e1	4.9	10
179	Prognostic values of bundle branch blocks for cardiovascular mortality in Japanese (24year follow-up of NIPPON DATA80). <i>Journal of Electrocardiology</i> , <b>2013</b> , 46, 360-5	1.4	10
178	Effects of a school-based stroke education program on stroke-related knowledge and behaviour modification-school class based intervention study for elementary school students and parental guardians in a Japanese rural area. <i>BMJ Open</i> , <b>2017</b> , 7, e017632	3	10
177	Changes in Waist Circumference and the Incidence of Type 2 Diabetes in Community-Dwelling Men and Women: The Suita Study. <i>Journal of Epidemiology</i> , <b>2015</b> , 25, 489-95	3.4	10
176	The relationship between Asian dust events and out-of-hospital cardiac arrests in Japan. <i>Journal of Epidemiology</i> , <b>2015</b> , 25, 289-96	3.4	10
175	Alcohol intake and 19-year mortality in diabetic men: NIPPON DATA80. <i>Alcohol</i> , <b>2009</b> , 43, 635-41	2.7	10
174	Normal fasting plasma glucose levels and type 2 diabetes: the high-risk and population strategy for occupational health promotion (HIPOP-OHP) [corrected] study. <i>Acta Diabetologica</i> , <b>2007</b> , 44, 164-6	3.9	10
173	Both angiotensinogen M235T and alpha-adducin G460W polymorphisms are associated with hypertension in the Japanese population. <i>Journal of Human Hypertension</i> , <b>2007</b> , 21, 253-5	2.6	10
172	Medical costs of patients with hypertension and/or diabetes: A 10-year follow-up study of National Health Insurance in Shiga, Japan. <i>Journal of Hypertension</i> , <b>2006</b> , 24, 2305-9	1.9	10
171	Employment and health after retirement in Japanese men. <i>Bulletin of the World Health Organization</i> , <b>2018</b> , 96, 826-833	8.2	10



170	Absolute risk of cardiovascular disease and lipid management targets. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2013</b> , 20, 689-97	4	10
169	Dietary Inflammatory Index Positively Associated With High-Sensitivity C-Reactive Protein Level in Japanese From NIPPON DATA2010. <i>Journal of Epidemiology</i> , <b>2020</b> , 30, 98-107	3.4	10
168	Socioeconomic Inequalities in Oral Health among Middle-Aged and Elderly Japanese: NIPPON DATA2010. <i>Journal of Epidemiology</i> , <b>2018</b> , 28 Suppl 3, S59-S65	3.4	10
167	Associations between Socioeconomic Status and the Prevalence and Treatment of Hypercholesterolemia in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2018</b> , 25, 606-620	4	10
166	Birthweight and cardiovascular risk factors in a Japanese general population. <i>Journal of Obstetrics and Gynaecology Research</i> , <b>2017</b> , 43, 1001-1007	1.9	9
165	Socioeconomic and lifestyle factors associated with depressive tendencies in general Japanese men and women: NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , <b>2019</b> , 24, 37	4.2	9
164	Long-term outcomes associated with prolonged PR interval in the general Japanese population. <i>International Journal of Cardiology</i> , <b>2015</b> , 184, 291-293	3.2	9
163	Relationship of type of work with health-related quality of life. <i>Quality of Life Research</i> , <b>2015</b> , 24, 2927-337		9
162	Development of a Cardiovascular Disease Risk Prediction Model Using the Suita Study, a Population-Based Prospective Cohort Study in Japan. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2020</b> , 27, 1160-1175	4	9
161	Fatty liver index predicts incident diabetes in a Japanese general population with and without impaired fasting glucose. <i>Hepatology Research</i> , <b>2018</b> , 48, 708-716	5.1	9
160	Associations of serum LDL particle concentration with carotid intima-media thickness and coronary artery calcification. <i>Journal of Clinical Lipidology</i> , <b>2016</b> , 10, 1195-1202.e1	4.9	9
159	High long-chain n-3 fatty acid intake attenuates the effect of high resting heart rate on cardiovascular mortality risk: a 24-year follow-up of Japanese general population. <i>Journal of Cardiology</i> , <b>2014</b> , 64, 218-24	3	9
158	Relation of Serum Leptin and Adiponectin Level to Serum C-Reactive Protein: The INTERLIPID Study. <i>International Journal of Vascular Medicine</i> , <b>2013</b> , 2013, 601364	1.2	9
157	Medical expenditures of men with hypertension and/or a smoking habit: a 10-year follow-up study of National Health Insurance in Shiga, Japan. <i>Hypertension Research</i> , <b>2010</b> , 33, 802-7	4.7	9
156	Relationships between protein intake and renal function in a Japanese general population: NIPPON DATA90. <i>Journal of Epidemiology</i> , <b>2010</b> , 20 Suppl 3, S537-43	3.4	9
155	Serum Eglutamyltransferase and Mortality due to Cardiovascular Disease in Japanese Men and Women. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2016</b> , 23, 792-9	4	9
154	Relationship between outpatient visit frequency and hypertension control: a 9-year occupational cohort study. <i>Hypertension Research</i> , <b>2016</b> , 39, 376-81	4.7	9
153	Associations of High-Density Lipoprotein Particle and High-Density Lipoprotein Cholesterol With Alcohol Intake, Smoking, and Body Mass Index - The INTERLIPID Study. <i>Circulation Journal</i> , <b>2018</b> , 82, 2557-2565 <sup>9</sup>	2.9	9

152	Socioeconomic Status and Knowledge of Cardiovascular Risk Factors: NIPPON DATA2010. <i>Journal of Epidemiology</i> , <b>2018</b> , 28 Suppl 3, S46-S52	3.4	9
151	Diabetes and lifetime risk of coronary heart disease. <i>Primary Care Diabetes</i> , <b>2017</b> , 11, 461-466	2.4	8
150	Higher serum uric acid level is inversely associated with renal function assessed by cystatin C in a Japanese general population without chronic kidney disease: the KOBE study. <i>BMC Nephrology</i> , <b>2019</b> , 20, 117	2.7	8
149	Longitudinal Trajectories of Fasting Plasma Glucose and Risks of Cardiovascular Diseases in Middle Age to Elderly People Within the General Japanese Population: The Suita Study. <i>Journal of the American Heart Association</i> , <b>2019</b> , 8, e010628	6	8
148	Standardized Prevalence Ratios for Atrial Fibrillation in Adult Dialysis Patients in Japan. <i>Journal of Epidemiology</i> , <b>2016</b> , 26, 272-6	3.4	8
147	Effect size and cost-effectiveness estimates of breast and cervical cancer screening reminders by population size through complete enumeration of Japanese local municipalities. <i>BMC Public Health</i> , <b>2014</b> , 14, 43	4.1	8
146	Association between serum long-chain n-3 and n-6 polyunsaturated fatty acid profiles and glomerular filtration rate assessed by serum creatinine and cystatin C levels in Japanese community-dwellers. <i>Journal of Epidemiology</i> , <b>2015</b> , 25, 303-11	3.4	8
145	Association of total marine fatty acids, eicosapentaenoic and docosahexaenoic acids, with aortic stiffness in Koreans, whites, and Japanese Americans. <i>American Journal of Hypertension</i> , <b>2013</b> , 26, 1321-7	3.3	8
144	Relationship of moderate metabolic risk factor clustering to cardiovascular disease mortality in non-lean Japanese: a 15-year follow-up of NIPPON DATA90. <i>Atherosclerosis</i> , <b>2011</b> , 215, 209-13	3.1	8
143	Relation of Gamma-glutamyltransferase and alcohol drinking with incident diabetes: the HIPOP-OHP study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2010</b> , 17, 195-202	4	8
142	Protein intakes and serum albumin levels in a Japanese general population: NIPPON DATA90. <i>Journal of Epidemiology</i> , <b>2010</b> , 20 Suppl 3, S531-6	3.4	8
141	Relationships between family histories of stroke and of hypertension and stroke mortality: NIPPON DATA80, 1980-1999. <i>Hypertension Research</i> , <b>2008</b> , 31, 1525-31	4.7	8
140	Effects of the Y chromosome on cardiovascular risk factors in Japanese men. <i>Hypertension Research</i> , <b>2008</b> , 31, 1687-94	4.7	8
139	Is weak association between cigarette smoking and cardiovascular disease mortality observed in Japan explained by low total cholesterol? NIPPON DATA80. <i>International Journal of Epidemiology</i> , <b>2007</b> , 36, 1060-7	7.8	8
138	Methodological issues for a large-scale intervention trial of lifestyle modification: Interim assessment of the high-risk and population strategy for occupational health promotion (HIPOP-OHP) study. <i>Environmental Health and Preventive Medicine</i> , <b>2004</b> , 9, 137-43	4.2	8
137	The evaluation of materials to provide health-related information as a population strategy in the worksite: The high-risk and population strategy for occupational health promotion (HIPOP-OHP) study. <i>Environmental Health and Preventive Medicine</i> , <b>2004</b> , 9, 144-51	4.2	8
136	Determinants of platelet aggregation in 50-70-year-old men from three Japanese communities. <i>Atherosclerosis</i> , <b>2002</b> , 165, 327-34	3.1	8
135	Hypertension and medical expenditure in the Japanese population: Review of prospective studies. <i>World Journal of Cardiology</i> , <b>2014</b> , 6, 531-8	2.1	8

134	Does the flushing response modify the relationship between alcohol intake and hypertension in the Japanese population? NIPPON DATA2010. <i>Hypertension Research</i> , <b>2016</b> , 39, 670-9	4.7	8
133	Association Between Body Mass Index and All-Cause Death in Japanese Population: Pooled Individual Participant Data Analysis of 13 Cohort Studies. <i>Journal of Epidemiology</i> , <b>2019</b> , 29, 457-463	3.4	8
132	A Novel Cell-Free, Non-Fluorescent Method to Measure LOX-1-Binding Activity Corresponding to The Functional Activity of HDL. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2019</b> , 26, 947-958	4	7
131	Dietary tofu intake and long-term risk of death from stroke in a general population. <i>Clinical Nutrition</i> , <b>2018</b> , 37, 182-188	5.9	7
130	Ethnic difference in liver fat content: a cross-sectional observation among Japanese American in Hawaii, Japanese in Japan, and non-Hispanic whites in United States. <i>Obesity Research and Clinical Practice</i> , <b>2013</b> , 7, e198-205	5.4	7
129	Relative and absolute risks of all-cause and cause-specific deaths attributable to atrial fibrillation in middle-aged and elderly community dwellers. <i>International Journal of Cardiology</i> , <b>2015</b> , 184, 692-698	3.2	7
128	Association of short sleep duration with impaired glucose tolerance or diabetes mellitus. <i>Journal of Diabetes Investigation</i> , <b>2011</b> , 2, 366-72	3.9	7
127	Association of intima-media thickening of carotid artery with genetic polymorphisms of the regulator of G-protein signaling 2 gene in patients with hypertension and in the general population. <i>Hypertension Research</i> , <b>2011</b> , 34, 740-6	4.7	7
126	Comparison of the National Nutritional Survey in Japan estimated individual-based nutritional data and NIPPON DATA80 food frequency questionnaires. <i>Journal of Epidemiology</i> , <b>2010</b> , 20 Suppl 3, S582-6	3.4	7
125	Serum ghrelin levels are higher in Caucasian men than Japanese men aged 40-49 years. <i>Diabetes, Obesity and Metabolism</i> , <b>2007</b> , 9, 591-3	6.7	7
124	Are comprehensive environmental changes as effective as health education for smoking cessation?. <i>Tobacco Control</i> , <b>2006</b> , 15, 26-9	5.3	7
123	Smoking scenes in popular Japanese serial television dramas: descriptive analysis during the same 3-month period in two consecutive years. <i>Health Promotion International</i> , <b>2006</b> , 21, 98-103	3	7
122	The determinants of plasma plasminogen activator inhibitor-1 levels differ for American and Japanese men aged 40-49. <i>Diabetes Research and Clinical Practice</i> , <b>2006</b> , 72, 176-82	7.4	7
121	Isolated systolic hypertension and 29-year cardiovascular mortality risk in Japanese adults aged 30-49 years. <i>Journal of Hypertension</i> , <b>2020</b> , 38, 2230-2236	1.9	7
120	Statin Intolerance Clinical Guide 2018. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2020</b> , 27, 375-396	4	7
119	The impact of sex on risk of cardiovascular disease and all-cause mortality in adults with or without diabetes mellitus: A comparison between the U.S. and Japan. <i>Journal of Diabetes and Its Complications</i> , <b>2019</b> , 33, 417-423	3.2	6
118	Spreading Awareness of Stroke through School-Based Education: A Pooled Analysis of Three Community-Based Studies. <i>Journal of Stroke and Cerebrovascular Diseases</i> , <b>2018</b> , 27, 1810-1814	2.8	6
117	The Associations of C-Reactive Protein with Serum Levels of Polyunsaturated Fatty Acids and Trans Fatty Acids Among Middle-Aged Men from Three Populations. <i>Journal of Nutrition, Health and Aging</i> , <b>2016</b> , 20, 16-21	5.2	6

116	Relationship of three different types of low-carbohydrate diet to cardiometabolic risk factors in a Japanese population: the INTERMAP/INTERLIPID Study. <i>European Journal of Nutrition</i> , <b>2016</b> , 55, 1515-24	5.2	6
115	Association between impairment of salty taste recognition and masked hypertension based on home blood pressure in Japanese residents: the KOBE study. <i>Hypertension Research</i> , <b>2018</b> , 41, 756-762	4.7	6
114	Interaction between dietary marine-derived n-3 fatty acids intake and J-point elevation on the risk of cardiac death: a 24-year follow-up of Japanese men. <i>Heart</i> , <b>2013</b> , 99, 1024-9	5.1	6
113	Medical expenditure for diabetic patients: a 10-year follow-up study of National Health Insurance in Shiga, Japan. <i>Public Health</i> , <b>2008</b> , 122, 1226-8	4	6
112	Association of Alcohol Consumption With Fat Deposition in a Community-Based Sample of Japanese Men: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Epidemiology</i> , <b>2019</b> , 29, 205-212	3.4	6
111	Relationship of household salt intake level with long-term all-cause and cardiovascular disease mortality in Japan: NIPPON DATA80. <i>Hypertension Research</i> , <b>2020</b> , 43, 132-139	4.7	6
110	Calibration between the Estimated Probability of the Risk Assessment Chart of Japan Atherosclerosis Society and Actual Mortality Using External Population: Evidence for Cardiovascular Prevention from Observational Cohorts in Japan (EPOCH-JAPAN). <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2016</b> , 23, 176-95	4	6
109	Validation of the european SCORE risk chart in the healthy middle-aged Japanese. <i>Atherosclerosis</i> , <b>2016</b> , 252, 116-121	3.1	6
108	Impacts of chronic kidney disease and diabetes on cardiovascular mortality in a general Japanese population: A 20-year follow-up of the NIPPON DATA90 study. <i>European Journal of Preventive Cardiology</i> , <b>2017</b> , 24, 505-513	3.9	5
107	Cardiovascular Risk Assessment Chart by Dietary Factors in Japan - NIPPON DATA80. <i>Circulation Journal</i> , <b>2019</b> , 83, 1254-1260	2.9	5
106	International Comparison of Abdominal Fat Distribution Among Four Populations: The ERA-JUMP Study. <i>Metabolic Syndrome and Related Disorders</i> , <b>2018</b> , 16, 166-173	2.6	5
105	Overweight or underweight and the risk of decline in activities of daily living in a 22-year cohort study of a Japanese sample. <i>Geriatrics and Gerontology International</i> , <b>2018</b> , 18, 799-805	2.9	5
104	Cardiovascular risk and blood pressure lowering treatment among elderly individuals: Evidence for Cardiovascular Prevention from Observational Cohorts in Japan. <i>Journal of Hypertension</i> , <b>2018</b> , 36, 410-418	1.9	5
103	Cardiometabolic index as a useful discriminator for the risk of increased arterial stiffness. <i>Clinica Chimica Acta</i> , <b>2018</b> , 486, 42-43	6.2	5
102	Serum leptin and total dietary energy intake: the INTERLIPID Study. <i>European Journal of Nutrition</i> , <b>2013</b> , 52, 1641-8	5.2	5
101	γGlutamyltransferase and mortality risk from heart disease and stroke in Japanese men and women: NIPPON DATA90. <i>CVD Prevention and Control</i> , <b>2010</b> , 5, 27-34		5
100	Dietary habits in middle age and future changes in activities of daily living - NIPPON DATA80. <i>Gerontology</i> , <b>2009</b> , 55, 707-13	5.5	5
99	Impairments in activities of daily living in older Japanese men in hawaii and Japan. <i>Journal of Aging Research</i> , <b>2011</b> , 2011, 324592	2.3	5

98	What unfavorable factors are associated with low serum total cholesterol in a Japanese population?. <i>Journal of Epidemiology</i> , <b>2002</b> , 12, 271-9	3.4	5
97	Hypercholesterolemia and Lifetime Risk of Coronary Heart Disease in the General Japanese Population: Results from the Suita Cohort Study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2020</b> , 27, 60-70	4	5
96	Relationships among Food Group Intakes, Household Expenditure, and Education Attainment in a General Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , <b>2018</b> , 28 Suppl 3, S23-S28	3.4	5
95	Diabetes and lifetime risk of stroke and subtypes in an urban middle-aged population. <i>Journal of Diabetes and Its Complications</i> , <b>2017</b> , 31, 831-835	3.2	4
94	Association of Work Situation With Cardiovascular Disease Mortality Risk Among Working-Age Japanese Men - A 20-Year Follow-up of NIPPON DATA90. <i>Circulation Journal</i> , <b>2019</b> , 83, 1506-1513	2.9	4
93	Concordance of CKD stages in estimation by the CKD-EPI equation and estimation by the MDRD equation in the Japanese general population: the Iwate KENCO Study. <i>International Journal of Cardiology</i> , <b>2013</b> , 165, 377-9	3.2	4
92	Risk factors for heart failure and coronary heart disease mortality over 24-year follow-up period in Japan: NIPPON DATA80. <i>CVD Prevention and Control</i> , <b>2010</b> , 5, 97-103		4
91	Relation of serum leptin to blood pressure of Japanese in Japan and Japanese-Americans in Hawaii. <i>Hypertension</i> , <b>2009</b> , 54, 1416-22	8.5	4
90	Risk factor effects and total mortality in older Japanese men in Japan and Hawaii. <i>Annals of Epidemiology</i> , <b>2008</b> , 18, 913-8	6.4	4
89	Does self-reported history of hypertension predict cardiovascular death? Comparison with blood pressure measurement in a 19-year prospective study. <i>Journal of Hypertension</i> , <b>2007</b> , 25, 959-64	1.9	4
88	Residual lifetime risk of cardiovascular diseases in Japan. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2011</b> , 18, 443-7	4	4
87	The association of home and accurately measured office blood pressure with coronary artery calcification among general Japanese men. <i>Journal of Hypertension</i> , <b>2019</b> , 37, 1676-1681	1.9	4
86	Relationship between carbohydrate and dietary fibre intake and the risk of cardiovascular disease mortality in Japanese: 24-year follow-up of NIPPON DATA80. <i>European Journal of Clinical Nutrition</i> , <b>2020</b> , 74, 67-76	5.2	4
85	A Combination of Blood Pressure and Total Cholesterol Increases the Lifetime Risk of Coronary Heart Disease Mortality: EPOCH-JAPAN. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2021</b> , 28, 6-24	4	4
84	Association of alcohol consumption and aortic calcification in healthy men aged 40-49 years for the ERA JUMP Study. <i>Atherosclerosis</i> , <b>2018</b> , 268, 84-91	3.1	4
83	Factors Related to Participation in Health Examinations for Japanese National Health Insurance: NIPPON DATA2010. <i>Journal of Epidemiology</i> , <b>2018</b> , 28 Suppl 3, S53-S58	3.4	4
82	Estimation of 10-Year Risk of Death from Coronary Heart Disease, Stroke, and Cardiovascular Disease in a Pooled Analysis of Japanese Cohorts: EPOCH-JAPAN. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2021</b> , 28, 816-825	4	4
81	Left ventricular hypertrophy by electrocardiogram as a predictor of success in home blood pressure control: HOMED-BP study. <i>Hypertension Research</i> , <b>2017</b> , 40, 504-510	4.7	3

80	One-year weight loss maintenance outcomes following a worksite-based weight reduction program among Japanese men with cardiovascular risk factors. <i>Journal of Occupational Health</i> , <b>2019</b> , 61, 189-196 <sup>2,3</sup>	3
79	Impact of hypertension stratified by diabetes on the lifetime risk of cardiovascular disease mortality in Japan: a pooled analysis of data from the Evidence for Cardiovascular Prevention from Observational Cohorts in Japan study. <i>Hypertension Research</i> , <b>2020</b> , 43, 1437-1444	4.7 3
78	Associations of Overweight, Obesity, and Underweight With High Serum Total Cholesterol Level Over 30 Years Among the Japanese Elderly: NIPPON DATA 80, 90, and 2010. <i>Journal of Epidemiology</i> , <b>2019</b> , 29, 133-138	3.4 3
77	Association of Asn221Ser mutation in tissue factor pathway inhibitor-beta with plasma total tissue factor pathway inhibitor level. <i>Blood Coagulation and Fibrinolysis</i> , <b>2009</b> , 20, 22-6	1 3
76	Independent Prognostic Value of Single and Multiple Non-Specific 12-Lead Electrocardiographic Findings for Long-Term Cardiovascular Outcomes: A Prospective Cohort Study. <i>PLoS ONE</i> , <b>2016</b> , 11, e0157563 <sup>3</sup>	3.7 563 3
75	Pooled Analysis of the Associations between Body Mass Index, Total Cholesterol, and Liver Cancer-related Mortality in Japan. <i>Asian Pacific Journal of Cancer Prevention</i> , <b>2018</b> , 19, 2089-2095	1.7 3
74	Chronic kidney disease. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 173-4	4 3
73	Estimated 24 h Urinary Sodium-to-Potassium Ratio Is Related to Renal Function Decline: A 6-Year Cohort Study of Japanese Urban Residents. <i>International Journal of Environmental Research and Public Health</i> , <b>2020</b> , 17,	4.6 3
72	Association between socioeconomic status and prolonged television viewing time in a general Japanese population: NIPPON DATA2010. <i>Environmental Health and Preventive Medicine</i> , <b>2021</b> , 26, 57	4.2 3
71	A Novel Risk Stratification System for Ischemic Stroke in Japanese Patients With Non-Valvular Atrial Fibrillation. <i>Circulation Journal</i> , <b>2021</b> , 85, 1254-1262	2.9 3
70	Validity Assessment of Self-reported Medication Use for Hypertension, Diabetes, and Dyslipidemia in a Pharmacoepidemiologic Study by Comparison With Health Insurance Claims. <i>Journal of Epidemiology</i> , <b>2021</b> , 31, 495-502	3.4 3
69	Detection of Subclinical Peripheral Artery Ischemia in Healthy Male Smokers by an Ankle-Brachial Index After Exercise: Sasayama Study. <i>Angiology</i> , <b>2017</b> , 68, 769-775	2.1 2
68	Data on alcohol consumption and coronary artery calcification among asymptomatic middle-aged men for the ERA-JUMP study. <i>Data in Brief</i> , <b>2018</b> , 17, 1091-1098	1.2 2
67	Being Conscious of Water Intake Positively Associated with Sufficient Non-Alcohol Drink Intake Regardless of Seasons and Reasons in Healthy Japanese; the KOBE Study: A Cross Sectional Study. <i>International Journal of Environmental Research and Public Health</i> , <b>2019</b> , 16,	4.6 2
66	Women. Executive Summary of the Japan Atherosclerosis Society (JAS) Guidelines for the Diagnosis and Prevention of Atherosclerotic Cardiovascular Diseases in Japan 2012 Version. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 291-5	4 2
65	Correlation of a self-report and direct measure of physical activity level in the electron-beam tomography and risk assessment among Japanese and US Men in the post World War II birth cohort (ERA JUMP) study. <i>Journal of Epidemiology</i> , <b>2013</b> , 23, 411-7	3.4 2
64	Fasting plasma insulin is associated with metabolic syndrome in farmers but not in nomads among the Mongolian population, China. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2011</b> , 18, 291-7	4 2
63	Relationship between Dietary and Other Lifestyle Habits and Cardiometabolic Risk Factors in Men. <i>Diabetology and Metabolic Syndrome</i> , <b>2011</b> , 3, 30	5.6 2

62	Medical costs of individuals with proteinuria: A 10-year follow-up study of National Health Insurance in Shiga, Japan. <i>Public Health</i> , <b>2007</b> , 121, 174-6	4	2
61	The Perinatal Condition Around Birth and Cardiovascular Risk Factors in the Japanese General Population: The Suita Study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2020</b> , 27, 204-214	4	2
60	Metabolic profiling of charged metabolites in association with menopausal status in Japanese community-dwelling midlife women: Tsuruoka Metabolomic Cohort Study. <i>Maturitas</i> , <b>2022</b> , 155, 54-62	5	2
59	Fatty Acid Intakes and Coronary Heart Disease Mortality in Japan: NIPPON DATA90, 1990-2005. <i>Current Nutrition and Food Science</i> , <b>2013</b> , 9, 26-32	0.7	2
58	Other types of primary hyperlipoproteinemia(hyperlipidemia). Executive summary of the Japan Atherosclerosis Society (JAS) guidelines for the diagnosis and prevention of atherosclerotic cardiovascular diseases in Japan--2012 version. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 82-5	4	2
57	Impact of Hypertension on Hospitalizations for Cardiovascular Diseases in a Worksite Population: An Epidemiologic Study Using Claims Data for Workers. <i>American Journal of Hypertension</i> , <b>2019</b> , 32, 298-307	3.7	2
56	Differences between home blood pressure and strictly measured office blood pressure and their determinants in Japanese men. <i>Hypertension Research</i> , <b>2021</b> , 44, 80-87	4.7	2
55	The Risk of Fasting Triglycerides and its Related Indices for Ischemic Cardiovascular Diseases in Japanese Community Dwellers: the Suita Study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2021</b> , 28, 1273-1288 <sup>2</sup>	4	2
54	Relationship of higher-level functional capacity with long-term mortality in Japanese older people: NIPPON DATA90. <i>Journal of Epidemiology</i> , <b>2021</b> ,	3.4	2
53	Alcohol drinking and brain morphometry in apparently healthy community-dwelling Japanese men. <i>Alcohol</i> , <b>2021</b> , 90, 57-65	2.7	2
52	Differences in Lifestyle Improvements With the Intention to Prevent Cardiovascular Diseases by Socioeconomic Status in a Representative Japanese Population: NIPPON DATA2010. <i>Journal of Epidemiology</i> , <b>2018</b> , 28 Suppl 3, S35-S39	3.4	2
51	Time-Related Changes in Relationships Between the Keys Score, Dietary Lipids, and Serum Total Cholesterol in Japan - NIPPON DATA80/90/2010. <i>Circulation Journal</i> , <b>2018</b> , 83, 147-155	2.9	2
50	Association between socioeconomic status and physical inactivity in a general Japanese population: NIPPON DATA2010. <i>PLoS ONE</i> , <b>2021</b> , 16, e0254706	3.7	2
49	Daily Habit of Water Intake in Patients with Cerebral Infarction before its Onset; Comparison with a Healthy Population: A Cross-Sectional Study. <i>Cerebrovascular Diseases</i> , <b>2019</b> , 47, 143-150	3.2	1
48	Electrocardiographic ST-T Abnormalities Are Associated With Stroke Risk in the REGARDS Study. <i>Stroke</i> , <b>2020</b> , 51, 1100-1106	6.7	1
47	Casual serum triglyceride as a predictor of premature type 2 diabetes mellitus: an 8-year cohort study of middle-aged Japanese workers. <i>Diabetology International</i> , <b>2016</b> , 7, 252-258	2.3	1
46	Impact of care coordination on oral anticoagulant therapy among patients with atrial fibrillation in routine clinical practice in Japan: a prospective, observational study. <i>BMC Cardiovascular Disorders</i> , <b>2019</b> , 19, 235	2.3	1
45	Alcohol consumption, hospitalization and medical expenditure: a large epidemiological study on the medical insurance system in Japan. <i>Alcohol and Alcoholism</i> , <b>2015</b> , 50, 236-43	3.5	1

44	Cerebrovascular diseases. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 175-9	4	1
43	Relation of dietary and lifestyle traits to difference in serum leptin of Japanese in Japan and Hawaii: the INTERLIPID study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , <b>2012</b> , 22, 14-22	4.5	1
42	Fatty Acid Intakes and Coronary Heart Disease Mortality in Japan: NIPPON DATA90, 1990-2005. <i>Current Nutrition and Food Science</i> , <b>2013</b> , 9, 26-32	0.7	1
41	Association between polymorphism of the AGTR1 and cardiovascular events in a Japanese general sample (The Shigaraki Study). <i>International Journal of Cardiology</i> , <b>2009</b> , 136, 354-5	3.2	1
40	Cardiovascular Risk Clustering With Obesity: A Good Target to Reduce Medical Expenditures as a First Step of High-Risk Approach in Communities and Worksites. <i>Circulation Journal</i> , <b>2007</b> , 71, 1667	2.9	1
39	Physique at Birth and Cardiovascular Disease Risk Factors in Japanese Urban Residents: the KOBE Study. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2020</b> ,	4	1
38	Guidelines on the Clinical Evaluation of Medicinal Products for Treatment of Dyslipidemia. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2020</b> , 27, 1246-1254	4	1
37	The elderly. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 180-5	4	1
36	What psychosocial characteristics are associated with smoking cessation behavior and readiness to quit smoking among Japanese male ever-smokers with type 2 diabetes mellitus?. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2010</b> , 17, 361-8	4	1
35	The relationship between serum levels of LOX-1 ligand containing ApoAI as a novel marker of dysfunctional HDL and coronary artery calcification in middle-aged Japanese men. <i>Atherosclerosis</i> , <b>2020</b> , 313, 20-25	3.1	1
34	Reliability of urinary charged metabolite concentrations in a large-scale cohort study using capillary electrophoresis-mass spectrometry. <i>Scientific Reports</i> , <b>2021</b> , 11, 7407	4.9	1
33	Risk Factors That Most Accurately Predict Coronary Artery Disease Based on the Duration of Follow-up - NIPPON DATA80. <i>Circulation Journal</i> , <b>2021</b> , 85, 908-913	2.9	1
32	Relationship of urinary liver-type fatty acid-binding protein with cardiovascular risk factors in the Japanese population without chronic kidney disease: Sasayama study. <i>BMC Nephrology</i> , <b>2021</b> , 22, 189	2.7	1
31	Echocardiographic Parameters and the Risk of Incident Atrial Fibrillation: The Suita Study. <i>Journal of Epidemiology</i> , <b>2020</b> , 30, 183-187	3.4	1
30	Electrocardiographic Left Atrial Abnormality and B-Type Natriuretic Peptide in a General Japanese Population: NIPPON DATA2010. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2021</b> , 28, 34-43	4	1
29	Association of the Estimated Coronary Artery Incidence Risk According to the Japan Atherosclerosis Society Guidelines 2017 with Cardio-Ankle Vascular Index. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2021</b> , 28, 1266-1274	4	1
28	Relationship Between Calcium Intake and Impaired Activities of Daily Living in a Japanese Population: NIPPON DATA90. <i>Journal of Epidemiology</i> , <b>2021</b> , 31, 119-124	3.4	1
27	Tracking Pattern of Total Cholesterol Levels from Childhood to Adolescence in Japan. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2021</b> ,	4	1



26	Trends in Prevalence, Treatment, and Control of Hypertension According to 40-Year-Old Life Expectancy at Prefectures in Japan from the National Health and Nutrition Surveys.. <i>Nutrients</i> , <b>2022</b> , 14,	6.7	1
25	Neurologically favourable outcomes of cardiogenic out-of-hospital cardiac arrest with relation to the type of witness in Japan. <i>Resuscitation</i> , <b>2020</b> , 152, 97-104	4	0
24	Predictors of lower limb fractures in general Japanese: NIPPON DATA90.. <i>PLoS ONE</i> , <b>2022</b> , 17, e0261716	3.7	0
23	Prediction of Lifetime Risk of Cardiovascular Disease Deaths Stratified by Sex in the Japanese Population. <i>Journal of the American Heart Association</i> , <b>2021</b> , 10, e021753	6	0
22	Risk Classification for Metabolic Syndrome and the Incidence of Cardiovascular Disease in Japan With Low Prevalence of Obesity: A Pooled Analysis of 10 Prospective Cohort Studies. <i>Journal of the American Heart Association</i> , <b>2021</b> , 10, e020760	6	0
21	Exercise Habits Are Associated with Improved Long-Term Mortality Risks in the Nationwide General Japanese Population: A 20-Year Follow-Up of the NIPPON DATA90 Study. <i>Tohoku Journal of Experimental Medicine</i> , <b>2020</b> , 252, 253-262	2.4	0
20	Acceleration of opportunistic atrial fibrillation screening for elderly patients in routine primary care. <i>PLoS ONE</i> , <b>2020</b> , 15, e0244240	3.7	0
19	Long-Term Exposure to Particulate Matter and Mortality from Cardiovascular Diseases in Japan: The Ibaraki Prefectural Health Study (IPHS). <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2021</b> , 28, 230-240 <sup>4</sup>		0
18	Comparison of the Japan Society of Clinical Chemistry reference method and CDC method for HDL and LDL cholesterol measurements using fresh sera. <i>Practical Laboratory Medicine</i> , <b>2021</b> , 25, e00228	1.7	0
17	Accumulated unhealthy behaviours and insomnia in Japanese dwellers with and without cardiovascular risk factors: a cross-sectional study.. <i>BMJ Open</i> , <b>2022</b> , 12, e052787	3	0
16	Relationship between thin physique at 6 years and metabolic disease risks in middle-aged Japanese women: The Suita study. <i>Journal of Obstetrics and Gynaecology Research</i> , <b>2020</b> , 46, 517-526	1.9	
15	Is Childhood Exposure to Parental Smoking a Risk Factor for Future Cardiovascular Disease?. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2017</b> , 24, 1204-1205	4	
14	Carotid Intima-Media Thickness and Plaque in Apparently Healthy Japanese Individuals with an Estimated 10-Year Absolute Risk of CAD Death According to the Japan Atherosclerosis Society (JAS) Guidelines 2012: The Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA). <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2019</b> , 26, 746	4	
13	PO3-1PRELIMINARY RESULTS ON THE IMPLEMENTATION OF THE SCREENING, BRIEF INTERVENTION, AND REFERRAL TO TREATMENT (SBRT) APPROACH IN A PORTUGUESE EMERGENCY DEPARTMENT. <i>Alcohol and Alcoholism</i> , <b>2017</b> , 52, i31-i49	3.5	
12	Familial Hypercholesterolemia. <i>Journal of Atherosclerosis and Thrombosis</i> , <b>2014</b> , 21, 289-289	4	
11	P1-530 Blood pressure categories and long-term mortality risk of total and components of cardiovascular diseases among Japanese: a 24 year follow-up of representative Japanese (NIPPON DATA80). <i>Journal of Epidemiology and Community Health</i> , <b>2011</b> , 65, A213-A214	5.1	
10	Risk-prediction tools for cardiovascular disease based on Japanese cohort studies. <i>Hypertension Research</i> , <b>2009</b> , 32, 1053-4	4.7	
9	P1-257 Relationship between the number of established cardiovascular risk factors and annual medical costs in Japan: cost analysis in 38 890 Japanese individuals. <i>Journal of Epidemiology and Community Health</i> , <b>2011</b> , 65, A137-A137	5.1	

- 8 Pros and cons of evidence-based medicine from the viewpoint of epidemiology. *Nosotchu*, **2008**, 30, 920-924
- 7 Lifestyle behaviors associated with the initiation of renal replacement therapy in Japanese patients with chronic kidney disease: a retrospective cohort study using a claims database linked with specific health checkup results. *Environmental Health and Preventive Medicine*, **2021**, 26, 102 4.2
- 6 518-P: Association between Plasma Metabolites in the Urea Cycle and Diabetic Kidney Disease (DKD) in a Japanese Population. *Diabetes*, **2020**, 69, 518-P 0.9
- 5 Lipoprotein particles and coronary artery calcium in middle-aged US-White and Japanese men. *Open Heart*, **2019**, 6, e001119 3
- 4 Response to the letter entitled "Extremely high levels of high-density lipoprotein cholesterol and increased risk of cardiovascular mortality" by Dr Kawada. *Journal of Clinical Lipidology*, **2018**, 12, 1551 4.9
- 3 A18797 Lifetime Risk of Stroke and Coronary Heart Disease Death according to Blood Pressure Level. *Journal of Hypertension*, **2018**, 36, e92-e93 1.9
- 2 Validity of a selective recall method for assessing water intake and its relationship with hydration status. *European Review for Medical and Pharmacological Sciences*, **2021**, 25, 6623-6632 2.9
- 1 Tooth Loss Related with Prevalence of Metabolic Syndrome in a General Urban Japanese Population: The Suita Study. *International Journal of Environmental Research and Public Health*, **2022**, 19, 6441 4.6