

Nobuaki Matsumori

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7417777/nobuaki-matsumori-publications-by-year.pdf>

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

140 papers	4,612 citations	33 h-index	63 g-index
158 ext. papers	5,113 ext. citations	4.9 avg, IF	5.22 L-index

#	Paper	IF	Citations
140	Recent advances in microscale separation techniques for lipidome analysis. <i>Analyst, The</i> , 2021 , 146, 7418-7430	5.7	2
139	Preparation of Nitrogen Analogues of Ceramide and Studies of Their Aggregation in Sphingomyelin Bilayers. <i>Langmuir</i> , 2021 , 37, 12438-12446	4	0
138	Metal Complex Lipids for Fluid-Fluid Phase Separation in Coassembled Phospholipid Membranes. <i>Angewandte Chemie - International Edition</i> , 2021 , 60, 13603-13608	16.4	2
137	Metal Complex Lipids for Fluid-Fluid Phase Separation in Coassembled Phospholipid Membranes. <i>Angewandte Chemie</i> , 2021 , 133, 13715-13720	3.6	
136	Amphidinol 3 preferentially binds to cholesterol in disordered domains and disrupts membrane phase separation. <i>Biochemistry and Biophysics Reports</i> , 2021 , 26, 100941	2.2	2
135	Low-flux scanning electron diffraction reveals substructures inside the ordered membrane domain. <i>Scientific Reports</i> , 2020 , 10, 22188	4.9	1
134	Pseudo-Membrane Jackets: Two-Dimensional Coordination Polymers Achieving Visible Phase Separation in Cell Membrane. <i>Angewandte Chemie - International Edition</i> , 2020 , 59, 17931-17937	16.4	6
133	Biophysics at Kyushu University. <i>Biophysical Reviews</i> , 2020 , 12, 245-247	3.7	0
132	Total Synthesis of Amphidinol 3: A General Strategy for Synthesizing Amphidinol Analogues and Structure-Activity Relationship Study. <i>Journal of the American Chemical Society</i> , 2020 , 142, 3472-3478	16.4	14
131	Chemical diversity and mode of action of natural products targeting lipids in the eukaryotic cell membrane. <i>Natural Product Reports</i> , 2020 , 37, 677-702	15.1	15
130	Defining raft domains in the plasma membrane. <i>Traffic</i> , 2020 , 21, 106-137	5.7	39
129	Assembly formation of minor dihydrosphingomyelin in sphingomyelin-rich ordered membrane domains. <i>Scientific Reports</i> , 2020 , 10, 11794	4.9	2
128	Pseudo-Membrane Jackets: Two-Dimensional Coordination Polymers Achieving Visible Phase Separation in Cell Membrane. <i>Angewandte Chemie</i> , 2020 , 132, 18087-18093	3.6	2
127	Archaeal Glycolipid S-TGA-1 Is Crucial for Trimer Formation and Photocycle Activity of Bacteriorhodopsin. <i>ACS Chemical Biology</i> , 2020 , 15, 197-204	4.9	2
126	The influence of ceramide and its dihydro analog on the physico-chemical properties of sphingomyelin bilayers. <i>Chemistry and Physics of Lipids</i> , 2020 , 226, 104835	3.7	5
125	Mechanism of local anesthetic-induced disruption of raft-like ordered membrane domains. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2019 , 1863, 1381-1389	4	7
124	The Perpendicular Orientation of Amphotericin B Methyl Ester in Hydrated Lipid Bilayers Supports the Barrel-Stave Model. <i>Biochemistry</i> , 2019 , 58, 2282-2291	3.2	11

123	A concise method for quantitative analysis of interactions between lipids and membrane proteins. <i>Analytica Chimica Acta</i> , 2019 , 1059, 103-112	6.6	9
122	Theonellamide A, a marine-sponge-derived bicyclic peptide, binds to cholesterol in aqueous DMSO: Solution NMR-based analysis of peptide-sterol interactions using hydroxylated sterol. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2019 , 1861, 228-235	3.8	6
121	Cholesterol-Induced Conformational Change in the Sphingomyelin Headgroup. <i>Biophysical Journal</i> , 2019 , 117, 307-318	2.9	8
120	The Amphotericin B-Ergosterol Complex Spans a Lipid Bilayer as a Single-Length Assembly. <i>Biochemistry</i> , 2019 , 58, 5188-5196	3.2	9
119	Preparation and Membrane Distribution of Fluorescent Derivatives of Ceramide. <i>Langmuir</i> , 2019 , 35, 2392-2398	4	6
118	Synthesis and Stereochemical Revision of the C31-C67 Fragment of Amphidinol 3. <i>Angewandte Chemie</i> , 2018 , 130, 6168-6172	3.6	4
117	Synthesis and Stereochemical Revision of the C31-C67 Fragment of Amphidinol 3. <i>Angewandte Chemie - International Edition</i> , 2018 , 57, 6060-6064	16.4	18
116	Preparation and Membrane Properties of Oxidized Ceramide Derivatives. <i>Langmuir</i> , 2018 , 34, 465-471	4	4
115	Synthesis and Complete Structure Determination of a Sperm-Activating and -Attracting Factor Isolated from the Ascidian <i>Ascidia sydneiensis</i> . <i>Journal of Natural Products</i> , 2018 , 81, 985-997	4.9	7
114	Dynamic membrane interactions of antibacterial and antifungal biomolecules, and amyloid peptides, revealed by solid-state NMR spectroscopy. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2018 , 1862, 307-323	4	29
113	Evidence of lipid rafts based on the partition and dynamic behavior of sphingomyelins. <i>Chemistry and Physics of Lipids</i> , 2018 , 215, 84-95	3.7	17
112	Recent Solid-State NMR Studies of Hydrated Lipid Membranes. <i>Annual Reports on NMR Spectroscopy</i> , 2018 , 41-72	1.7	2
111	Excellent Fluorescent Sphingomyelin Analog Reveals the Existence of Lipid Rafts. <i>Seibutsu Butsuri</i> , 2018 , 58, 321-323	0	
110	NMR Studies on Natural Product Stereochemical Determination and Conformational Analysis in Solution and in Membrane 2018 , 383-414		1
109	On the Importance of the C(1)-OH and C(3)-OH Functional Groups of the Long-Chain Base of Ceramide for Interlipid Interaction and Lateral Segregation into Ceramide-Rich Domains. <i>Langmuir</i> , 2018 , 34, 15864-15870	4	7
108	Sphingomyelin Stereoisomers Reveal That Homophilic Interactions Cause Nanodomain Formation. <i>Biophysical Journal</i> , 2018 , 115, 1530-1540	2.9	16
107	Raft-based sphingomyelin interactions revealed by new fluorescent sphingomyelin analogs. <i>Journal of Cell Biology</i> , 2017 , 216, 1183-1204	7.3	79
106	Channel Formation and Membrane Deformation via Sterol-Aided Polymorphism of Amphidinol 3. <i>Scientific Reports</i> , 2017 , 7, 10782	4.9	12

105	Emphatic visualization of sphingomyelin-rich domains by inter-lipid FRET imaging using fluorescent sphingomyelins. <i>Scientific Reports</i> , 2017 , 7, 16801	4.9	11
104	The impact of metal complex lipids on viscosity and curvature of hybrid liposomes. <i>Chemical Communications</i> , 2017 , 53, 13249-13252	5.8	10
103	Structures of the Largest Amphidinol Homologues from the Dinoflagellate <i>Amphidinium carterae</i> and Structure-Activity Relationships. <i>Journal of Natural Products</i> , 2017 , 80, 2883-2888	4.9	23
102	Sterol-dependent membrane association of the marine sponge-derived bicyclic peptide Theonellamide A as examined by H NMR. <i>Bioorganic and Medicinal Chemistry</i> , 2016 , 24, 5235-5242	3.4	6
101	Membrane protein structure determination by SAD, SIR, or SIRAS phasing in serial femtosecond crystallography using an iododetergent. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016 , 113, 13039-13044	11.5	38
100	Marine sponge cyclic peptide theonellamide A disrupts lipid bilayer integrity without forming distinct membrane pores. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2016 , 1858, 1373-9	3.8	19
99	(13) C-TmDOTA as versatile thermometer compound for solid-state NMR of hydrated lipid bilayer membranes. <i>Magnetic Resonance in Chemistry</i> , 2016 , 54, 227-33	2.1	1
98	The Structure of the Bimolecular Complex between Amphotericin B and Ergosterol in Membranes Is Stabilized by Face-to-Face van der Waals Interaction with Their Rigid Cyclic Cores. <i>Biochemistry</i> , 2016 , 55, 3392-402	3.2	16
97	Lipid Interactions and Organization in Complex Bilayer Membranes. <i>Biophysical Journal</i> , 2016 , 110, 1563-1573	4.73	19
96	Orientation and Order of the Amide Group of Sphingomyelin in Bilayers Determined by Solid-State NMR. <i>Biophysical Journal</i> , 2015 , 108, 2816-24	2.9	21
95	Stereoselective synthesis of the head group of archaeal phospholipid PGP-Me to investigate bacteriorhodopsin-lipid interactions. <i>Organic and Biomolecular Chemistry</i> , 2015 , 13, 10279-84	3.9	7
94	Role of polyol moiety of amphotericin B in ion channel formation and sterol selectivity in bilayer membrane. <i>Bioorganic and Medicinal Chemistry</i> , 2015 , 23, 5782-8	3.4	9
93	Deuterium NMR of raft model membranes reveals domain-specific order profiles and compositional distribution. <i>Biophysical Journal</i> , 2015 , 108, 2502-2506	2.9	44
92	Phosphatidylcholine bearing 6,6-dideuterated oleic acid: a useful solid-state (2)H NMR probe for investigating membrane properties. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2015 , 25, 203-6	2.9	8
91	Bioactive Structure of Membrane Lipids and Natural Products Elucidated by a Chemistry-Based Approach. <i>Chemical Record</i> , 2015 , 15, 675-90	6.6	14
90	Modification of bafilomycin structure to efficiently synthesize solid-state NMR probes that selectively bind to vacuolar-type ATPase. <i>Chemistry - an Asian Journal</i> , 2015 , 10, 915-24	4.5	7
89	Novel Raman-tagged sphingomyelin that closely mimics original raft-forming behavior. <i>Bioorganic and Medicinal Chemistry</i> , 2015 , 23, 2989-94	3.4	11
88	Formation of Gel-like Nanodomains in Cholesterol-Containing Sphingomyelin or Phosphatidylcholine Binary Membrane As Examined by Fluorescence Lifetimes and (2)H NMR Spectra. <i>Langmuir</i> , 2015 , 31, 13783-92	4	19

87	Axial hydrogen at C7 position and bumpy tetracyclic core markedly reduce sterols affinity to amphotericin B in membrane. <i>Biochemistry</i> , 2015 , 54, 303-12	3.2	13
86	Effect of sterol side chain on ion channel formation by amphotericin B in lipid bilayers. <i>Biochemistry</i> , 2014 , 53, 3088-94	3.2	14
85	Direct and stereospecific interaction of amphidinol 3 with sterol in lipid bilayers. <i>Biochemistry</i> , 2014 , 53, 3287-93	3.2	31
84	Detailed comparison of deuterium quadrupole profiles between sphingomyelin and phosphatidylcholine bilayers. <i>Biophysical Journal</i> , 2014 , 106, 631-8	2.9	49
83	Structure and biological activity of 8-deoxyheronamide C from a marine-derived <i>Streptomyces</i> sp.: heronamides target saturated hydrocarbon chains in lipid membranes. <i>Journal of the American Chemical Society</i> , 2014 , 136, 5209-12	16.4	47
82	Design and Synthesis of 24-Fluorinated Bafilomycin Analogue as an NMR Probe with Potent Inhibitory Activity to Vacuolar-type ATPase. <i>Chemistry Letters</i> , 2014 , 43, 474-476	1.7	5
81	Coexistence of two liquid crystalline phases in dihydrosphingomyelin and dioleoylphosphatidylcholine binary mixtures. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2014 , 1838, 1372-81	3.8	14
80	Structure and Interaction in Lipid Bilayers Analyzed Using Bicelles. <i>Yuki Gosei Kagaku Kyokaiishi/Journal of Synthetic Organic Chemistry</i> , 2014 , 72, 596-603	0.2	
79	A novel sperm-activating and attracting factor from the ascidian <i>Ascidia sydneiensis</i> . <i>Organic Letters</i> , 2013 , 15, 294-7	6.2	15
78	Synthesis and structure revision of the C43-C67 part of amphidinol 3. <i>Organic Letters</i> , 2013 , 15, 2846-9	6.2	25
77	Interaction between the marine sponge cyclic peptide theonellamide A and sterols in lipid bilayers as viewed by surface plasmon resonance and solid-state (2)H nuclear magnetic resonance. <i>Biochemistry</i> , 2013 , 52, 2410-8	3.2	38
76	Characterization of the ordered phase formed by sphingomyelin analogues and cholesterol binary mixtures. <i>Biophysics (Nagoya-shi, Japan)</i> , 2013 , 9, 37-49		9
75	NMR-based conformational analysis of sphingomyelin in bicelles. <i>Bioorganic and Medicinal Chemistry</i> , 2012 , 20, 270-8	3.4	26
74	Artificial ladder-shaped polyethers that inhibit maitotoxin-induced Ca ²⁺ influx in rat glioma C6 cells. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2012 , 22, 3619-22	2.9	8
73	Confirmation of the absolute configuration at C45 of amphidinol 3. <i>Journal of Natural Products</i> , 2012 , 75, 2003-6	4.9	17
72	Head-to-tail interaction between amphotericin B and ergosterol occurs in hydrated phospholipid membrane. <i>Biochemistry</i> , 2012 , 51, 83-9	3.2	30
71	Effects of chemical modification of sphingomyelin ammonium group on formation of liquid-ordered phase. <i>Bioorganic and Medicinal Chemistry</i> , 2012 , 20, 4012-9	3.4	7
70	Possible conformation of amphotericin B dimer in membrane-bound assembly as deduced from solid-state NMR. <i>Bioorganic and Medicinal Chemistry</i> , 2012 , 20, 5699-704	3.4	3

69	Comprehensive molecular motion capture for sphingomyelin by site-specific deuterium labeling. <i>Biochemistry</i> , 2012 , 51, 8363-70	3.2	46
68	An Approach Toward Identification of Target Proteins of Maitotoxin Based on Organic Synthesis 2012 , 23-35		
67	Channels formed by amphotericin B covalent dimers exhibit rectification. <i>Journal of Membrane Biology</i> , 2011 , 240, 159-64	2.3	11
66	Conformations of spermine in adenosine triphosphate complex: the structural basis for weak bimolecular interactions of major cellular electrolytes. <i>Chemistry - A European Journal</i> , 2011 , 17, 4788-95	4.8	4
65	Design and synthesis of sphingomyelin-cholesterol conjugates and their formation of ordered membranes. <i>Chemistry - A European Journal</i> , 2011 , 17, 8568-75	4.8	8
64	Fluorinated cholesterol retains domain-forming activity in sphingomyelin bilayers. <i>Chemistry and Physics of Lipids</i> , 2011 , 164, 401-8	3.7	12
63	Synthesis of 6-F-ergosterol and its influence on membrane-permeabilization of amphotericin B and amphidinol 3. <i>Organic and Biomolecular Chemistry</i> , 2011 , 9, 1437-42	3.9	8
62	Lysine proximity significantly affects glycation of lysine-containing collagen model peptides. <i>Bioorganic and Medicinal Chemistry</i> , 2011 , 19, 2125-9	3.4	3
61	Structural Reevaluations of Amphidinol 3, a Potent Antifungal Compound from Dinoflagellate. <i>Heterocycles</i> , 2010 , 82, 1359	0.8	3
60	3D structures of membrane-associated small molecules as determined in isotropic bicelles. <i>Natural Product Reports</i> , 2010 , 27, 1480-92	15.1	25
59	Sterol effect on interaction between amphidinol 3 and liposomal membrane as evidenced by surface plasmon resonance. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2010 , 20, 2215-8	2.9	23
58	Detection of Rap1A as a yessotoxin binding protein from blood cell membranes. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2010 , 20, 6443-6	2.9	9
57	Ion channel complex of antibiotics as viewed by NMR. <i>Pure and Applied Chemistry</i> , 2009 , 81, 1123-1129	2.1	14
56	Conformational change of spermidine upon interaction with adenosine triphosphate in aqueous solution. <i>Chemistry - A European Journal</i> , 2009 , 15, 1618-26	4.8	7
55	Amphotericin B-induced ion flux is markedly attenuated in phosphatidylglycerol membrane as evidenced by a newly devised fluorometric method. <i>Bioorganic and Medicinal Chemistry</i> , 2009 , 17, 6301-4	2.4	9
54	Surface plasmon resonance-based detection of ladder-shaped polyethers by inhibition detection method. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009 , 19, 2824-8	2.9	14
53	Direct interaction between amphotericin B and ergosterol in lipid bilayers as revealed by ² H NMR spectroscopy. <i>Journal of the American Chemical Society</i> , 2009 , 131, 11855-60	16.4	54
52	Synthesis of 25- ¹³ C-Amphotericin B Methyl Ester: A Molecular Probe for Solid-state NMR Measurements. <i>Chemistry Letters</i> , 2009 , 38, 114-115	1.7	7

51	Roles of integral protein in membrane permeabilization by amphidinols. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2008 , 1778, 1453-9	3.8	17
50	Orientation of fluorinated cholesterol in lipid bilayers analyzed by ¹⁹ F tensor calculation and solid-state NMR. <i>Journal of the American Chemical Society</i> , 2008 , 130, 4757-66	16.4	22
49	Structure of membrane-bound amphidinol 3 in isotropic small bicelles. <i>Organic Letters</i> , 2008 , 10, 4191-4	6.2	30
48	Design and synthesis of ladder-shaped tetracyclic, heptacyclic, and decacyclic ethers and evaluation of the interaction with transmembrane proteins. <i>Journal of the American Chemical Society</i> , 2008 , 130, 10217-26	16.4	30
47	Combinatorial synthesis of the 1,5-polyol system based on cross metathesis: structure revision of amphidinol 3. <i>Organic Letters</i> , 2008 , 10, 5203-6	6.2	56
46	Complex formation of amphotericin B in sterol-containing membranes as evidenced by surface plasmon resonance. <i>Biochemistry</i> , 2008 , 47, 7807-15	3.2	56
45	Convergent synthesis and biological activity of the WXYZABCSring system of maitotoxin. <i>Organic Letters</i> , 2008 , 10, 3599-602	6.2	33
44	Ergosterol increases the intermolecular distance of amphotericin B in the membrane-bound assembly as evidenced by solid-state NMR. <i>Biochemistry</i> , 2008 , 47, 13463-9	3.2	33
43	Structural Features of Dinoflagellate Toxins Underlying Biological Activity as Viewed by NMR. <i>Bulletin of the Chemical Society of Japan</i> , 2008 , 81, 307-319	5.1	24
42	Accurate Measurement of Vicinal Carbon-Hydrogen Coupling Constants via Ammonium Nitrogen Based on HMBC Experiments. <i>Chemistry Letters</i> , 2008 , 37, 1172-1173	1.7	3
41	Self-assembled amphotericin B is probably surrounded by ergosterol: bimolecular interactions as evidenced by solid-state NMR and CD spectra. <i>Chemistry - A European Journal</i> , 2008 , 14, 1178-85	4.8	36
40	Effects of lipid constituents on membrane-permeabilizing activity of amphidinols. <i>Bioorganic and Medicinal Chemistry</i> , 2008 , 16, 3084-90	3.4	33
39	Interaction of ladder-shaped polyethers with transmembrane alpha-helix of glycophorin A as evidenced by saturation transfer difference NMR and surface plasmon resonance. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2008 , 18, 6115-8	2.9	16
38	Conformation and position of membrane-bound amphotericin B deduced from NMR in SDS micelles. <i>Journal of Organic Chemistry</i> , 2007 , 72, 700-6	4.2	16
37	Conformation and location of membrane-bound salinomycin-sodium complex deduced from NMR in isotropic bicelles. <i>Journal of the American Chemical Society</i> , 2007 , 129, 14989-95	16.4	38
36	Assignment of the absolute configuration of blasticidin A and revision of that of aflastatin A. <i>Tetrahedron Letters</i> , 2007 , 48, 2527-2531	2	17
35	Amphotericin B covalent dimers with carbonyl-amino linkage: a new probe for investigating ion channel assemblies. <i>Tetrahedron Letters</i> , 2007 , 48, 3393-3396	2	15
34	Large molecular assembly of amphotericin B formed in ergosterol-containing membrane evidenced by solid-state NMR of intramolecular bridged derivative. <i>Journal of the American Chemical Society</i> , 2006 , 128, 11977-84	16.4	26

33	Membrane interaction of amphotericin B as single-length assembly examined by solid state NMR for uniformly ^{13}C -enriched agent. <i>Bioorganic and Medicinal Chemistry</i> , 2006 , 14, 6608-14	3.4	23
32	Structures of new amphidinols with truncated polyhydroxyl chain and their membrane-permeabilizing activities. <i>Bioorganic and Medicinal Chemistry</i> , 2006 , 14, 6548-54	3.4	68
31	Design and synthesis of an artificial ladder-shaped polyether that interacts with glycophorin A. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2006 , 16, 6355-9	2.9	21
30	Synthesis of 28-19F-amphotericin B methyl ester. <i>Tetrahedron Letters</i> , 2006 , 47, 6187-6191	2	34
29	Detailed description of the conformation and location of membrane-bound erythromycin a using isotropic bicelles. <i>Journal of Medicinal Chemistry</i> , 2006 , 49, 3501-8	8.3	19
28	Derivatization and Isotope Labeling of Amphotericin B Aiming at Elucidation of the Ion-channel Structure. <i>Yuki Gosei Kagaku Kyokaishi/Journal of Synthetic Organic Chemistry</i> , 2006 , 64, 502-514	0.2	2
27	Dominant formation of a single-length channel by amphotericin B in dimyristoylphosphatidylcholine membrane evidenced by ^{13}C - ^{31}P rotational echo double resonance. <i>Biochemistry</i> , 2005 , 44, 704-10	3.2	45
26	Mycosamine orientation of amphotericin B controlling interaction with ergosterol: sterol-dependent activity of conformation-restricted derivatives with an amino-carbonyl bridge. <i>Journal of the American Chemical Society</i> , 2005 , 127, 10667-75	16.4	69
25	Ladder-shaped polyether compound, desulfated yessotoxin, interacts with membrane-integral alpha-helix peptides. <i>Bioorganic and Medicinal Chemistry</i> , 2005 , 13, 5099-103	3.4	26
24	Hairpin conformation of amphidinols possibly accounting for potent membrane permeabilizing activities. <i>Tetrahedron</i> , 2005 , 61, 2795-2802	2.4	56
23	Isolation and structure elucidation of a new amphidinol with a truncated polyhydroxyl chain from <i>Amphidinium klebsii</i> . <i>Tetrahedron</i> , 2005 , 61, 8606-8610	2.4	71
22	Bioactive fluorinated derivative of amphotericin B. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2005 , 15, 3565-7	2.9	22
21	Amphotericin B covalent dimers bearing a tartarate linkage. <i>Chemistry and Biodiversity</i> , 2004 , 1, 346-52	2.5	15
20	Synthesis and conformation of deuterated spermidine for investigating weak interaction with polyanionic biomolecules. <i>Tetrahedron</i> , 2004 , 60, 5163-5170	2.4	4
19	An amphotericin B-ergosterol covalent conjugate with powerful membrane permeabilizing activity. <i>Chemistry and Biology</i> , 2004 , 11, 673-9		34
18	Membrane-permeabilizing activities of amphidinol 3, polyene-polyhydroxy antifungal from a marine dinoflagellate. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2004 , 1667, 91-100	3.8	46
17	Amphotericin B-phospholipid covalent conjugates: dependence of membrane-permeabilizing activity on acyl-chain length. <i>Organic and Biomolecular Chemistry</i> , 2003 , 1, 3882-4	3.9	24
16	Amphotericin B dimers with bisamide linkage bearing powerful membrane-permeabilizing activity. <i>Organic Letters</i> , 2002 , 4, 2087-9	6.2	30

15	Amphotericin B covalent dimers forming sterol-dependent ion-permeable membrane channels. <i>Journal of the American Chemical Society</i> , 2002 , 124, 4180-1	16.4	52
14	Absolute configuration of aflastatin A, a specific inhibitor of aflatoxin production by <i>Aspergillus parasiticus</i> . <i>Journal of Organic Chemistry</i> , 2000 , 65, 438-44	4.2	44
13	Leptomycin B inactivates CRM1/exportin 1 by covalent modification at a cysteine residue in the central conserved region. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999 , 96, 9112-7	11.5	858
12	Stereochemical Determination of Acyclic Structures Based on Carbon-Proton Spin-Coupling Constants. A Method of Configuration Analysis for Natural Products. <i>Journal of Organic Chemistry</i> , 1999 , 64, 866-876	4.2	610
11	Absolute Configuration of Amphidinol 3, the First Complete Structure Determination from Amphidinol Homologues: Application of a New Configuration Analysis Based on Carbon-Hydrogen Spin-Coupling Constants. <i>Journal of the American Chemical Society</i> , 1999 , 121, 870-871	16.4	169
10	Absolute Structure and Total Synthesis of Lipogrammistin-A, a Lipophilic Ichthyotoxin of the Soapfish. <i>Journal of Organic Chemistry</i> , 1998 , 63, 3925-3932	4.2	13
9	Involvement of afsA in A-factor biosynthesis as a key enzyme. <i>Journal of Antibiotics</i> , 1997 , 50, 847-52	3.7	38
8	Die Struktur von Maitotoxin II: Konfiguration der C1-C14-Seitenkette. <i>Angewandte Chemie</i> , 1996 , 108, 1782-1785	3.6	26
7	Die Struktur von Maitotoxin III: Konfiguration der C135-C142-Seitenkette und absolute Konfiguration des gesamten Moleküls. <i>Angewandte Chemie</i> , 1996 , 108, 1786-1789	3.6	22
6	The Complete Structure of Maitotoxin, Part I: Configuration of the C1-C14 Side Chain. <i>Angewandte Chemie International Edition in English</i> , 1996 , 35, 1672-1675		93
5	The Complete Structure of Maitotoxin, Part II: Configuration of the C135-C142 Side Chain and Absolute Configuration of the Entire Molecule. <i>Angewandte Chemie International Edition in English</i> , 1996 , 35, 1675-1678		91
4	Long-range carbon-proton coupling constants for stereochemical assignment of acyclic structures in natural products: Configuration of the C5-C9 portion of maitotoxin. <i>Tetrahedron Letters</i> , 1996 , 37, 1269-1272	2	46
3	Isolation and chemical structure of amphidinol 2, a potent hemolytic compound from marine dinoflagellate <i>Amphidinium klebsii</i> . <i>Tetrahedron Letters</i> , 1995 , 36, 6279-6282	2	103
2	Stereochemical assignment of the C35-C39 Acyclic linkage in maitotoxin: completion of stereochemical determination of C15-C134. <i>Tetrahedron Letters</i> , 1995 , 36, 9011-9014	2	44
1	Conformational analysis of natural products using long-range carbon-proton coupling constants: Three-dimensional structure of okadaic acid in solution. <i>Tetrahedron</i> , 1995 , 51, 12229-12238	2.4	54