

Jiadi Luo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7420095/publications.pdf>

Version: 2024-02-01

29
papers

791
citations

623734

14
h-index

501196

28
g-index

30
all docs

30
docs citations

30
times ranked

1385
citing authors

#	ARTICLE	IF	CITATIONS
1	The roles of tumor-derived exosomes in non-small cell lung cancer and their clinical implications. <i>Journal of Experimental and Clinical Cancer Research</i> , 2018, 37, 226.	8.6	107
2	AIM2 Inflammasome Is Critical for Influenza-Induced Lung Injury and Mortality. <i>Journal of Immunology</i> , 2017, 198, 4383-4393.	0.8	85
3	MicroRNAs associated with therapy of non-small cell lung cancer. <i>International Journal of Biological Sciences</i> , 2018, 14, 390-397.	6.4	78
4	Genome-wide identification of interferon-sensitive mutations enables influenza vaccine design. <i>Science</i> , 2018, 359, 290-296.	12.6	64
5	Activation of Akt/mTOR Pathway Is Associated with Poor Prognosis of Nasopharyngeal Carcinoma. <i>PLoS ONE</i> , 2014, 9, e106098.	2.5	57
6	Suppression Of β -catenin Nuclear Translocation By CGP57380 Decelerates Poor Progression And Potentiates Radiation-Induced Apoptosis in Nasopharyngeal Carcinoma. <i>Theranostics</i> , 2017, 7, 2134-2149.	10.0	51
7	CGP57380 enhances efficacy of RAD001 in non-small cell lung cancer through abrogating mTOR inhibition-induced phosphorylation of eIF4E and activating mitochondrial apoptotic pathway. <i>Oncotarget</i> , 2016, 7, 27787-27801.	1.8	47
8	Roles of exosomes in the carcinogenesis and clinical therapy of non-small cell lung cancer. <i>Biomedicine and Pharmacotherapy</i> , 2019, 111, 338-346.	5.6	44
9	Phosphorylated Mnk1 and eIF4E Are Associated with Lymph Node Metastasis and Poor Prognosis of Nasopharyngeal Carcinoma. <i>PLoS ONE</i> , 2014, 9, e89220.	2.5	43
10	Flot-2 Expression Correlates with EGFR Levels and Poor Prognosis in Surgically Resected Non-Small Cell Lung Cancer. <i>PLoS ONE</i> , 2015, 10, e0132190.	2.5	28
11	Increased Expression of Flotillin-2 Protein as a Novel Biomarker for Lymph Node Metastasis in Nasopharyngeal Carcinoma. <i>PLoS ONE</i> , 2014, 9, e101676.	2.5	21
12	Elevated expression of HSP10 protein inhibits apoptosis and associates with poor prognosis of astrocytoma. <i>PLoS ONE</i> , 2017, 12, e0185563.	2.5	21
13	Exosome-mediated miR-7-5p delivery enhances the anticancer effect of Everolimus via blocking MNK/eIF4E axis in non-small cell lung cancer. <i>Cell Death and Disease</i> , 2022, 13, 129.	6.3	21
14	Elevated expression of mcl-1 inhibits apoptosis and predicts poor prognosis in patients with surgically resected non-small cell lung cancer. <i>Diagnostic Pathology</i> , 2019, 14, 108.	2.0	16
15	Increased expression of IRS-1 is associated with lymph node metastasis in nasopharyngeal carcinoma. <i>International Journal of Clinical and Experimental Pathology</i> , 2014, 7, 6117-24.	0.5	15
16	Increased expression of Cks1 protein is associated with lymph node metastasis and poor prognosis in nasopharyngeal carcinoma. <i>Diagnostic Pathology</i> , 2017, 12, 2.	2.0	13
17	Epigenetic Regulation of IL-17-Induced Chemokines in Lung Epithelial Cells. <i>Mediators of Inflammation</i> , 2019, 2019, 1-11.	3.0	13
18	4EGI-1 induces apoptosis and enhances radiotherapy sensitivity in nasopharyngeal carcinoma cells via DR5 induction on 4E-BP1 dephosphorylation. <i>Oncotarget</i> , 2016, 7, 21728-21741.	1.8	12

#	ARTICLE	IF	CITATIONS
19	Fatty acid synthase-associated protein with death domain: a prognostic factor for survival in patients with nasopharyngeal carcinoma. <i>Human Pathology</i> , 2014, 45, 2447-2452.	2.0	10
20	Comprehensive analysis of the mechanism and treatment significance of Mucins in lung cancer. <i>Journal of Experimental and Clinical Cancer Research</i> , 2020, 39, 162.	8.6	10
21	FLOT-2 is an independent prognostic marker in oral squamous cell carcinoma. <i>International Journal of Clinical and Experimental Pathology</i> , 2015, 8, 8236-43.	0.5	8
22	High expression of heat shock protein 10 correlates negatively with estrogen/progesterone receptor status and predicts poor prognosis in invasive ductal breast carcinoma. <i>Human Pathology</i> , 2017, 61, 173-180.	2.0	7
23	Elevated expression of IRS-1 associates with phosphorylated Akt expression and predicts poor prognosis of breast invasive ductal carcinoma. <i>Human Pathology</i> , 2018, 79, 9-17.	2.0	7
24	Lipoadenoma of the adrenal gland: report of a rare entity and review of literature. <i>International Journal of Clinical and Experimental Pathology</i> , 2015, 8, 9693-7.	0.5	5
25	Expression of cancer cell-intrinsic PD-1 associates with PD-L1 and p-S6 and predicts a good prognosis in nasopharyngeal carcinoma. <i>Journal of Cancer</i> , 2021, 12, 6118-6125.	2.5	4
26	Primary synovial sarcoma of the larynx in a 14-year-old boy diagnosed by immunohistochemistry and fluorescence in situ hybridization: a rare case report. <i>International Journal of Clinical and Experimental Pathology</i> , 2017, 10, 7913-7919.	0.5	2
27	Primary cutaneous $\hat{I}^3\hat{I}$ -T-cell lymphoma (CGD-TCL) with unilateral lower extremity swelling as first-onset symptom: a rare case report. <i>International Journal of Clinical and Experimental Pathology</i> , 2014, 7, 5337-42.	0.5	1
28	High expression of heat shock protein 10 (Hsp10) is associated with poor prognosis in oral squamous cell carcinoma. <i>International Journal of Clinical and Experimental Pathology</i> , 2017, 10, 7784-7791.	0.5	1
29	Insight into the regulatory mechanism of VEGFA isoforms in p53 mutant breast cancer: can it be substantiated with solely in vitro studies?. <i>Journal of Thoracic Disease</i> , 2018, 10, 567-569.	1.4	0