
Jiadi Luo

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7420095/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Exosome-mediated miR-7-5p delivery enhances the anticancer effect of Everolimus via blocking MNK/eIF4E axis in non-small cell lung cancer. Cell Death and Disease, 2022, 13, 129.	6.3	21
2	Expression of cancer cell-intrinsic PD-1 associates with PD-L1 and p-S6 and predicts a good prognosis in nasopharyngeal carcinoma. Journal of Cancer, 2021, 12, 6118-6125.	2.5	4
3	Comprehensive analysis of the mechanism and treatment significance of Mucins in lung cancer. Journal of Experimental and Clinical Cancer Research, 2020, 39, 162.	8.6	10
4	Elevated expression of mcl-1 inhibits apoptosis and predicts poor prognosis in patients with surgically resected non-small cell lung cancer. Diagnostic Pathology, 2019, 14, 108.	2.0	16
5	Epigenetic Regulation of IL-17-Induced Chemokines in Lung Epithelial Cells. Mediators of Inflammation, 2019, 2019, 1-11.	3.0	13
6	Roles of exosomes in the carcinogenesis and clinical therapy of non-small cell lung cancer. Biomedicine and Pharmacotherapy, 2019, 111, 338-346.	5.6	44
7	Elevated expression of IRS-1 associates with phosphorylated Akt expression and predicts poor prognosis of breast invasive ductal carcinoma. Human Pathology, 2018, 79, 9-17.	2.0	7
8	Genome-wide identification of interferon-sensitive mutations enables influenza vaccine design. Science, 2018, 359, 290-296.	12.6	64
9	Insight into the regulatory mechanism of VEGFA isoforms in p53 mutant breast cancer: can it be substantiated with solely in vitro studies?. Journal of Thoracic Disease, 2018, 10, 567-569.	1.4	0
10	The roles of tumor-derived exosomes in non-small cell lung cancer and their clinical implications. Journal of Experimental and Clinical Cancer Research, 2018, 37, 226.	8.6	107
11	MicroRNAs associated with therapy of non-small cell lung cancer. International Journal of Biological Sciences, 2018, 14, 390-397.	6.4	78
12	Increased expression of Cks1 protein is associated with lymph node metastasis and poor prognosis in nasopharyngeal carcinoma. Diagnostic Pathology, 2017, 12, 2.	2.0	13
13	AIM2 Inflammasome Is Critical for Influenza-Induced Lung Injury and Mortality. Journal of Immunology, 2017, 198, 4383-4393.	0.8	85
14	High expression of heat shock protein 10 correlates negatively with estrogen/progesterone receptor status and predicts poor prognosis in invasive ductal breast carcinoma. Human Pathology, 2017, 61, 173-180.	2.0	7
15	Suppression Of β-catenin Nuclear Translocation By CGP57380 Decelerates Poor Progression And Potentiates Radiation-Induced Apoptosis in Nasopharyngeal Carcinoma. Theranostics, 2017, 7, 2134-2149.	10.0	51
16	Elevated expression of HSP10 protein inhibits apoptosis and associates with poor prognosis of astrocytoma. PLoS ONE, 2017, 12, e0185563.	2.5	21
17	Primary synovial sarcoma of the larynx in a 14-year-old boy diagnosed by immunohistochemistry and fluorescence in situ hybridization: a rare case report. International Journal of Clinical and Experimental Pathology, 2017, 10, 7913-7919.	0.5	2
18	High expression of heat shock protein 10 (Hsp10) is associated with poor prognosis in oral squamous cell carcinoma. International Journal of Clinical and Experimental Pathology, 2017, 10, 7784-7791.	0.5	1

Jiadi Luo

#	Article	IF	CITATIONS
19	4EGI-1 induces apoptosis and enhances radiotherapy sensitivity in nasopharyngeal carcinoma cells via DR5 induction on 4E-BP1 dephosphorylation. Oncotarget, 2016, 7, 21728-21741.	1.8	12
20	CGP57380 enhances efficacy of RAD001 in non-small cell lung cancer through abrogating mTOR inhibition-induced phosphorylation of eIF4E and activating mitochondrial apoptotic pathway. Oncotarget, 2016, 7, 27787-27801.	1.8	47
21	Flot-2 Expression Correlates with EGFR Levels and Poor Prognosis in Surgically Resected Non-Small Cell Lung Cancer. PLoS ONE, 2015, 10, e0132190.	2.5	28
22	FLOT-2 is an independent prognostic marker in oral squamous cell carcinoma. International Journal of Clinical and Experimental Pathology, 2015, 8, 8236-43.	0.5	8
23	Lipoadenoma of the adrenal gland: report of a rare entity and review of literature. International Journal of Clinical and Experimental Pathology, 2015, 8, 9693-7.	0.5	5
24	Activation of Akt/mTOR Pathway Is Associated with Poor Prognosis of Nasopharyngeal Carcinoma. PLoS ONE, 2014, 9, e106098.	2.5	57
25	Fatty acid synthase–associated protein with death domain: a prognostic factor for survival in patients with nasopharyngeal carcinoma. Human Pathology, 2014, 45, 2447-2452.	2.0	10
26	Phosphorylated Mnk1 and eIF4E Are Associated with Lymph Node Metastasis and Poor Prognosis of Nasopharyngeal Carcinoma. PLoS ONE, 2014, 9, e89220.	2.5	43
27	Increased Expression of Flotillin-2 Protein as a Novel Biomarker for Lymph Node Metastasis in Nasopharyngeal Carcinoma. PLoS ONE, 2014, 9, e101676.	2.5	21
28	Primary cutaneous Î ³ Î-T-cell lymphoma (CGD-TCL) with unilateral lower extremity swelling as first-onset symptom: a rare case report. International Journal of Clinical and Experimental Pathology, 2014, 7, 5337-42.	0.5	1
29	Increased expression of IRS-1 is associated with lymph node metastasis in nasopharyngeal carcinoma. International Journal of Clinical and Experimental Pathology, 2014, 7, 6117-24.	0.5	15