

ValÃ©rie Schreiber

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7423756/publications.pdf>

Version: 2024-02-01

77
papers

11,748
citations

53751

45
h-index

64755

79
g-index

82
all docs

82
docs citations

82
times ranked

10460
citing authors

#	ARTICLE	IF	CITATIONS
1	Poly(ADP-ribose): novel functions for an old molecule. <i>Nature Reviews Molecular Cell Biology</i> , 2006, 7, 517-528.	16.1	1,719
2	XRCC1 Is Specifically Associated with Poly(ADP-Ribose) Polymerase and Negatively Regulates Its Activity following DNA Damage. <i>Molecular and Cellular Biology</i> , 1998, 18, 3563-3571.	1.1	843
3	PARP-1 Inhibition Increases Mitochondrial Metabolism through SIRT1 Activation. <i>Cell Metabolism</i> , 2011, 13, 461-468.	7.2	673
4	PARP-2, A Novel Mammalian DNA Damage-dependent Poly(ADP-ribose) Polymerase. <i>Journal of Biological Chemistry</i> , 1999, 274, 17860-17868.	1.6	644
5	Poly(ADP-ribose) Polymerase-2 (PARP-2) Is Required for Efficient Base Excision DNA Repair in Association with PARP-1 and XRCC1. <i>Journal of Biological Chemistry</i> , 2002, 277, 23028-23036.	1.6	602
6	Functional interaction between PARP-1 and PARP-2 in chromosome stability and embryonic development in mouse. <i>EMBO Journal</i> , 2003, 22, 2255-2263.	3.5	544
7	Base Excision Repair Is Impaired in Mammalian Cells Lacking Poly(ADP-ribose) Polymerase-1. <i>Biochemistry</i> , 2000, 39, 7559-7569.	1.2	440
8	The diverse roles and clinical relevance of PARPs in DNA damage repair: Current state of the art. <i>Biochemical Pharmacology</i> , 2012, 84, 137-146.	2.0	428
9	Involvement of poly(ADP-ribose) polymerase in base excision repair. <i>Biochimie</i> , 1999, 81, 69-75.	1.3	317
10	Feedback-regulated poly(ADP-ribosyl)ation by PARP-1 is required for rapid response to DNA damage in living cells. <i>Nucleic Acids Research</i> , 2007, 35, 7665-7675.	6.5	271
11	Poly(ADP-ribose) polymerases in double-strand break repair: Focus on PARP1, PARP2 and PARP3. <i>Experimental Cell Research</i> , 2014, 329, 18-25.	1.2	238
12	Parp-1 protects homologous recombination from interference by Ku and Ligase IV in vertebrate cells. <i>EMBO Journal</i> , 2006, 25, 1305-1314.	3.5	237
13	Poly(ADP-ribose) polymerase 3 (PARP3), a newcomer in cellular response to DNA damage and mitotic progression. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 2783-2788.	3.3	235
14	PARP-2 Regulates SIRT1 Expression and Whole-Body Energy Expenditure. <i>Cell Metabolism</i> , 2011, 13, 450-460.	7.2	231
15	A Nuclear Poly(ADP-Ribose)-Dependent Signalosome Confers DNA Damage-Induced β Kinase Activation. <i>Molecular Cell</i> , 2009, 36, 365-378.	4.5	216
16	Structure and function of poly(ADP-ribose) polymerase. <i>Molecular and Cellular Biochemistry</i> , 1994, 138, 15-24.	1.4	203
17	A dominant-negative mutant of human poly(ADP-ribose) polymerase affects cell recovery, apoptosis, and sister chromatid exchange following DNA damage. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 4753-4757.	3.3	203
18	Functional Interaction between Poly(ADP-Ribose) Polymerase 2 (PARP-2) and TRF2: PARP Activity Negatively Regulates TRF2. <i>Molecular and Cellular Biology</i> , 2004, 24, 1595-1607.	1.1	166

#	ARTICLE	IF	CITATIONS
19	PARP1â€™TDP1 coupling for the repair of topoisomerase â€™induced DNA damage. <i>Nucleic Acids Research</i> , 2014, 42, 4435-4449.	6.5	163
20	PARP-1 and PARP-2 interact with nucleophosmin/B23 and accumulate in transcriptionally active nucleoli. <i>Journal of Cell Science</i> , 2005, 118, 211-222.	1.2	156
21	Poly(ADPâ€™ribose) Polymeraseâ€™1 Activation During DNA Damage and Repair. <i>Methods in Enzymology</i> , 2006, 409, 493-510.	0.4	150
22	Toward specific functions of poly(ADP-ribose) polymerase-2. <i>Trends in Molecular Medicine</i> , 2008, 14, 169-178.	3.5	142
23	The expanding field of poly(ADPâ€™ribosyl)ation reactions. <i>EMBO Reports</i> , 2008, 9, 1094-1100.	2.0	140
24	Laspâ€™1 (MLN 50) defines a new LIM protein subfamily characterized by the association of LIM and SH3 domains. <i>FEBS Letters</i> , 1995, 373, 245-249.	1.3	132
25	Radiation-induced mitotic catastrophe in PARG-deficient cells. <i>Journal of Cell Science</i> , 2009, 122, 1990-2002.	1.2	114
26	PARP-2 deficiency affects the survival of CD4+CD8+ double-positive thymocytes. <i>EMBO Journal</i> , 2006, 25, 4350-4360.	3.5	112
27	PARG is recruited to DNA damage sites through poly(ADP-ribose)- and PCNA-dependent mechanisms. <i>Nucleic Acids Research</i> , 2011, 39, 5045-5056.	6.5	108
28	PARP1 Is a TRF2-associated Poly(ADP-Ribose)Polymerase and Protects Eroded Telomeres. <i>Molecular Biology of the Cell</i> , 2006, 17, 1686-1696.	0.9	106
29	PARP-1 transcriptional activity is regulated by sumoylation upon heat shock. <i>EMBO Journal</i> , 2009, 28, 3534-3548.	3.5	103
30	Poly(ADP-ribose) polymerase: Molecular biological aspects. <i>BioEssays</i> , 1991, 13, 455-462.	1.2	99
31	Peroxisome Proliferator-activated Receptor (PPAR)-2 Controls Adipocyte Differentiation and Adipose Tissue Function through the Regulation of the Activity of the Retinoid X Receptor/PPARÎ³ Heterodimer. <i>Journal of Biological Chemistry</i> , 2007, 282, 37738-37746.	1.6	97
32	The role of poly(ADP-ribosyl)ation in epigenetic events. <i>International Journal of Biochemistry and Cell Biology</i> , 2009, 41, 60-65.	1.2	96
33	Parp-2 is required to maintain hematopoiesis following sublethal Î³-irradiation in mice. <i>Blood</i> , 2013, 122, 44-54.	0.6	96
34	PARP-2 sustains erythropoiesis in mice by limiting replicative stress in erythroid progenitors. <i>Cell Death and Differentiation</i> , 2015, 22, 1144-1157.	5.0	95
35	Poly(ADP-ribose) polymerase 1 regulates both the exonuclease and helicase activities of the Werner syndrome protein. <i>Nucleic Acids Research</i> , 2004, 32, 4003-4014.	6.5	89
36	Lasp-1, a Novel Type of Actin-Binding Protein Accumulating in Cell Membrane Extensions. <i>Molecular Medicine</i> , 1998, 4, 675-687.	1.9	86

#	ARTICLE	IF	CITATIONS
37	PARP3 affects the relative contribution of homologous recombination and nonhomologous end-joining pathways. <i>Nucleic Acids Research</i> , 2014, 42, 5616-5632.	6.5	82
38	New readers and interpretations of poly(ADP-ribosyl)ation. <i>Trends in Biochemical Sciences</i> , 2012, 37, 381-390.	3.7	75
39	Poly(ADP-ribose) Polymerase-1 (PARP-1) Is Required in Murine Cell Lines for Base Excision Repair of Oxidative DNA Damage in the Absence of DNA Polymerase β . <i>Journal of Biological Chemistry</i> , 2003, 278, 18471-18477.	1.6	71
40	PARP3 controls TGF β ² and ROS driven epithelial-to-mesenchymal transition and stemness by stimulating a TG2-Snail-E-cadherin axis. <i>Oncotarget</i> , 2016, 7, 64109-64123.	0.8	71
41	Expanding functions of ADP-ribosylation in the maintenance of genome integrity. <i>Seminars in Cell and Developmental Biology</i> , 2017, 63, 92-101.	2.3	69
42	Regulation of NFAT by poly(ADP-ribose) polymerase activity in T cells. <i>Molecular Immunology</i> , 2008, 45, 1863-1871.	1.0	68
43	PARC is dispensable for recovery from transient replicative stress but required to prevent detrimental accumulation of poly(ADP-ribose) upon prolonged replicative stress. <i>Nucleic Acids Research</i> , 2014, 42, 7776-7792.	6.5	58
44	Functional interplay between Parp-1 and SirT1 in genome integrity and chromatin-based processes. <i>Cellular and Molecular Life Sciences</i> , 2009, 66, 3219-3234.	2.4	53
45	Interaction of PARP-2 with DNA structures mimicking DNA repair intermediates and consequences on activity of base excision repair proteins. <i>Biochimie</i> , 2013, 95, 1208-1215.	1.3	52
46	PARP-1/PARP-2 double deficiency in mouse T cells results in faulty immune responses and T lymphomas. <i>Scientific Reports</i> , 2017, 7, 41962.	1.6	51
47	PARP-2 Interacts with TTF-1 and Regulates Expression of Surfactant Protein-B. <i>Journal of Biological Chemistry</i> , 2006, 281, 9600-9606.	1.6	48
48	Poly (ADP-Ribose) Glycohydrolase Regulates Retinoic Acid Receptor-Mediated Gene Expression. <i>Molecular Cell</i> , 2012, 48, 785-798.	4.5	48
49	Kin17, a mouse nuclear zinc finger protein that binds preferentially to curved DNA. <i>Nucleic Acids Research</i> , 1994, 22, 4335-4341.	6.5	44
50	Autophagy requires poly(adp-ribosyl)ation-dependent AMPK nuclear export. <i>Cell Death and Differentiation</i> , 2016, 23, 2007-2018.	5.0	44
51	A Bidirectional Promoter Connects the Poly(ADP-ribose) Polymerase 2 (PARP-2) Gene to the Gene for RNase P RNA. <i>Journal of Biological Chemistry</i> , 2001, 276, 11092-11099.	1.6	43
52	Nucleolar localization of aprataxin is dependent on interaction with nucleolin and on active ribosomal DNA transcription. <i>Human Molecular Genetics</i> , 2006, 15, 2239-2249.	1.4	40
53	Poly(ADP-ribose) Polymerase 1 (PARP1) Associates with E3 Ubiquitin-Protein Ligase UHRF1 and Modulates UHRF1 Biological Functions. <i>Journal of Biological Chemistry</i> , 2014, 289, 16223-16238.	1.6	39
54	Rfx6 promotes the differentiation of peptide-secreting enteroendocrine cells while repressing genetic programs controlling serotonin production. <i>Molecular Metabolism</i> , 2019, 29, 24-39.	3.0	39

#	ARTICLE	IF	CITATIONS
55	The expanding field of poly(ADP-ribose)ation reactions. <i>Protein Modifications: Beyond the Usual Suspects</i> ™ Review Series. <i>EMBO Reports</i> , 2008, 9, 1252-1252.	2.0	35
56	Chromosomal assignment and expression pattern of the murine Lasp-1 gene. <i>Gene</i> , 1998, 207, 171-175.	1.0	31
57	XRCC1 interacts with the p58 subunit of DNA Pol δ -primase and may coordinate DNA repair and replication during S phase. <i>Nucleic Acids Research</i> , 2009, 37, 3177-3188.	6.5	28
58	Functional aspects of PARylation in induced and programmed DNA repair processes: Preserving genome integrity and modulating physiological events. <i>Molecular Aspects of Medicine</i> , 2013, 34, 1138-1152.	2.7	28
59	Poly(ADP-ribose)ation of Methyl CpG Binding Domain Protein 2 Regulates Chromatin Structure. <i>Journal of Biological Chemistry</i> , 2016, 291, 4873-4881.	1.6	28
60	The macroPARP genes <i>parp9</i> and <i>parp14</i> are developmentally and differentially regulated in mouse tissues. <i>Developmental Dynamics</i> , 2008, 237, 209-215.	0.8	25
61	Activation of the abundant nuclear factor poly(ADP-ribose) polymerase-1 by <i>Helicobacter pylori</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 19998-20003.	3.3	25
62	Genetic Ablation of PARP-1 Protects Against Oxazolone-Induced Contact Hypersensitivity by Modulating Oxidative Stress. <i>Journal of Investigative Dermatology</i> , 2010, 130, 2629-2637.	0.3	23
63	PARP3, a new therapeutic target to alter Rictor/mTORC2 signaling and tumor progression in BRCA1-associated cancers. <i>Cell Death and Differentiation</i> , 2019, 26, 1615-1630.	5.0	23
64	Phenotypic Characterization of Parp-1 and Parp-2 Deficient Mice and Cells. <i>Methods in Molecular Biology</i> , 2011, 780, 313-336.	0.4	23
65	Functional interaction between human papillomavirus type 18 E2 and poly(ADP-ribose) polymerase 1. <i>Oncogene</i> , 2002, 21, 5877-5885.	2.6	22
66	Purification of Recombinant Poly(ADP-Ribose) Polymerases. <i>Methods in Molecular Biology</i> , 2011, 780, 135-152.	0.4	22
67	PARG deficiency is neither synthetic lethal with BRCA1 nor PTEN deficiency. <i>Cancer Cell International</i> , 2016, 16, 53.	1.8	20
68	Extensive NEUROG3 occupancy in the human pancreatic endocrine gene regulatory network. <i>Molecular Metabolism</i> , 2021, 53, 101313.	3.0	20
69	Poly(ADP-ribose) polymerase: Structure-function relationship. <i>Biochimie</i> , 1995, 77, 456-461.	1.3	19
70	Parp2 is required for the differentiation of post-meiotic germ cells: Identification of a spermatid-specific complex containing Parp1, Parp2, TP2 and HSPA2. <i>Experimental Cell Research</i> , 2009, 315, 2824-2834.	1.2	19
71	Detection of the Nuclear Poly(ADP-ribose)-Metabolizing Enzymes and Activities in Response to DNA Damage. <i>Methods in Molecular Biology</i> , 2008, 464, 267-283.	0.4	15
72	A eukaryotic expression vector for the study of nuclear localization signals. <i>Gene</i> , 1994, 150, 411-412.	1.0	10

#	ARTICLE	IF	CITATIONS
73	Robust immunoglobulin class switch recombination and end joining in <i>Parp9</i> deficient mice. <i>European Journal of Immunology</i> , 2017, 47, 665-676.	1.6	8
74	PARP-2: Structure-Function Relationship. , 2006, , 13-31.		6
75	Purification of Recombinant Human PARC and Activity Assays. <i>Methods in Molecular Biology</i> , 2017, 1608, 395-413.	0.4	6
76	Discovery of the PARP Superfamily and Focus on the Lesser Exhibited But Not Lesser Talented Members. <i>Cancer Drug Discovery and Development</i> , 2015, , 15-46.	0.2	3
77	Purification of Recombinant Human PARP-3. <i>Methods in Molecular Biology</i> , 2017, 1608, 373-394.	0.4	1