

Wolfgang Otto

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7431320/publications.pdf>

Version: 2024-02-01

53
papers

1,226
citations

430442

18
h-index

414034

32
g-index

53
all docs

53
docs citations

53
times ranked

1781
citing authors

#	ARTICLE	IF	CITATIONS
1	The Tumor Immune Microenvironment Drives a Prognostic Relevance That Correlates with Bladder Cancer Subtypes. <i>Cancer Immunology Research</i> , 2019, 7, 923-938.	1.6	148
2	The WHO classification of 1973 is more suitable than the WHO classification of 2004 for predicting survival in pT1 urothelial bladder cancer. <i>BJU International</i> , 2011, 107, 404-408.	1.3	86
3	Performance of the Food and Drug Administration/EMA-approved programmed cell death ligand-1 assays in urothelial carcinoma with emphasis on therapy stratification for first-line use of atezolizumab and pembrolizumab. <i>European Journal of Cancer</i> , 2019, 106, 234-243.	1.3	75
4	Analysis of Sex Differences in Cancer-Specific Survival and Perioperative Mortality Following Radical Cystectomy: Results of a Large German Multicenter Study of Nearly 2500 Patients with Urothelial Carcinoma of the Bladder. <i>Gender Medicine</i> , 2012, 9, 481-489.	1.4	65
5	In stage pT1 non-muscle-invasive bladder cancer (NMIBC), high KRT20 and low KRT5 mRNA expression identify the luminal subtype and predict recurrence and survival. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2017, 470, 267-274.	1.4	58
6	FGFR3 Mutation Status and FGFR3 Expression in a Large Bladder Cancer Cohort Treated by Radical Cystectomy: Implications for Anti-FGFR3 Treatment?â€. <i>European Urology</i> , 2020, 78, 682-687.	0.9	57
7	High PDL1 mRNA expression predicts better survival of stage pT1 non-muscle-invasive bladder cancer (NMIBC) patients. <i>Cancer Immunology, Immunotherapy</i> , 2018, 67, 403-412.	2.0	54
8	Cytotoxic T-cell-related gene expression signature predicts improved survival in muscle-invasive urothelial bladder cancer patients after radical cystectomy and adjuvant chemotherapy. , 2020, 8, e000162.		45
9	Biochemical Foundations of Health and Energy Conservation in Hibernating Free-ranging Subadult Brown Bear <i>Ursus arctos</i> . <i>Journal of Biological Chemistry</i> , 2016, 291, 22509-22523.	1.6	37
10	Epithelialâ€mesenchymal transformation markers E-cadherin and survivin predict progression of stage pT1a urothelial bladder carcinoma. <i>World Journal of Urology</i> , 2016, 34, 709-716.	1.2	36
11	mRNA-Expression of KRT5 and KRT20 Defines Distinct Prognostic Subgroups of Muscle-Invasive Urothelial Bladder Cancer Correlating with Histological Variants. <i>International Journal of Molecular Sciences</i> , 2018, 19, 3396.	1.8	35
12	A multicenter round robin test of PD-L1 expression assessment in urothelial bladder cancer by immunohistochemistry and RT-qPCR with emphasis on prognosis prediction after radical cystectomy. <i>Oncotarget</i> , 2018, 9, 15001-15014.	0.8	33
13	CDKN2A as transcriptomic marker for muscle-invasive bladder cancer risk stratification and therapy decision-making. <i>Scientific Reports</i> , 2018, 8, 14383.	1.6	32
14	ESR1, ERBB2, and Ki67 mRNA expression predicts stage and grade of non-muscle-invasive bladder carcinoma (NMIBC). <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2016, 469, 547-552.	1.4	25
15	The World Health Organization 1973 classification system for grade is an important prognosticator in T1 nonâ€muscleâ€invasive bladder cancer. <i>BJU International</i> , 2018, 122, 978-985.	1.3	25
16	High proliferation rate and TNM stage but not histomorphological subtype are independent prognostic markers for overall survival in papillary renal cell carcinoma. <i>Human Pathology</i> , 2019, 83, 212-223.	1.1	23
17	High Androgen Receptor mRNA Expression Is Independently Associated with Prolonged Cancer-Specific and Recurrence-Free Survival in Stage T1 Bladder Cancer. <i>Translational Oncology</i> , 2017, 10, 340-345.	1.7	22
18	Prognostic markers in invasive bladder cancer: FGFR3 mutation status versus P53 and KI-67 expression: a multi-center, multi-laboratory analysis in 1058 radical cystectomy patients. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2022, 40, 110.e1-110.e9.	0.8	22

#	ARTICLE	IF	CITATIONS
19	No mutations of FGFR3 in normal urothelium in the vicinity of urothelial carcinoma of the bladder harbouring activating FGFR3 mutations in patients with bladder cancer. <i>International Journal of Cancer</i> , 2009, 125, 2205-2208.	2.3	20
20	Metric substage according to micro and extensive lamina propria invasion improves prognostics in T1 bladder cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2018, 36, 361.e7-361.e13.	0.8	20
21	<sc>FOXM</sc>1 overexpression is associated with adverse outcome and predicts response to intravesical instillation therapy in stage <sc>pT</sc>1 nonâ€muscleâ€invasive bladder cancer. <i>BJU International</i> , 2019, 123, 187-196.	1.3	19
22	Comparative analysis of comorbidity and performance indices for prediction of oncological outcomes in patients with upper tract urothelial carcinoma who were treated with radical nephroureterectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014, 32, 1141-1150.	0.8	18
23	Infiltrative lamina propria invasion pattern as an independent predictor for cancerâ€specific and overall survival of instillation treatmentâ€naÃve stageÂT1 highâ€grade urothelial bladder cancer. <i>International Journal of Urology</i> , 2018, 25, 442-449.	0.5	17
24	Differentiationâ€associated urothelial cytochrome P450 oxidoreductase predicates the xenobioticâ€metabolizing activity of â€luminalâ€muscleâ€invasive bladder cancers. <i>Molecular Carcinogenesis</i> , 2018, 57, 606-618.	1.3	17
25	Expression, localisation and potential significance of aquaporins in benign and malignant human prostate tissue. <i>BMC Urology</i> , 2018, 18, 75.	0.6	17
26	High CDKN2A/p16 and Low FGFR3 Expression Predict Progressive Potential of Stage pT1 Urothelial Bladder Carcinoma. <i>Clinical Genitourinary Cancer</i> , 2018, 16, 248-256.e2.	0.9	16
27	Androgen Receptor mRNA Expression in Urothelial Carcinoma of the Bladder: A Retrospective Analysis of Two Independent Cohorts. <i>Translational Oncology</i> , 2019, 12, 661-668.	1.7	16
28	FOXM1 predicts overall and disease specific survival in muscle-invasive urothelial carcinoma and presents a differential expression between bladder cancer subtypes. <i>Oncotarget</i> , 2017, 8, 47595-47606.	0.8	16
29	Predictive value of molecular subtyping in NMIBC by RT-qPCR of ERBB2, ESR1, PGR and MKI67 from formalin fixed TUR biopsies. <i>Oncotarget</i> , 2017, 8, 67684-67695.	0.8	16
30	Photodynamic Diagnosis for Superficial Bladder Cancer: Do All Risk-Groups Profit Equally from Oncological and Economic Long-Term Results?. <i>Clinical Medicine Oncology</i> , 2009, 3, CMO.S1012.	0.2	14
31	High Stroma T-Cell Infiltration is Associated with Better Survival in Stage pT1 Bladder Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8407.	1.8	14
32	FOXA1 Gene Expression for Defining Molecular Subtypes of Muscle-Invasive Bladder Cancer after Radical Cystectomy. <i>Journal of Clinical Medicine</i> , 2020, 9, 994.	1.0	14
33	Loss of AQP3 protein expression is associated with worse progression-free and cancer-specific survival in patients with muscle-invasive bladder cancer. <i>World Journal of Urology</i> , 2015, 33, 1959-1964.	1.2	12
34	WHO 1973 grade 3 and infiltrative growth pattern proved, aberrant E-cadherin expression tends to be of predictive value for progression in a series of stage T1 high-grade bladder cancer after organ-sparing approach. <i>International Urology and Nephrology</i> , 2017, 49, 431-437.	0.6	11
35	A switch from epithelial to mesenchymal properties correlates with lymphovascular invasion in squamous cell carcinoma of the penis. <i>Pathology Research and Practice</i> , 2015, 211, 641-645.	1.0	10
36	Analysis of the prognostic relevance of sex-steroid hormonal receptor mRNA expression in muscle-invasive urothelial carcinoma of the urinary bladder. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2019, 474, 209-217.	1.4	10

#	ARTICLE	IF	CITATIONS
37	The Prognostic Value of FGFR3 Expression in Patients with T1 Non-Muscle Invasive Bladder Cancer. <i>Cancer Management and Research</i> , 2021, Volume 13, 6567-6578.	0.9	10
38	Tumor budding correlates with tumor invasiveness and predicts worse survival in pT1 non-muscle-invasive bladder cancer. <i>Scientific Reports</i> , 2021, 11, 17981.	1.6	9
39	Increased Proliferation as Independent Predictor of Disease Recurrence in Initial Stage pTa Urothelial Bladder Cancer. <i>Bladder Cancer</i> , 2017, 3, 173-180.	0.2	7
40	Improved prediction of nephron-sparing surgery versus radical nephrectomy by the optimized R.E.N.A.L. Score in patients undergoing surgery for renal masses. <i>Minerva Urologica E Nefrologica = the Italian Journal of Urology and Nephrology</i> , 2019, 71, 249-257.	3.9	7
41	Aquaporin 3 Expression Loss in Urothelial Carcinoma: Association with Tumor Invasion Depth, but not with Grading?. <i>Bladder Cancer</i> , 2017, 3, 31-34.	0.2	6
42	Strong Expression of Cancer-testis Antigens CTAG1B and MAGEA3 Is Correlated with Unfavourable Histopathological Features and MAGEA3 Is Associated with Worse Progression-Free Survival in Urothelial Bladder Cancer. <i>Urologia Internationalis</i> , 2019, 102, 77-82.	0.6	6
43	Characterization of PD-1 and PD-L1 Expression in Papillary Renal Cell Carcinoma: Results of a Large Multicenter Study. <i>Clinical Genitourinary Cancer</i> , 2021, 19, 53-59.e1.	0.9	6
44	Prognostic and Predictive Value of Fibroblast Growth Factor Receptor Alterations in High-grade Non-muscle-invasive Bladder Cancer Treated with and Without Bacillus Calmette-Guérin Immunotherapy. <i>European Urology</i> , 2022, 81, 606-614.	0.9	6
45	Loss of CHEK2 Predicts Progression in Stage pT1 Non-Muscle-Invasive Bladder Cancer (NMIBC). <i>Pathology and Oncology Research</i> , 2020, 26, 1625-1632.	0.9	4
46	Impact of E-Cadherin and β -Catenin as Prognostic Factor in Renal Cell Carcinoma with Tumor Thrombus of the Vena Cava. <i>Urologia Internationalis</i> , 2019, 102, 413-420.	0.6	3
47	cMET: a prognostic marker in papillary renal cell carcinoma?. <i>Human Pathology</i> , 2022, 121, 1-10.	1.1	3
48	BioScore (B7-H1, survivin, and Ki-67) does not predict cancer-specific mortality in surgically treated patients with renal cell carcinoma: An external validation study. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2019, 37, 510-518.	0.8	2
49	Stage T1 bladder cancer: historic background and latest tracks for its demystification. <i>Translational Andrology and Urology</i> , 2018, 7, 760-763.	0.6	1
50	Expression of Low Affinity Nerve Growth Factor Receptor p75 in Classic Bladder Exstrophy. <i>Frontiers in Pediatrics</i> , 2021, 9, 634343.	0.9	1
51	Serine threonine kinase 15 amplification in normal urothelium of cystectomy specimens is no prognostic factor in urothelial carcinoma of the bladder. <i>Pathology Research and Practice</i> , 2011, 207, 161-163.	1.0	0
52	Predictive Value of Positive Surgical Margins after Radical Prostatectomy for Lymph Node Metastasis in Locally Advanced Prostate Carcinoma. <i>Advances in Urology</i> , 2012, 2012, 1-3.	0.6	0
53	Impact of Male Patient Information on Quality of Urine Examination (PIQUE Study). <i>Urologia Internationalis</i> , 2021, , 1-5.	0.6	0