

Wolfgang Otto

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7431320/wolfgang-otto-publications-by-citations.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

50
papers

751
citations

16
h-index

25
g-index

53
ext. papers

1,057
ext. citations

4.1
avg, IF

3.49
L-index

#	Paper	IF	Citations
50	The WHO classification of 1973 is more suitable than the WHO classification of 2004 for predicting survival in pT1 urothelial bladder cancer. <i>BJU International</i> , 2011 , 107, 404-8	5.6	72
49	The Tumor Immune Microenvironment Drives a Prognostic Relevance That Correlates with Bladder Cancer Subtypes. <i>Cancer Immunology Research</i> , 2019 , 7, 923-938	12.5	69
48	Analysis of sex differences in cancer-specific survival and perioperative mortality following radical cystectomy: results of a large German multicenter study of nearly 2500 patients with urothelial carcinoma of the bladder. <i>Gender Medicine</i> , 2012 , 9, 481-9		58
47	Performance of the Food and Drug Administration/EMA-approved programmed cell death ligand-1 assays in urothelial carcinoma with emphasis on therapy stratification for first-line use of atezolizumab and pembrolizumab. <i>European Journal of Cancer</i> , 2019 , 106, 234-243	7.5	39
46	High PDL1 mRNA expression predicts better survival of stage pT1 non-muscle-invasive bladder cancer (NMIBC) patients. <i>Cancer Immunology, Immunotherapy</i> , 2018 , 67, 403-412	7.4	39
45	In stage pT1 non-muscle-invasive bladder cancer (NMIBC), high KRT20 and low KRT5 mRNA expression identify the luminal subtype and predict recurrence and survival. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2017 , 470, 267-274	5.1	36
44	Epithelial-mesenchymal transformation markers E-cadherin and survivin predict progression of stage pTa urothelial bladder carcinoma. <i>World Journal of Urology</i> , 2016 , 34, 709-16	4	27
43	Biochemical Foundations of Health and Energy Conservation in Hibernating Free-ranging Subadult Brown Bear <i>Ursus arctos</i> . <i>Journal of Biological Chemistry</i> , 2016 , 291, 22509-22523	5.4	25
42	mRNA-Expression of and Defines Distinct Prognostic Subgroups of Muscle-Invasive Urothelial Bladder Cancer Correlating with Histological Variants. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	24
41	A multicenter round robin test of PD-L1 expression assessment in urothelial bladder cancer by immunohistochemistry and RT-qPCR with emphasis on prognosis prediction after radical cystectomy. <i>Oncotarget</i> , 2018 , 9, 15001-15014	3.3	22
40	ESR1, ERBB2, and Ki67 mRNA expression predicts stage and grade of non-muscle-invasive bladder carcinoma (NMIBC). <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2016 , 469, 547-552	5.1	20
39	FGFR3 Mutation Status and FGFR3 Expression in a Large Bladder Cancer Cohort Treated by Radical Cystectomy: Implications for Anti-FGFR3 Treatment?. <i>European Urology</i> , 2020 , 78, 682-687	10.2	20
38	No mutations of FGFR3 in normal urothelium in the vicinity of urothelial carcinoma of the bladder harbouring activating FGFR3 mutations in patients with bladder cancer. <i>International Journal of Cancer</i> , 2009 , 125, 2205-8	7.5	19
37	Cytotoxic T-cell-related gene expression signature predicts improved survival in muscle-invasive urothelial bladder cancer patients after radical cystectomy and adjuvant chemotherapy 2020 , 8,		18
36	High Androgen Receptor mRNA Expression Is Independently Associated with Prolonged Cancer-Specific and Recurrence-Free Survival in Stage T1 Bladder Cancer. <i>Translational Oncology</i> , 2017 , 10, 340-345	4.9	17
35	The World Health Organization 1973 classification system for grade is an important prognosticator in T1 non-muscle-invasive bladder cancer. <i>BJU International</i> , 2018 , 122, 978-985	5.6	16
34	CDKN2A as transcriptomic marker for muscle-invasive bladder cancer risk stratification and therapy decision-making. <i>Scientific Reports</i> , 2018 , 8, 14383	4.9	16

33	High proliferation rate and TNM stage but not histomorphological subtype are independent prognostic markers for overall survival in papillary renal cell carcinoma. <i>Human Pathology</i> , 2019 , 83, 212-223	3.7	15
32	Photodynamic diagnosis for superficial bladder cancer: do all risk-groups profit equally from oncological and economic long-term results?. <i>Clinical Medicine Oncology</i> , 2009 , 3, 53-8		14
31	FOXM1 predicts overall and disease specific survival in muscle-invasive urothelial carcinoma and presents a differential expression between bladder cancer subtypes. <i>Oncotarget</i> , 2017 , 8, 47595-47606	3.3	14
30	Metric substage according to micro and extensive lamina propria invasion improves prognostics in T1 bladder cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2018 , 36, 361.e7-361.e13	2.8	14
29	FOXM1 overexpression is associated with adverse outcome and predicts response to intravesical instillation therapy in stage pT1 non-muscle-invasive bladder cancer. <i>BJU International</i> , 2019 , 123, 187-196	5.6	13
28	Expression, localisation and potential significance of aquaporins in benign and malignant human prostate tissue. <i>BMC Urology</i> , 2018 , 18, 75	2.2	13
27	Infiltrative lamina propria invasion pattern as an independent predictor for cancer-specific and overall survival of instillation treatment-naïve stage T1 high-grade urothelial bladder cancer. <i>International Journal of Urology</i> , 2018 , 25, 442-449	2.3	12
26	Predictive value of molecular subtyping in NMIBC by RT-qPCR of ERBB2, ESR1, PGR and MKI67 from formalin fixed TUR biopsies. <i>Oncotarget</i> , 2017 , 8, 67684-67695	3.3	11
25	Comparative analysis of comorbidity and performance indices for prediction of oncological outcomes in patients with upper tract urothelial carcinoma who were treated with radical nephroureterectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 1141-50	2.8	10
24	Loss of AQP3 protein expression is associated with worse progression-free and cancer-specific survival in patients with muscle-invasive bladder cancer. <i>World Journal of Urology</i> , 2015 , 33, 1959-64	4	9
23	Androgen Receptor mRNA Expression in Urothelial Carcinoma of the Bladder: A Retrospective Analysis of Two Independent Cohorts. <i>Translational Oncology</i> , 2019 , 12, 661-668	4.9	8
22	High CDKN2A/p16 and Low FGFR3 Expression Predict Progressive Potential of Stage pT1 Urothelial Bladder Carcinoma. <i>Clinical Genitourinary Cancer</i> , 2018 , 16, 248-256.e2	3.3	8
21	WHO 1973 grade 3 and infiltrative growth pattern proved, aberrant E-cadherin expression tends to be of predictive value for progression in a series of stage T1 high-grade bladder cancer after organ-sparing approach. <i>International Urology and Nephrology</i> , 2017 , 49, 431-437	2.3	7
20	A switch from epithelial to mesenchymal properties correlates with lymphovascular invasion in squamous cell carcinoma of the penis. <i>Pathology Research and Practice</i> , 2015 , 211, 641-5	3.4	7
19	Differentiation-associated urothelial cytochrome P450 oxidoreductase predicates the xenobiotic-metabolizing activity of "luminal" muscle-invasive bladder cancers. <i>Molecular Carcinogenesis</i> , 2018 , 57, 606-618	5	7
18	Analysis of the prognostic relevance of sex-steroid hormonal receptor mRNA expression in muscle-invasive urothelial carcinoma of the urinary bladder. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2019 , 474, 209-217	5.1	7
17	FOXA1 Gene Expression for Defining Molecular Subtypes of Muscle-Invasive Bladder Cancer after Radical Cystectomy. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	7
16	Improved prediction of nephron-sparing surgery versus radical nephrectomy by the optimized R.E.N.A.L. Score in patients undergoing surgery for renal masses. <i>Minerva Urologica E Nefrologica = the Italian Journal of Urology and Nephrology</i> , 2019 , 71, 249-257	4.4	6

15	Strong Expression of Cancer Testis Antigens CTAG1B and MAGEA3 Is Correlated with Unfavourable Histopathological Features and MAGEA3 Is Associated with Worse Progression-Free Survival in Urothelial Bladder Cancer. <i>Urologia Internationalis</i> , 2019 , 102, 77-82	1.9	6
14	Increased Proliferation as Independent Predictor of Disease Recurrence in Initial Stage pTa Urothelial Bladder Cancer. <i>Bladder Cancer</i> , 2017 , 3, 173-180	1	5
13	Aquaporin 3 Expression Loss in Urothelial Carcinoma: Association with Tumor Invasion Depth, but not with Grading?. <i>Bladder Cancer</i> , 2017 , 3, 31-34	1	4
12	High Stroma T-Cell Infiltration is Associated with Better Survival in Stage pT1 Bladder Cancer. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	3
11	Loss of CHEK2 Predicts Progression in Stage pT1 Non-Muscle-Invasive Bladder Cancer (NMIBC). <i>Pathology and Oncology Research</i> , 2020 , 26, 1625-1632	2.6	3
10	Characterization of PD-1 and PD-L1 Expression in Papillary Renal Cell Carcinoma: Results of a Large Multicenter Study. <i>Clinical Genitourinary Cancer</i> , 2021 , 19, 53-59.e1	3.3	3
9	BioScore (B7-H1, survivin, and Ki-67) does not predict cancer-specific mortality in surgically treated patients with renal cell carcinoma: An external validation study. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2019 , 37, 510-518	2.8	2
8	Impact of E-Cadherin and β Catenin as Prognostic Factor in Renal Cell Carcinoma with Tumor Thrombus of the Vena Cava. <i>Urologia Internationalis</i> , 2019 , 102, 413-420	1.9	2
7	Prognostic markers in invasive bladder cancer: FGFR3 mutation status versus P53 and KI-67 expression: a multi-center, multi-laboratory analysis in 1058 radical cystectomy patients.. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2021 ,	2.8	2
6	Tumor budding correlates with tumor invasiveness and predicts worse survival in pT1 non-muscle-invasive bladder cancer. <i>Scientific Reports</i> , 2021 , 11, 17981	4.9	1
5	The Prognostic Value of FGFR3 Expression in Patients with T1 Non-Muscle Invasive Bladder Cancer. <i>Cancer Management and Research</i> , 2021 , 13, 6567-6578	3.6	0
4	Serine threonine kinase 15 amplification in normal urothelium of cystectomy specimens is no prognostic factor in urothelial carcinoma of the bladder. <i>Pathology Research and Practice</i> , 2011 , 207, 161-3	3.4	
3	Predictive value of positive surgical margins after radical prostatectomy for lymph node metastasis in locally advanced prostate carcinoma. <i>Advances in Urology</i> , 2012 , 2012, 618574	1.6	
2	Expression of Low Affinity Nerve Growth Factor Receptor p75 in Classic Bladder Exstrophy. <i>Frontiers in Pediatrics</i> , 2021 , 9, 634343	3.4	
1	Impact of Male Patient Information on Quality of Urine Examination (PIQUE Study). <i>Urologia Internationalis</i> , 2021 , 1-5	1.9	