

Yang Han

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7434235/publications.pdf>

Version: 2024-02-01

41
papers

1,112
citations

516215

16
h-index

414034

32
g-index

56
all docs

56
docs citations

56
times ranked

2474
citing authors

#	ARTICLE	IF	CITATIONS
1	High-Dose Intravenous Immunoglobulin as a Therapeutic Option for Deteriorating Patients With Coronavirus Disease 2019. <i>Open Forum Infectious Diseases</i> , 2020, 7, ofaa102.	0.4	327
2	Association Between Gut Microbiota and CD4 Recovery in HIV-1 Infected Patients. <i>Frontiers in Microbiology</i> , 2018, 9, 1451.	1.5	90
3	Long-Term Persistence of Robust Antibody and Cytotoxic T Cell Responses in Recovered Patients Infected with SARS Coronavirus. <i>PLoS ONE</i> , 2006, 1, e24.	1.1	69
4	Cross-reactive neutralization of SARS-CoV-2 by serum antibodies from recovered SARS patients and immunized animals. <i>Science Advances</i> , 2020, 6, .	4.7	57
5	Significance of serology testing to assist timely diagnosis of SARS-CoV-2 infections: implication from a family cluster. <i>Emerging Microbes and Infections</i> , 2020, 9, 924-927.	3.0	51
6	High-Dose Intravenous Immunoglobulin in Severe Coronavirus Disease 2019: A Multicenter Retrospective Study in China. <i>Frontiers in Immunology</i> , 2021, 12, 627844.	2.2	40
7	Prevalence of hepatitis B and C viruses in HIV-positive patients in China: a cross-sectional study. <i>Journal of the International AIDS Society</i> , 2016, 19, 20659.	1.2	39
8	Phylogenetics of major CRF01_AE epidemic clusters circulating in mainland of China. <i>Scientific Reports</i> , 2017, 7, 6330.	1.6	37
9	A higher CD4/CD8 ratio correlates with an ultralow cell-associated HIV-1 DNA level in chronically infected patients on antiretroviral therapy: a case control study. <i>BMC Infectious Diseases</i> , 2017, 17, 771.	1.3	37
10	Longitudinal profiles of immunoglobulin G antibodies against severe acute respiratory syndrome coronavirus components and neutralizing activities in recovered patients. <i>Scandinavian Journal of Infectious Diseases</i> , 2011, 43, 515-521.	1.5	36
11	Detection of HIV-1 viruses in tears of patients even under long-term HAART. <i>Aids</i> , 2011, 25, 1925-1927.	1.0	31
12	Tryptophan Metabolism Activates Aryl Hydrocarbon Receptor-Mediated Pathway To Promote HIV-1 Infection and Reactivation. <i>MBio</i> , 2019, 10, .	1.8	28
13	Cardiovascular disease risk among Chinese antiretroviral-naïve adults with advanced HIV disease. <i>BMC Infectious Diseases</i> , 2017, 17, 287.	1.3	21
14	Prolonged presence of viral nucleic acid in clinically recovered COVID-19 patients was not associated with effective infectiousness. <i>Emerging Microbes and Infections</i> , 2020, 9, 2315-2321.	3.0	21
15	Super-dominant pathobiontic bacteria in the nasopharyngeal microbiota as causative agents of secondary bacterial infection in influenza patients. <i>Emerging Microbes and Infections</i> , 2020, 9, 605-615.	3.0	18
16	Emergence of Lamivudine-Resistant HBV during Antiretroviral Therapy Including Lamivudine for Patients Coinfected with HIV and HBV in China. <i>PLoS ONE</i> , 2015, 10, e0134539.	1.1	18
17	An antiretroviral regimen containing 6 months of stavudine followed by long-term zidovudine for first-line HIV therapy is optimal in resource-limited settings: a prospective, multicenter study in China. <i>Chinese Medical Journal</i> , 2014, 127, 59-65.	0.9	16
18	The effects of antiretroviral therapy initiation time on HIV reservoir size in Chinese chronically HIV infected patients: a prospective, multi-site cohort study. <i>BMC Infectious Diseases</i> , 2019, 19, 257.	1.3	14

#	ARTICLE	IF	CITATIONS
19	Baseline Naive CD4+ T-cell Level Predicting Immune Reconstitution in Treated HIV-infected Late Presenters. <i>Chinese Medical Journal</i> , 2016, 129, 2683-2690.	0.9	13
20	The prevalence of drug resistance among treatment-naïve HIV-1-infected individuals in China during pre- and post- 2004. <i>BMC Infectious Diseases</i> , 2016, 16, 605.	1.3	12
21	Incidence of hypertension among persons living with HIV in China: a multicenter cohort study. <i>BMC Public Health</i> , 2020, 20, 834.	1.2	10
22	CRF07_BC is associated with slow HIV disease progression in Chinese patients. <i>Scientific Reports</i> , 2022, 12, 3773.	1.6	10
23	HIV sequence diversity during the early phase of infection is associated with HIV DNA reductions during antiretroviral therapy. <i>Journal of Medical Virology</i> , 2017, 89, 982-988.	2.5	9
24	Comparative Transcriptional Analysis Identified Characteristic Genes and Patterns in HIV-Infected Immunological Non-Responders. <i>Frontiers in Immunology</i> , 2022, 13, 807890.	2.2	9
25	Very high baseline HIV viremia impairs efficacy of non-nucleoside reverse transcriptase inhibitor-based ART: a long-term observation in treatment-naïve patients. <i>Infectious Diseases of Poverty</i> , 2020, 9, 75.	1.5	8
26	CD8+ T cell responses specific for hepatitis B virus core protein in patients with chronic hepatitis B virus infection. <i>Journal of Clinical Virology</i> , 2014, 61, 40-46.	1.6	7
27	Week 120 Efficacy of Tenofovir, Lamivudine and Lopinavir/r-Based Second-Line Antiretroviral Therapy in Treatment-Experienced HIV Patients. <i>PLoS ONE</i> , 2015, 10, e0120705.	1.1	7
28	Combined multi-omics and network pharmacology approach reveals the role of Tripterygium Wilfordii Hook F in treating HIV immunological non-responders. <i>Phytomedicine</i> , 2022, 101, 154103.	2.3	7
29	HIV-1 CRF01_AE subtype and HIV-1 DNA level among patients with chronic HIV-1 infection: a correlation study. <i>BMC Infectious Diseases</i> , 2020, 20, 66.	1.3	6
30	Cerebral Vasoreactivity Evaluated by the Breath-Holding Challenge Correlates With Performance on a Cognitive Screening Test in Persons Living With Treated HIV Infection in China. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2018, 79, e101-e104.	0.9	4
31	CD16-expressing monocytes correlate with arterial stiffness in HIV-infected ART-naïve men. <i>HIV Clinical Trials</i> , 2018, 19, 39-45.	2.0	4
32	High APRIL Levels Are Associated With Slow Disease Progression and Low Immune Activation in Chronic HIV-1-Infected Patients. <i>Frontiers in Medicine</i> , 2020, 7, 299.	1.2	4
33	Pharmacodynamics of efavirenz 400 mg in treatment-naïve Chinese HIV-infected patients in a prospective cohort study. <i>BMC Infectious Diseases</i> , 2021, 21, 112.	1.3	4
34	Therapeutic prediction of HIV-1 DNA decay: a multicenter longitudinal cohort study. <i>BMC Infectious Diseases</i> , 2021, 21, 592.	1.3	4
35	Elevated pre-treatment IL-18 level is associated with HBeAg seroconversion in HIV-HBV coinfection. <i>Antiviral Therapy</i> , 2017, 22, 523-527.	0.6	4
36	Comparison of Renal Function Biomarkers of Serum Creatinine and Cystatin C in HIV-Infected People on Dolutegravir-Containing Therapy. <i>Infection and Drug Resistance</i> , 2022, Volume 15, 1695-1706.	1.1	4

#	ARTICLE	IF	CITATIONS
37	Gut lactate-producing bacteria promote CD4 T cell recovery on Anti-retroviral therapy in HIV-infected patients. Computational and Structural Biotechnology Journal, 2021, 19, 2928-2937.	1.9	3
38	Whole blood as an alternative to peripheral blood mononuclear cell for detection of total HIV-1 DNA. BMC Infectious Diseases, 2020, 20, 941.	1.3	2
39	Na ⁺ ve CD4+ cell counts significantly decay and high HIV RNA levels contribute to immunological progression in long-term non-progressors infected with HIV by blood products: a cohort study. BMC Immunology, 2021, 22, 36.	0.9	2
40	HBV pgRNA profiles in Chinese HIV / HBV coinfecting patients under pre- and posttreatment: a multicentre observational cohort study. Journal of Viral Hepatitis, 2022, , .	1.0	2
41	A4.14...Increased bone turnover after switch to tenofovir + lopinavir/ritonavir in chinese HIV + patients. Annals of the Rheumatic Diseases, 2014, 73, A62.1-A62.	0.5	0