

Abdul R Qureshi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7437578/publications.pdf>

Version: 2024-02-01

194
papers

8,548
citations

44042

48
h-index

51562

86
g-index

196
all docs

196
docs citations

196
times ranked

9189
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Factors predicting malnutrition in hemodialysis patients: A cross-sectional study. <i>Kidney International</i> , 1998, 53, 773-782. | 2.6 | 507 |
| 2 | Serum Albumin, C-Reactive Protein, Interleukin 6, and Fetuin A as Predictors of Malnutrition, Cardiovascular Disease, and Mortality in Patients With ESRD. <i>American Journal of Kidney Diseases</i> , 2006, 47, 139-148. | 2.1 | 442 |
| 3 | Comparative Associations of Muscle Mass and Muscle Strength with Mortality in Dialysis Patients. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2014, 9, 1720-1728. | 2.2 | 386 |
| 4 | Obese sarcopenia in patients with end-stage renal disease is associated with inflammation and increased mortality. <i>American Journal of Clinical Nutrition</i> , 2007, 86, 633-638. | 2.2 | 246 |
| 5 | Hand-grip muscle strength, lean body mass, and plasma proteins as markers of nutritional status in patients with chronic renal failure close to start of dialysis therapy. <i>American Journal of Kidney Diseases</i> , 2000, 36, 1213-1225. | 2.1 | 241 |
| 6 | Muscle atrophy, inflammation and clinical outcome in incident and prevalent dialysis patients. <i>Clinical Nutrition</i> , 2008, 27, 557-564. | 2.3 | 230 |
| 7 | Elevated resistin levels in chronic kidney disease are associated with decreased glomerular filtration rate and inflammation, but not with insulin resistance. <i>Kidney International</i> , 2006, 69, 596-604. | 2.6 | 209 |
| 8 | Comparison of nutritional and inflammatory markers in dialysis patients with reduced appetite. <i>American Journal of Clinical Nutrition</i> , 2007, 85, 695-701. | 2.2 | 202 |
| 9 | Truncal fat mass as a contributor to inflammation in end-stage renal disease. <i>American Journal of Clinical Nutrition</i> , 2004, 80, 1222-1229. | 2.2 | 187 |
| 10 | Hyperhomocysteinemia, nutritional status, and cardiovascular disease in hemodialysis patients. <i>Kidney International</i> , 2000, 57, 1727-1735. | 2.6 | 177 |
| 11 | Telomere attrition is associated with inflammation, low fetuin-A levels and high mortality in prevalent haemodialysis patients. <i>Journal of Internal Medicine</i> , 2008, 263, 302-312. | 2.7 | 165 |
| 12 | Cholinergic Anti-Inflammatory Pathway Activity and High Mobility Group Box-1 (HMGB1) Serum Levels in Patients with Rheumatoid Arthritis. <i>Molecular Medicine</i> , 2007, 13, 210-215. | 1.9 | 162 |
| 13 | Accelerated lean body mass loss in incident chronic dialysis patients with diabetes mellitus. <i>Kidney International</i> , 2005, 68, 2368-2374. | 2.6 | 157 |
| 14 | Clinical and biochemical implications of low thyroid hormone levels (total and free forms) in euthyroid patients with chronic kidney disease. <i>Journal of Internal Medicine</i> , 2007, 262, 690-701. | 2.7 | 144 |
| 15 | Prevalence and clinical implications of testosterone deficiency in men with end-stage renal disease. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 184-190. | 0.4 | 144 |
| 16 | Serum Albumin as Predictor of Nutritional Status in Patients with ESRD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012, 7, 1446-1453. | 2.2 | 138 |
| 17 | Biomarkers of Cardiovascular Disease and Mortality Risk in Patients with Advanced CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2016, 11, 1163-1172. | 2.2 | 133 |
| 18 | Abdominal fat deposition is associated with increased inflammation, protein-energy wasting and worse outcome in patients undergoing haemodialysis. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 562-568. | 0.4 | 116 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Comorbidity and Acute Clinical Events as Determinants of C-Reactive Protein Variation in Hemodialysis Patients: Implications for Patient Survival. <i>American Journal of Kidney Diseases</i> , 2009, 53, 1024-1033. | 2.1 | 111 |
| 20 | Measures of chronic kidney disease and risk of incident peripheral artery disease: a collaborative meta-analysis of individual participant data. <i>Lancet Diabetes and Endocrinology</i> , 2017, 5, 718-728. | 5.5 | 110 |
| 21 | Circulating Levels of Visfatin/Pre-B-Cell Colony-Enhancing Factor 1 in Relation to Genotype, GFR, Body Composition, and Survival in Patients With CKD. <i>American Journal of Kidney Diseases</i> , 2007, 49, 237-244. | 2.1 | 109 |
| 22 | Additive Effects of Soluble TWEAK and Inflammation on Mortality in Hemodialysis Patients. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2009, 4, 110-118. | 2.2 | 106 |
| 23 | Albuminuria changes are associated with subsequent risk of end-stage renal disease and mortality. <i>Kidney International</i> , 2017, 91, 244-251. | 2.6 | 104 |
| 24 | Novel Links between the Long Pentraxin 3, Endothelial Dysfunction, and Albuminuria in Early and Advanced Chronic Kidney Disease. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008, 3, 976-985. | 2.2 | 103 |
| 25 | Increased circulating sclerostin levels in end-stage renal disease predict biopsy-verified vascular medial calcification and coronary artery calcification. <i>Kidney International</i> , 2015, 88, 1356-1364. | 2.6 | 102 |
| 26 | Prevalence and recognition of chronic kidney disease in Stockholm healthcare. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, 2086-2094. | 0.4 | 101 |
| 27 | High Mobility Group Box Protein-1 Correlates with Renal Function in Chronic Kidney Disease (CKD). <i>Molecular Medicine</i> , 2008, 14, 109-115. | 1.9 | 92 |
| 28 | The higher mortality associated with low serum albumin is dependent on systemic inflammation in end-stage kidney disease. <i>PLoS ONE</i> , 2018, 13, e0190410. | 1.1 | 91 |
| 29 | Effect of Circulating Soluble Receptor for Advanced Glycation End Products (sRAGE) and the Proinflammatory RAGE Ligand (EN-RAGE, S100A12) on Mortality in Hemodialysis Patients. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2010, 5, 2213-2219. | 2.2 | 83 |
| 30 | Body Fat Mass and Serum Leptin Levels Influence Epoetin Sensitivity in Patients With ESRD. <i>American Journal of Kidney Diseases</i> , 2005, 46, 628-634. | 2.1 | 78 |
| 31 | Whole blood cytokine attenuation by cholinergic agonists <i>in vivo</i> and relationship to vagus nerve activity in rheumatoid arthritis. <i>Journal of Internal Medicine</i> , 2010, 268, 94-101. | 2.7 | 78 |
| 32 | Inflammation and wasting in chronic kidney disease: Partners in crime. <i>Kidney International</i> , 2006, 70, S8-S13. | 2.6 | 77 |
| 33 | Testosterone deficiency is a cause of anaemia and reduced responsiveness to erythropoiesis-stimulating agents in men with chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2012, 27, 709-715. | 0.4 | 74 |
| 34 | The Stockholm CREATinine Measurements (SCREAM) project: protocol overview and regional representativeness. <i>CKJ: Clinical Kidney Journal</i> , 2016, 9, 119-127. | 1.4 | 74 |
| 35 | Trimestral variations of C-reactive protein, interleukin-6 and tumour necrosis factor- α are similarly associated with survival in haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 1313-1318. | 0.4 | 70 |
| 36 | Early Vascular Ageing and Cellular Senescence in Chronic Kidney Disease. <i>Computational and Structural Biotechnology Journal</i> , 2019, 17, 721-729. | 1.9 | 65 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Does statins promote vascular calcification in chronic kidney disease?. European Journal of Clinical Investigation, 2017, 47, 137-148. | 1.7 | 62 |
| 38 | The reverse epidemiology of plasma total homocysteine as a mortality risk factor is related to the impact of wasting and inflammation. Nephrology Dialysis Transplantation, 2006, 22, 209-217. | 0.4 | 61 |
| 39 | Protein-energy wasting modifies the association of ghrelin with inflammation, leptin, and mortality in hemodialysis patients. Kidney International, 2011, 79, 749-756. | 2.6 | 60 |
| 40 | Clinical global assessment of nutritional status as predictor of mortality in chronic kidney disease patients. PLoS ONE, 2017, 12, e0186659. | 1.1 | 60 |
| 41 | Effects of Probiotic Supplementation on Trimethylamine-N-Oxide Plasma Levels in Hemodialysis Patients: a Pilot Study. Probiotics and Antimicrobial Proteins, 2019, 11, 648-654. | 1.9 | 59 |
| 42 | Time in Therapeutic Range and Outcomes After Warfarin Initiation in Newly Diagnosed Atrial Fibrillation Patients With Renal Dysfunction. Journal of the American Heart Association, 2017, 6, . | 1.6 | 57 |
| 43 | The long pentraxin PTX-3 in prevalent hemodialysis patients: associations with comorbidities and mortality. QJM - Monthly Journal of the Association of Physicians, 2008, 101, 397-405. | 0.2 | 55 |
| 44 | Baseline Levels and Trimestral Variation of Triiodothyronine and Thyroxine and Their Association with Mortality in Maintenance Hemodialysis Patients. Clinical Journal of the American Society of Nephrology: CJASN, 2012, 7, 131-138. | 2.2 | 54 |
| 45 | Matrix Gla protein is an independent predictor of both intimal and medial vascular calcification in chronic kidney disease. Scientific Reports, 2020, 10, 6586. | 1.6 | 53 |
| 46 | Plasma S100A12 and soluble receptor of advanced glycation end product levels and mortality in chronic kidney disease Stage 5 patients. Nephrology Dialysis Transplantation, 2015, 30, 84-91. | 0.4 | 52 |
| 47 | CDKN2A/p16INK4a expression is associated with vascular progeria in chronic kidney disease. Aging, 2017, 9, 494-507. | 1.4 | 52 |
| 48 | Elevated Serum Macrophage Migration Inhibitory Factor (MIF) Concentrations in Chronic Kidney Disease (CKD) Are Associated with Markers of Oxidative Stress and Endothelial Activation. Molecular Medicine, 2009, 15, 70-75. | 1.9 | 50 |
| 49 | Metabolic Changes in Summer Active and Anuric Hibernating Free-Ranging Brown Bears (Ursus) Tj ETQq1 1 0.784314 rgBT /Overlock 1.1 50 | | |
| 50 | Clinical determinants of reduced physical activity in hemodialysis and peritoneal dialysis patients. Journal of Nephrology, 2015, 28, 503-510. | 0.9 | 50 |
| 51 | Influence of Body Mass Index on the Association of Weight Changes with Mortality in Hemodialysis Patients. Clinical Journal of the American Society of Nephrology: CJASN, 2013, 8, 1725-1733. | 2.2 | 49 |
| 52 | Inverse Relationship between the Inflammatory Marker Pentraxin-3, Fat Body Mass, and Abdominal Obesity in End-Stage Renal Disease. Clinical Journal of the American Society of Nephrology: CJASN, 2011, 6, 2785-2791. | 2.2 | 47 |
| 53 | Essential polyunsaturated fatty acids, inflammation and mortality in dialysis patients. Nephrology Dialysis Transplantation, 2012, 27, 3615-3620. | 0.4 | 47 |
| 54 | Reduced skeletal muscle expression of mitochondrial-derived peptides humanin and MOTS-C and Nrf2 in chronic kidney disease. American Journal of Physiology - Renal Physiology, 2019, 317, F1122-F1131. | 1.3 | 44 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Dialysis modality and nutritional status are associated with variability of inflammatory markers. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, 1320-1327. | 0.4 | 42 |
| 56 | Vertebral bone density associates with coronary artery calcification and is an independent predictor of poor outcome in end-stage renal disease patients. <i>Bone</i> , 2016, 92, 50-57. | 1.4 | 42 |
| 57 | Circulating proteins as predictors of cardiovascular mortality in end-stage renal disease. <i>Journal of Nephrology</i> , 2019, 32, 111-119. | 0.9 | 42 |
| 58 | Elevated Circulating Levels and Tissue Expression of Pentraxin 3 in Uremia: A Reflection of Endothelial Dysfunction. <i>PLoS ONE</i> , 2013, 8, e63493. | 1.1 | 41 |
| 59 | Malnutrition and inflammation are associated with impaired pulmonary function in patients with chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2004, 19, 1823-1828. | 0.4 | 40 |
| 60 | Clinical importance of an elevated circulating chemerin level in incident dialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 4017-4023. | 0.4 | 40 |
| 61 | Circulating vascular endothelial growth factor (VEGF) and its soluble receptor 1 (sVEGFR-1) are associated with inflammation and mortality in incident dialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2013, 28, 2356-2363. | 0.4 | 39 |
| 62 | Oxidative Dna Damage and Mortality in Hemodialysis and Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2015, 35, 206-215. | 1.1 | 37 |
| 63 | Vitamin D Deficiency in Dialysis Patients: Effect of Dialysis Modality and Implications on Outcome. , 2010, 20, 359-367. | | 36 |
| 64 | Validation of insulin sensitivity surrogate indices and prediction of clinical outcomes in individuals with and without impaired renal function. <i>Kidney International</i> , 2014, 86, 383-391. | 2.6 | 36 |
| 65 | Bone mineral density and mortality in end-stage renal disease patients. <i>CKJ: Clinical Kidney Journal</i> , 2020, 13, 307-321. | 1.4 | 36 |
| 66 | Nonthyroidal illness: a risk factor for coronary calcification and arterial stiffness in patients undergoing peritoneal dialysis?. <i>Journal of Internal Medicine</i> , 2013, 274, 584-593. | 2.7 | 34 |
| 67 | IGF-1 and Survival in ESRD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2014, 9, 120-127. | 2.2 | 34 |
| 68 | Uric acid is not associated with decline in renal function or time to renal replacement therapy initiation in a referred cohort of patients with Stage III, IV and V chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, 2039-2045. | 0.4 | 34 |
| 69 | Plasma Pentosidine and Its Association with Mortality in Patients with Chronic Kidney Disease. <i>PLoS ONE</i> , 2016, 11, e0163826. | 1.1 | 34 |
| 70 | Lung Dysfunction and Mortality in Patients with Chronic Kidney Disease. <i>Kidney and Blood Pressure Research</i> , 2018, 43, 522-535. | 0.9 | 33 |
| 71 | Health-related quality of life as predictor of mortality in end-stage renal disease patients: an observational study. <i>BMC Nephrology</i> , 2019, 20, 144. | 0.8 | 33 |
| 72 | Subclinical versus overt obesity in dialysis patients: more than meets the eye. <i>Nephrology Dialysis Transplantation</i> , 2013, 28, iv175-iv181. | 0.4 | 32 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Associations between Thyroid Hormones, Calcification Inhibitor Levels and Vascular Calcification in End-Stage Renal Disease. PLoS ONE, 2015, 10, e0132353. | 1.1 | 31 |
| 74 | Plasma pentosidine and total homocysteine levels in relation to change in common carotid intima-media area in the first year of dialysis therapy. Clinical Nephrology, 2006, 66, 418-425. | 0.4 | 29 |
| 75 | Serum albumin, inflammation, and nutrition in end-stage renal disease: C-reactive protein is needed for optimal assessment. Seminars in Dialysis, 2018, 31, 435-439. | 0.7 | 28 |
| 76 | Association between levels of pentraxin 3 and incidence of chronic kidney disease in the elderly. Journal of Internal Medicine, 2016, 279, 173-179. | 2.7 | 27 |
| 77 | Total and bone-specific alkaline phosphatase are associated with bone mineral density over time in end-stage renal disease patients starting dialysis. Journal of Nephrology, 2017, 30, 255-262. | 0.9 | 27 |
| 78 | Determinants and survival implications of low bone mineral density in end-stage renal disease patients. Journal of Nephrology, 2013, 26, 485-494. | 0.9 | 27 |
| 79 | Chemerin inhibits vascular calcification through ChemR23 and is associated with lower coronary calcium in chronic kidney disease. Journal of Internal Medicine, 2019, 286, 449-457. | 2.7 | 26 |
| 80 | Bone Mineral Density in End-Stage Renal Disease Patients: Association with Wasting, Cardiovascular Disease and Mortality. Blood Purification, 2008, 26, 284-290. | 0.9 | 25 |
| 81 | Serum 8-hydroxydeoxyguanosine, a marker of oxidative DNA damage, is associated with mortality independent of inflammation in chronic kidney disease. European Journal of Internal Medicine, 2019, 68, 60-65. | 1.0 | 25 |
| 82 | Skin autofluorescence, arterial stiffness and Framingham risk score as predictors of clinical outcome in chronic kidney disease patients: a cohort study. Nephrology Dialysis Transplantation, 2019, 34, 442-448. | 0.4 | 25 |
| 83 | Low levels of IgM antibodies against phosphorylcholine-A increase mortality risk in patients undergoing haemodialysis. Nephrology Dialysis Transplantation, 2009, 24, 3454-3460. | 0.4 | 24 |
| 84 | Impact of Baseline Health-Related Quality of Life Scores on Survival of Incident Patients on Peritoneal Dialysis: A Cohort Study. Nephron, 2015, 129, 97-103. | 0.9 | 24 |
| 85 | Trends in haemoglobin, erythropoietin-stimulating agents and iron use in Swedish chronic kidney disease patients between 2008 and 2013. Nephrology Dialysis Transplantation, 2016, 31, 628-635. | 0.4 | 24 |
| 86 | Lower serum calcium is independently associated with CKD progression. Scientific Reports, 2018, 8, 5148. | 1.6 | 24 |
| 87 | Restrictive lung disorder is common in patients with kidney failure and associates with protein-energy wasting, inflammation and cardiovascular disease. PLoS ONE, 2018, 13, e0195585. | 1.1 | 23 |
| 88 | N-Terminal Pro-Brain Natriuretic Peptide Independently Predicts Protein Energy Wasting and Is Associated with All-Cause Mortality in Prevalent HD Patients. American Journal of Nephrology, 2009, 29, 516-523. | 1.4 | 22 |
| 89 | C-reactive Protein: Repeated Measurements will Improve Dialysis Patient Care. Seminars in Dialysis, 2016, 29, 7-14. | 0.7 | 22 |
| 90 | New Algorithm for the Management of Orbital Blowout Fracture Based on Prospective Study. Craniomaxillofacial Trauma & Reconstruction, 2018, 11, 285-295. | 0.6 | 21 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Insights in the regulation of trimethylamine N-oxide production using a comparative biomimetic approach suggest a metabolic switch in hibernating bears. <i>Scientific Reports</i> , 2020, 10, 20323. | 1.6 | 21 |
| 92 | The influence of hepatitis C and iron replacement therapy on plasma pentosidine levels in haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2004, 19, 3112-3116. | 0.4 | 20 |
| 93 | Associations between the CYBA 242C/T and the MPO 463G/A Polymorphisms, Oxidative Stress and Cardiovascular Disease in Chronic Kidney Disease Patients. <i>Blood Purification</i> , 2007, 25, 210-218. | 0.9 | 20 |
| 94 | Health-related quality of life in peritoneal dialysis patients: A narrative review. <i>Seminars in Dialysis</i> , 2019, 32, 452-462. | 0.7 | 20 |
| 95 | Inverse J-shaped relation between coronary arterial calcium density and mortality in advanced chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, 1202-1211. | 0.4 | 20 |
| 96 | Bone mineral density at different sites and 5 years mortality in end-stage renal disease patients: A cohort study. <i>Bone</i> , 2020, 130, 115075. | 1.4 | 20 |
| 97 | Temporal discrepancies in the association between the apoB/apoA ratio and mortality in incident dialysis patients. <i>Journal of Internal Medicine</i> , 2009, 265, 708-716. | 2.7 | 19 |
| 98 | Self-Rated Appetite as a Predictor of Mortality in Patients With Stage 5 Chronic Kidney Disease. , 2013, 23, 106-113. | | 19 |
| 99 | Increased Levels of Modified Advanced Oxidation Protein Products are Associated with Central and Peripheral Blood Pressure in Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2015, 35, 460-470. | 1.1 | 19 |
| 100 | Offering Patients Therapy Options in Unplanned Start (OPTiONS): Implementation of an educational program is feasible and effective. <i>BMC Nephrology</i> , 2017, 18, 18. | 0.8 | 19 |
| 101 | Nutritional status, muscle composition and plasma and muscle free amino acids in renal transplant patients. <i>Clinical Nephrology</i> , 1994, 42, 237-45. | 0.4 | 19 |
| 102 | Fractures after kidney transplantation: Incidence, predictors, and association with mortality. <i>Bone</i> , 2020, 140, 115554. | 1.4 | 18 |
| 103 | Postprandial metabolic response to a fat- and carbohydrate-rich meal in patients with chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 2231-2237. | 0.4 | 16 |
| 104 | Increased Telomere Attrition After Renal Transplantation—Impact of Antimetabolite Therapy. <i>Transplantation Direct</i> , 2016, 2, e116. | 0.8 | 16 |
| 105 | Major fractures after initiation of dialysis: Incidence, predictors and association with mortality. <i>Bone</i> , 2020, 133, 115242. | 1.4 | 16 |
| 106 | Factors influencing access to education, decision making, and receipt of preferred dialysis modality in unplanned dialysis start patients. <i>Patient Preference and Adherence</i> , 2016, Volume 10, 2229-2237. | 0.8 | 15 |
| 107 | Association of Serum Sclerostin with Bone Sclerostin in Chronic Kidney Disease is Lost in Glucocorticoid Treated Patients. <i>Calcified Tissue International</i> , 2019, 104, 214-223. | 1.5 | 15 |
| 108 | Low renal replacement therapy incidence among slowly progressing elderly chronic kidney disease patients referred to nephrology care: an observational study. <i>BMC Nephrology</i> , 2017, 18, 59. | 0.8 | 14 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Fractures and their sequelae in non-dialysis-dependent chronic kidney disease: the Stockholm CREATinine Measurement project. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, 1908-1915. | 0.4 | 14 |
| 110 | Sevelamer Use in End-Stage Kidney Disease (ESKD) Patients Associates with Poor Vitamin K Status and High Levels of Gut-Derived Uremic Toxins: A Drugâ€“Bug Interaction?. <i>Toxins</i> , 2020, 12, 351. | 1.5 | 14 |
| 111 | Functional vitamin K insufficiency, vascular calcification and mortality in advanced chronic kidney disease: A cohort study. <i>PLoS ONE</i> , 2021, 16, e0247623. | 1.1 | 14 |
| 112 | Bicarbonate-Based Peritoneal Dialysis Solution has Less Effect on Ingestive Behavior than Lactate-Based Peritoneal Dialysis Solution. <i>Peritoneal Dialysis International</i> , 2009, 29, 656-663. | 1.1 | 13 |
| 113 | Longitudinal Changes in Health-Related Quality of Life Scores in Brazilian Incident Peritoneal Dialysis Patients (Brazpd): Socio-Economic Status Not a Barrier. <i>Peritoneal Dialysis International</i> , 2013, 33, 687-696. | 1.1 | 13 |
| 114 | High alkaline phosphatase and low intact parathyroid hormone associate with worse clinical outcome in peritoneal dialysis patients. <i>Peritoneal Dialysis International</i> , 2021, 41, 236-243. | 1.1 | 13 |
| 115 | Three-month variation of plasma pentraxin 3 compared with C-reactive protein, albumin and homocysteine levels in haemodialysis patients. <i>CKJ: Clinical Kidney Journal</i> , 2014, 7, 373-379. | 1.4 | 12 |
| 116 | Cholinergic anti-inflammatory pathway activity in dialysis patients: a role for neuroimmunomodulation?. <i>CKJ: Clinical Kidney Journal</i> , 2015, 8, 599-605. | 1.4 | 12 |
| 117 | Inflammation down-regulates CYP3A4-catalysed drug metabolism in hemodialysis patients. <i>BMC Pharmacology & Toxicology</i> , 2018, 19, 33. | 1.0 | 12 |
| 118 | Comparative Analysis Between Computed Tomography and Surrogate Methods to Detect Low Muscle Mass Among Colorectal Cancer Patients. <i>Journal of Parenteral and Enteral Nutrition</i> , 2020, 44, 1328-1337. | 1.3 | 12 |
| 119 | Impaired postprandial fibroblast growth factor (FGF)-19 response in patients with stage 5 chronic kidney diseases is ameliorated following antioxidative therapy. <i>Nephrology Dialysis Transplantation</i> , 2013, 28, iv212-iv219. | 0.4 | 11 |
| 120 | Nonesterified Fatty Acids and Cardiovascular Mortality in Elderly Men with CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2015, 10, 584-591. | 2.2 | 11 |
| 121 | Elevated Circulating S100A12 Associates with Vascular Disease and Worse Clinical Outcome in Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2016, 36, 269-276. | 1.1 | 11 |
| 122 | Telomere Attrition and Elongation after Chronic Dialysis Initiation in Patients with End-Stage Renal Disease. <i>Blood Purification</i> , 2016, 41, 25-33. | 0.9 | 11 |
| 123 | Peritonitis: Episode Sequence, Microbiological Variation, Risk Factors and Clinical Outcomes in a North China Peritoneal Dialysis Center. <i>Kidney and Blood Pressure Research</i> , 2018, 43, 1573-1584. | 0.9 | 11 |
| 124 | Pro-neurotensin depends on renal function and is related to all-cause mortality in chronic kidney disease. <i>European Journal of Endocrinology</i> , 2020, 183, 233-244. | 1.9 | 11 |
| 125 | Longitudinal genome-wide DNA methylation changes in response to kidney failure replacement therapy. <i>Scientific Reports</i> , 2022, 12, 470. | 1.6 | 11 |
| 126 | Delta-He: a novel marker of inflammation predicting mortality and ESA response in peritoneal dialysis patients. <i>CKJ: Clinical Kidney Journal</i> , 2014, 7, 275-281. | 1.4 | 10 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | Serum hepatocyte growth factor is associated with truncal fat mass and increased mortality in chronic kidney disease stage 5 patients with protein-energy wasting. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, 274-282. | 0.4 | 10 |
| 128 | The cholinergic anti-inflammatory pathway in resistant hypertension treated with renal denervation. <i>Molecular Medicine</i> , 2019, 25, 39. | 1.9 | 10 |
| 129 | Phenotypic features of vascular calcification in chronic kidney disease. <i>Journal of Internal Medicine</i> , 2020, 287, 422-434. | 2.7 | 10 |
| 130 | Aortic Valve Calcium Associates with All-Cause Mortality Independent of Coronary Artery Calcium and Inflammation in Patients with End-Stage Renal Disease. <i>Journal of Clinical Medicine</i> , 2020, 9, 607. | 1.0 | 10 |
| 131 | Fatores associados À qualidade de vida de pacientes incidentes em diálise peritoneal no Brasil (BRAZPD). <i>Jornal Brasileiro De Nefrologia: Orgao Oficial De Sociedades Brasileira E Latino-Americana De Nefrologia</i> , 2011, 33, 38-44. | 0.4 | 10 |
| 132 | Type of Referral, Dialysis Start and Choice of Renal Replacement Therapy Modality in an International Integrated Care Setting. <i>PLoS ONE</i> , 2016, 11, e0155987. | 1.1 | 9 |
| 133 | Initiation of erythropoiesis-stimulating agents and outcomes: a nationwide observational cohort study in anaemic chronic kidney disease patients. <i>Nephrology Dialysis Transplantation</i> , 2017, 32, gfw328. | 0.4 | 9 |
| 134 | Copeptin is independently associated with vascular calcification in chronic kidney disease stage 5. <i>BMC Nephrology</i> , 2020, 21, 43. | 0.8 | 9 |
| 135 | High-sensitivity troponins in dialysis patients: variation and prognostic value. <i>CKJ: Clinical Kidney Journal</i> , 2021, 14, 1789-1797. | 1.4 | 9 |
| 136 | Scoring of medial arterial calcification predicts cardiovascular events and mortality after kidney transplantation. <i>Journal of Internal Medicine</i> , 2022, 291, 813-823. | 2.7 | 9 |
| 137 | Tryptophan and its metabolites in patients on continuous ambulatory peritoneal dialysis and following renal transplantation. <i>Nephrology Dialysis Transplantation</i> , 1994, 9, 791-6. | 0.4 | 9 |
| 138 | The increase in renal replacement therapy (RRT) incidence has come to an end in Sweden—analysis of variations by region over the period 1991-2010. <i>CKJ: Clinical Kidney Journal</i> , 2013, 6, 352-357. | 1.4 | 8 |
| 139 | Genotypic and phenotypic predictors of inflammation in patients with chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, 2033-2040. | 0.4 | 8 |
| 140 | Bone mineral density of extremities is associated with coronary calcification and biopsy-verified vascular calcification in living-donor renal transplant recipients. <i>Journal of Bone and Mineral Metabolism</i> , 2017, 35, 536-543. | 1.3 | 8 |
| 141 | Plasma Beta-Trace Protein as a Marker of Residual Renal Function: The Effect of Different Hemodialysis Modalities and Intra-Individual Variability over Time. <i>Kidney and Blood Pressure Research</i> , 2017, 42, 877-885. | 0.9 | 8 |
| 142 | Dialysis Access, Infections, and Hospitalisations in Unplanned Dialysis Start Patients: Results from the Options Study. <i>International Journal of Artificial Organs</i> , 2017, 40, 48-59. | 0.7 | 8 |
| 143 | Serum Glutaredoxin Activity as a Marker of Oxidative Stress in Chronic Kidney Disease: A Pilot Study. <i>Nephron</i> , 2018, 140, 249-256. | 0.9 | 8 |
| 144 | Differences in association of lower bone mineral density with higher coronary calcification in female and male end-stage renal disease patients. <i>BMC Nephrology</i> , 2019, 20, 59. | 0.8 | 8 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 145 | Causes of death across categories of estimated glomerular filtration rate: The Stockholm CREATinine Measurements (SCREAM) project. PLoS ONE, 2019, 14, e0209440. | 1.1 | 8 |
| 146 | Effects of acute fructose loading on levels of serum uric acid—a pilot study. European Journal of Clinical Investigation, 2019, 49, e13040. | 1.7 | 8 |
| 147 | Incidence of Fractures Before and After Dialysis Initiation. Journal of Bone and Mineral Research, 2020, 35, 2372-2380. | 3.1 | 8 |
| 148 | Different subclasses and isotypes of antibodies against phosphorylcholine in haemodialysis patients: association with mortality. Clinical and Experimental Immunology, 2020, 201, 94-104. | 1.1 | 8 |
| 149 | Association between reduced kidney function and incident hypoglycaemia in people with diabetes: The Stockholm Creatinine Measurements (SCREAM) project. Diabetes, Obesity and Metabolism, 2020, 22, 1425-1435. | 2.2 | 8 |
| 150 | Time-dependent lipid profile inversely associates with mortality in hemodialysis patients— independent of inflammation/malnutrition. Journal of Internal Medicine, 2021, 290, 910-921. | 2.7 | 8 |
| 151 | Determinants of N-Terminal Pro-Brain Natriuretic Peptide Variation in Hemodialysis Patients and Prediction of Survival. Blood Purification, 2014, 37, 138-145. | 0.9 | 7 |
| 152 | Prospective Randomized Controlled Pilot Study on Orbital Blowout Fracture. Craniomaxillofacial Trauma & Reconstruction, 2018, 11, 165-171. | 0.6 | 7 |
| 153 | Dialysis Adequacy Indices and Body Composition in Male and Female Patients on Peritoneal Dialysis. Peritoneal Dialysis International, 2014, 34, 417-425. | 1.1 | 6 |
| 154 | Sclerostin—A Debutant on the Autosomal Dominant Polycystic Kidney Disease Scene?. Kidney International Reports, 2017, 2, 481-485. | 0.4 | 6 |
| 155 | Pregnancy-associated plasma protein-A predicts survival in end-stage renal disease— confounding and modifying effects of cardiovascular disease, body composition and inflammation. Nephrology Dialysis Transplantation, 2018, 33, 971-977. | 0.4 | 6 |
| 156 | Sparing effect of peritoneal dialysis vs hemodialysis on BMD changes and its impact on mortality. Journal of Bone and Mineral Metabolism, 2021, 39, 260-269. | 1.3 | 6 |
| 157 | Increased Monocyte/Lymphocyte Ratio as Risk Marker for Cardiovascular Events and Infectious Disease Hospitalization in Dialysis Patients. Blood Purification, 2022, 51, 747-755. | 0.9 | 6 |
| 158 | Circulating Alpha-Tocopherol and Insulin Sensitivity Among Older Men With Chronic Kidney Disease. , 2016, 26, 177-182. | | 5 |
| 159 | Pentraxin-3 — a potential biomarker in ANCA-associated vasculitis. Scandinavian Journal of Rheumatology, 2023, 52, 293-301. | 0.6 | 5 |
| 160 | Selection of Genetic and Phenotypic Features Associated with Inflammatory Status of Patients on Dialysis Using Relaxed Linear Separability Method. PLoS ONE, 2014, 9, e86630. | 1.1 | 4 |
| 161 | Younger patients and smokers report a higher level of pain after knee arthroscopy: a clinical and experimental study including synovial metabolism. Knee Surgery, Sports Traumatology, Arthroscopy, 2019, 27, 471-477. | 2.3 | 4 |
| 162 | Antibodies against Malondialdehyde in Haemodialysis Patients and Its Association with Clinical Outcomes: Differences between Subclasses and Isotypes. Journal of Clinical Medicine, 2020, 9, 753. | 1.0 | 4 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 163 | Secular trends in hip fracture incidence and subsequent mortality in dialysis patients and the general population in Sweden. <i>Bone</i> , 2021, 147, 115909. | 1.4 | 4 |
| 164 | Observational study of risk factors associated with clinical outcome among elderly kidney transplant recipients in Sweden – a decade of follow-up. <i>Transplant International</i> , 2021, 34, 2363-2370. | 0.8 | 4 |
| 165 | Risk of hospitalization associated with body mass index and weight changes among prevalent haemodialysis patients. <i>Nefrologia</i> , 2018, 38, 520-527. | 0.2 | 3 |
| 166 | Risk of hospitalization associated with body mass index and weight changes among prevalent haemodialysis patients. <i>Nefrologia</i> , 2018, 38, 520-527. | 0.2 | 3 |
| 167 | Factors associated with the quality of life of incident patients on PD in Brazil (BRAZPD). <i>Jornal Brasileiro De Nefrologia: Orgao Oficial De Sociedades Brasileira E Latino-Americana De Nefrologia</i> , 2011, 33, 38-44. | 0.4 | 3 |
| 168 | Do metabolic derangements in end-stage polycystic kidney disease differ versus other primary kidney diseases?. <i>Nephrology</i> , 2018, 23, 31-36. | 0.7 | 2 |
| 169 | Inter-leg systolic blood pressure difference predicts cardiovascular events and mortality in incident hemodialysis patients. <i>International Journal of Artificial Organs</i> , 2020, 43, 217-224. | 0.7 | 2 |
| 170 | The impact of body mass index (BMI) variation on mortality of incident elderly patients on peritoneal dialysis: a joint model analysis. <i>Jornal Brasileiro De Nefrologia: Orgao Oficial De Sociedades Brasileira E Latino-Americana De Nefrologia</i> , 2017, 39, 267-274. | 0.4 | 2 |
| 171 | The Novel Bone Alkaline Phosphatase Isoform B1x Is Associated with Improved 5-Year Survival in Chronic Kidney Disease. <i>Nutrients</i> , 2021, 13, 4402. | 1.7 | 2 |
| 172 | P1389MAJOR OSTEOPOROTIC FRACTURES AFTER INITIATION OF DIALYSIS: INCIDENCE, PREDICTORS AND ASSOCIATION WITH MORTALITY. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, . | 0.4 | 1 |
| 173 | FC 123RENAL TRANSPLANTATION MITIGATES INCREASED BIOLOGICAL (EPIGENETIC) AGE IN CHRONIC KIDNEY DISEASE. <i>Nephrology Dialysis Transplantation</i> , 2021, 36, . | 0.4 | 1 |
| 174 | The Volume Difference Along the External Surface of the Zygomatic Bone. <i>Journal of Craniofacial Surgery</i> , 2021, Publish Ahead of Print, . | 0.3 | 1 |
| 175 | Effects of Probiotic Supplementation on Trimethylamine-N-Oxide Plasma Levels in Hemodialysis Patients: a Pilot Study. , 2019, 11, 648. | | 1 |
| 176 | The Impact of Occasional vs Persistent Inflammation on the Survival of HD Patients. <i>Hemodialysis International</i> , 2003, 7, 73-104. | 0.4 | 0 |
| 177 | Similar Increase Plasma Asymmetric Dimethylarginine (P-ADMA) and in Non Renal patients with Abnormal Myocardial Perfusion and in Hemodialysis Patients.. <i>Hemodialysis International</i> , 2003, 7, 73-104. | 0.4 | 0 |
| 178 | FP331INITIATION OF ERYTHROPOEISIS STIMULATING AGENTS AND MORTALITY IN A LARGE REFERRED CKD COHORT. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, iii179-iii179. | 0.4 | 0 |
| 179 | SP242GLOMERULAR FILTRATION RATE ESTIMATION USING BETA TRACE PROTEIN: EXTERNAL VALIDATION OF THREE EQUATIONS. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, i167-i167. | 0.4 | 0 |
| 180 | SP286PLASMA BETA TRACE PROTEIN PREDICTS MORTALITY IN CKD: RELATION WITH ENDOTHELIAL DYSFUNCTION. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, i183-i184. | 0.4 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 181 | SP306PREVALENCE, DIAGNOSIS AND NEPHROLOGY CARE OF CKD IN THE REGION OF STOCKHOLM. Nephrology Dialysis Transplantation, 2016, 31, i191-i192. | 0.4 | 0 |
| 182 | FP634HIGH ALL CAUSE AND CVD MORTALITY IN AN INCIDENT COHORT OF HEMODIALYSIS PATIENTS WITH LOW SERUM ALBUMIN AND INFLAMMATION. Nephrology Dialysis Transplantation, 2018, 33, i257-i257. | 0.4 | 0 |
| 183 | SP298HIGHER MEAN CORPUSCULAR VOLUME ASSOCIATES WITH POOR CLINICAL OUTCOME IN CKD5 PATIENTS. Nephrology Dialysis Transplantation, 2018, 33, i444-i444. | 0.4 | 0 |
| 184 | FP336HIGHER NEUTROPHIL TO LYMPHOCYTE RATIO ASSOCIATES WITH POOR CLINICAL OUTCOME IN CKD5 PATIENTS. Nephrology Dialysis Transplantation, 2018, 33, i144-i145. | 0.4 | 0 |
| 185 | SP323HIGH FIBRINOGEN LEVELS ARE INDEPENDENTLY ASSOCIATED WITH INCREASED MORTALITY IN PATIENTS WITH CHRONIC KIDNEY DISEASE (CKD). Nephrology Dialysis Transplantation, 2018, 33, i452-i452. | 0.4 | 0 |
| 186 | P1409HIP FRACTURE TRENDS IN SWEDISH GENERAL POPULATION AND DIALYSIS PATIENTS. Nephrology Dialysis Transplantation, 2020, 35, . | 0.4 | 0 |
| 187 | P0347COMPLEMENT FACTOR BB AND FACTOR C4D IN IGA NEPHROPATHY AND IGA VASCULITIS. Nephrology Dialysis Transplantation, 2020, 35, . | 0.4 | 0 |
| 188 | P0669HIGH ESTIMATED PHENOTYPIC AGE ASSOCIATES WITH WORSE CLINICAL OUTCOME IN CHRONIC KIDNEY DISEASE PATIENTS. Nephrology Dialysis Transplantation, 2020, 35, . | 0.4 | 0 |
| 189 | P0673INVERSE ASSOCIATION OF TRANSFERRIN SATURATION WITH MORTALITY RISK IN CHRONIC KIDNEY DISEASE. Nephrology Dialysis Transplantation, 2020, 35, . | 0.4 | 0 |
| 190 | P0791MATRIX GLA PROTEIN AND PREMATURE VASCULAR CALCIFICATION IN PATIENTS WITH END-STAGE RENAL DISEASE. Nephrology Dialysis Transplantation, 2020, 35, . | 0.4 | 0 |
| 191 | MO576ASSOCIATION OF BONE MATERIAL STRENGTH WITH BONE DENSITY IN PATIENTS WITH END-STAGE RENAL DISEASE. Nephrology Dialysis Transplantation, 2021, 36, . | 0.4 | 0 |
| 192 | MO673: Association of Mean Corpuscular Volume with Mortality in Kidney Failure Patients: Opposite Findings in Chinese and Swedish Kidney Failure Patients. Nephrology Dialysis Transplantation, 2022, 37, . | 0.4 | 0 |
| 193 | MO708: Association of Atherogenic Index of Plasma (AIP) with Mortality in Chinese and Swedish End-Stage Kidney Disease Patients: Results from Two Cohorts. Nephrology Dialysis Transplantation, 2022, 37, . | 0.4 | 0 |
| 194 | MO681: Association of Framingham Risk Score with Mortality in Kidney Failure Patients from China and Sweden. Nephrology Dialysis Transplantation, 2022, 37, . | 0.4 | 0 |