

# Peter Cresswell

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/7439398/peter-cresswell-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

150  
papers

19,677  
citations

68  
h-index

140  
g-index

154  
ext. papers

21,370  
ext. citations

15.4  
avg, IF

6.92  
L-index

#	Paper	IF	Citations
150	Nilabh Shastri (1952-2021). <i>Immunity</i> , <b>2021</b> , 54, 389-390	32.3	1
149	Translational shutdown and evasion of the innate immune response by SARS-CoV-2 NSP14 protein. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2021</b> , 118,	11.5	43
148	SARS-CoV-2 exacerbates proinflammatory responses in myeloid cells through C-type lectin receptors and Tweety family member 2. <i>Immunity</i> , <b>2021</b> , 54, 1304-1319.e9	32.3	41
147	Impact of Calreticulin and Its Mutants on Endoplasmic Reticulum Function in Health and Disease. <i>Progress in Molecular and Subcellular Biology</i> , <b>2021</b> , 59, 163-180	3	
146	Quantitating Endosomal Escape of a Library of Polymers for mRNA Delivery. <i>Nano Letters</i> , <b>2020</b> , 20, 1117-1123	11.5	31
145	Disruption of mosGILT in <i>Anopheles gambiae</i> impairs ovarian development and Plasmodium infection. <i>Journal of Experimental Medicine</i> , <b>2020</b> , 217,	16.6	12
144	HLA tapasin independence: broader peptide repertoire and HIV control. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2020</b> , 117, 28232-28238	11.5	18
143	A personal retrospective on the mechanisms of antigen processing. <i>Immunogenetics</i> , <b>2019</b> , 71, 141-160	3.2	11
142	Intrinsic expression of viperin regulates thermogenesis in adipose tissues. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2019</b> , 116, 17419-17428	11.5	13
141	Proteasomal degradation within endocytic organelles mediates antigen cross-presentation. <i>EMBO Journal</i> , <b>2019</b> , 38, e99266	13	24
140	A mosquito salivary gland protein partially inhibits Plasmodium sporozoite cell traversal and transmission. <i>Nature Communications</i> , <b>2018</b> , 9, 2908	17.4	21
139	A novel probe to assess cytosolic entry of exogenous proteins. <i>Nature Communications</i> , <b>2018</b> , 9, 3104	17.4	11
138	Cytosolic Processing Governs TAP-Independent Presentation of a Critical Melanoma Antigen. <i>Journal of Immunology</i> , <b>2018</b> , 201, 1875-1888	5.3	15
137	Tumor-associated calreticulin variants functionally compromise the peptide loading complex and impair its recruitment of MHC-I. <i>Journal of Biological Chemistry</i> , <b>2018</b> , 293, 9555-9569	5.4	40
136	miRNA-mediated TUSC3 deficiency enhances UPR and ERAD to promote metastatic potential of NSCLC. <i>Nature Communications</i> , <b>2018</b> , 9, 5110	17.4	22
135	The ongoing saga of the mechanism(s) of MHC class I-restricted cross-presentation. <i>Current Opinion in Immunology</i> , <b>2017</b> , 46, 89-96	7.8	32
134	Structural studies of viperin, an antiviral radical SAM enzyme. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2017</b> , 114, 6806-6811	11.5	46

133	Editing peptide presentation to T cells. <i>Science</i> , <b>2017</b> , 358, 992-993	33.3	1
132	Sec61 blockade by mycolactone inhibits antigen cross-presentation independently of endosome-to-cytosol export. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2017</b> , 114, E5910-E5919	11.5	52
131	Viperin interaction with mitochondrial antiviral signaling protein (MAVS) limits viperin-mediated inhibition of the interferon response in macrophages. <i>PLoS ONE</i> , <b>2017</b> , 12, e0172236	3.7	20
130	Antigen Processing and Presentation Mechanisms in Myeloid Cells. <i>Microbiology Spectrum</i> , <b>2016</b> , 4,	8.9	20
129	The transcription factor TFEB acts as a molecular switch that regulates exogenous antigen-presentation pathways. <i>Nature Immunology</i> , <b>2015</b> , 16, 729-36	19.1	89
128	Are ERAD components involved in cross-presentation?. <i>Molecular Immunology</i> , <b>2015</b> , 68, 112-5	4.3	20
127	Three tapasin docking sites in TAP cooperate to facilitate transporter stabilization and heterodimerization. <i>Journal of Immunology</i> , <b>2014</b> , 192, 2480-94	5.3	16
126	Invariant chain-MHC class II complexes: always odd and never invariant. <i>Immunology and Cell Biology</i> , <b>2014</b> , 92, 471-2	5	21
125	A congenital disorder of deglycosylation: biochemical characterization of N-glycanase 1 deficiency in patient fibroblasts (607.3). <i>FASEB Journal</i> , <b>2014</b> , 28, 607.3	0.9	
124	Expanding roles for GILT in immunity. <i>Current Opinion in Immunology</i> , <b>2013</b> , 25, 103-8	7.8	48
123	Pathways of antigen processing. <i>Annual Review of Immunology</i> , <b>2013</b> , 31, 443-73	34.7	889
122	In vitro reconstitution of the MHC class I peptide-loading complex. <i>Methods in Molecular Biology</i> , <b>2013</b> , 960, 67-79	1.4	3
121	Viperin regulates cellular lipid metabolism during human cytomegalovirus infection. <i>PLoS Pathogens</i> , <b>2013</b> , 9, e1003497	7.6	78
120	Deglycosylation-dependent fluorescent proteins provide unique tools for the study of ER-associated degradation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2013</b> , 110, 3393-8	11.5	33
119	Endoplasmic reticulum glycoprotein quality control regulates CD1d assembly and CD1d-mediated antigen presentation. <i>Journal of Biological Chemistry</i> , <b>2013</b> , 288, 16391-16402	5.4	6
118	Critical residues in the PMEL/Pmel17 N-terminus direct the hierarchical assembly of melanosomal fibrils. <i>Molecular Biology of the Cell</i> , <b>2013</b> , 24, 964-81	3.5	32
117	Intracellular regulation of cross-presentation during dendritic cell maturation. <i>PLoS ONE</i> , <b>2013</b> , 8, e76803	3.7	9
116	MHC class II-restricted presentation of the major house dust mite allergen Der p 1 Is GILT-dependent: implications for allergic asthma. <i>PLoS ONE</i> , <b>2013</b> , 8, e51343	3.7	19

115	Inefficient exogenous loading of a tapasin-dependent peptide onto HLA-B*44:02 can be improved by acid treatment or fixation of target cells. <i>European Journal of Immunology</i> , <b>2012</b> , 42, 1417-28	6.1	6
114	Intracellular events regulating cross-presentation. <i>Frontiers in Immunology</i> , <b>2012</b> , 3, 138	8.4	22
113	A switch in pathogenic mechanism in myelin oligodendrocyte glycoprotein-induced experimental autoimmune encephalomyelitis in IFN- $\beta$ -inducible lysosomal thiol reductase-free mice. <i>Journal of Immunology</i> , <b>2012</b> , 188, 6001-9	5.3	14
112	Dynamics of major histocompatibility complex class I association with the human peptide-loading complex. <i>Journal of Biological Chemistry</i> , <b>2012</b> , 287, 31172-84	5.4	40
111	Interleukin-2 signalling is modulated by a labile disulfide bond in the CD132 chain of its receptor. <i>Open Biology</i> , <b>2012</b> , 2, 110036	7	31
110	Disulfide reduction in the endocytic pathway: immunological functions of gamma-interferon-inducible lysosomal thiol reductase. <i>Antioxidants and Redox Signaling</i> , <b>2011</b> , 15, 657-684	8.4	69
109	A role for UDP-glucose glycoprotein glucosyltransferase in expression and quality control of MHC class I molecules. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2011</b> , 108, 4956-61	11.5	56
108	Viperin: a multifunctional, interferon-inducible protein that regulates virus replication. <i>Cell Host and Microbe</i> , <b>2011</b> , 10, 534-9	23.4	154
107	Human cytomegalovirus directly induces the antiviral protein viperin to enhance infectivity. <i>Science</i> , <b>2011</b> , 332, 1093-7	33.3	139
106	Labile disulfide bonds are common at the leucocyte cell surface. <i>Open Biology</i> , <b>2011</b> , 1, 110010	7	63
105	Viperin mRNA is a novel target for the human RNase MRP/RNase P endoribonuclease. <i>Cellular and Molecular Life Sciences</i> , <b>2011</b> , 68, 2469-80	10.3	27
104	Essential glycan-dependent interactions optimize MHC class I peptide loading. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2011</b> , 108, 4950-5	11.5	64
103	Proprotein convertases process Pmel17 during secretion. <i>Journal of Biological Chemistry</i> , <b>2011</b> , 286, 9323-37	3.7	39
102	The interferon-inducible gene viperin restricts West Nile virus pathogenesis. <i>Journal of Virology</i> , <b>2011</b> , 85, 11557-66	6.6	112
101	Defective cross-presentation of viral antigens in GILT-free mice. <i>Science</i> , <b>2010</b> , 328, 1394-8	33.3	105
100	Endoplasmic reticulum export, subcellular distribution, and fibril formation by Pmel17 require an intact N-terminal domain junction. <i>Journal of Biological Chemistry</i> , <b>2010</b> , 285, 16166-83	5.4	22
99	Calreticulin controls the rate of assembly of CD1d molecules in the endoplasmic reticulum. <i>Journal of Biological Chemistry</i> , <b>2010</b> , 285, 38283-92	5.4	9
98	GILT accelerates autoimmunity to the melanoma antigen tyrosinase-related protein 1. <i>Journal of Immunology</i> , <b>2010</b> , 185, 2828-35	5.3	38

97	The antiviral protein, viperin, localizes to lipid droplets via its N-terminal amphipathic alpha-helix. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2009</b> , 106, 20452-7	11.5	168
96	Natural lipid ligands associated with human CD1d targeted to different subcellular compartments. <i>Journal of Immunology</i> , <b>2009</b> , 182, 4784-91	5.3	79
95	The N-terminal amphipathic alpha-helix of viperin mediates localization to the cytosolic face of the endoplasmic reticulum and inhibits protein secretion. <i>Journal of Biological Chemistry</i> , <b>2009</b> , 284, 4705-12	5.4	108
94	Receptor-mediated phagocytosis elicits cross-presentation in nonprofessional antigen-presenting cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2009</b> , 106, 3324-9	11.5	50
93	Functional significance of tapasin membrane association and disulfide linkage to ERp57 in MHC class I presentation. <i>European Journal of Immunology</i> , <b>2009</b> , 39, 2371-6	6.1	22
92	Insights into MHC class I peptide loading from the structure of the tapasin-ERp57 thiol oxidoreductase heterodimer. <i>Immunity</i> , <b>2009</b> , 30, 21-32	32.3	214
91	Kinetics and cellular site of glycolipid loading control the outcome of natural killer T cell activation. <i>Immunity</i> , <b>2009</b> , 30, 888-98	32.3	143
90	Viperin is required for optimal Th2 responses and T-cell receptor-mediated activation of NF-kappaB and AP-1. <i>Blood</i> , <b>2009</b> , 113, 3520-9	2.2	52
89	GILT is a critical host factor for <i>Listeria monocytogenes</i> infection. <i>Nature</i> , <b>2008</b> , 455, 1244-7	50.4	114
88	Hsp90-mediated cytosolic refolding of exogenous proteins internalized by dendritic cells. <i>EMBO Journal</i> , <b>2008</b> , 27, 201-11	13	62
87	The quality control of MHC class I peptide loading. <i>Current Opinion in Cell Biology</i> , <b>2008</b> , 20, 624-31	9	156
86	Regulation of MHC class I assembly and peptide binding. <i>Annual Review of Cell and Developmental Biology</i> , <b>2008</b> , 24, 343-68	12.6	156
85	The redox activity of ERp57 is not essential for its functions in MHC class I peptide loading. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2008</b> , 105, 10477-82	11.5	52
84	Target peptide sequence within infectious human immunodeficiency virus type 1 does not ensure envelope-specific T-helper cell reactivation: influences of cysteine protease and gamma interferon-induced thiol reductase activities. <i>Vaccine Journal</i> , <b>2008</b> , 15, 713-9		25
83	C19orf48 encodes a minor histocompatibility antigen recognized by CD8+ cytotoxic T cells from renal cell carcinoma patients. <i>Clinical Cancer Research</i> , <b>2008</b> , 14, 5260-9	12.9	42
82	Selective loading of high-affinity peptides onto major histocompatibility complex class I molecules by the tapasin-ERp57 heterodimer. <i>Nature Immunology</i> , <b>2007</b> , 8, 873-81	19.1	183
81	Innate immune recognition triggers secretion of lysosomal enzymes by macrophages. <i>Traffic</i> , <b>2007</b> , 8, 1179-89	5.7	56
80	Aggregate formation by ERp57-deficient MHC class I peptide-loading complexes. <i>Traffic</i> , <b>2007</b> , 8, 1530-42	3.7	20

79	Saposin B is the dominant saposin that facilitates lipid binding to human CD1d molecules. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2007</b> , 104, 5551-6	11.5	88
78	Severe tryptophan starvation blocks onset of conventional persistence and reduces reactivation of <i>Chlamydia trachomatis</i> . <i>Infection and Immunity</i> , <b>2007</b> , 75, 5105-17	3.7	67
77	The interferon-inducible protein viperin inhibits influenza virus release by perturbing lipid rafts. <i>Cell Host and Microbe</i> , <b>2007</b> , 2, 96-105	23.4	333
76	Functional requirements for the lysosomal thiol reductase GILT in MHC class II-restricted antigen processing. <i>Journal of Immunology</i> , <b>2006</b> , 177, 8569-77	5.3	47
75	Exposure of the promonocytic cell line THP-1 to <i>Escherichia coli</i> induces IFN-gamma-inducible lysosomal thiol reductase expression by inflammatory cytokines. <i>Journal of Immunology</i> , <b>2006</b> , 177, 4833-40	5.2	51
74	An N-linked glycan modulates the interaction between the CD1d heavy chain and beta 2-microglobulin. <i>Journal of Biological Chemistry</i> , <b>2006</b> , 281, 40369-78	5.4	22
73	A role for the endoplasmic reticulum protein retrotranslocation machinery during crosspresentation by dendritic cells. <i>Immunity</i> , <b>2006</b> , 25, 607-17	32.3	234
72	Stoichiometric tapasin interactions in the catalysis of major histocompatibility complex class I molecule assembly. <i>Immunology</i> , <b>2005</b> , 114, 346-53	7.8	26
71	Access of soluble antigens to the endoplasmic reticulum can explain cross-presentation by dendritic cells. <i>Nature Immunology</i> , <b>2005</b> , 6, 107-13	19.1	154
70	Tapasin and ERp57 form a stable disulfide-linked dimer within the MHC class I peptide-loading complex. <i>EMBO Journal</i> , <b>2005</b> , 24, 3613-23	13	141
69	Mechanisms of MHC class I-restricted antigen processing and cross-presentation. <i>Immunological Reviews</i> , <b>2005</b> , 207, 145-57	11.3	333
68	Antigen processing and presentation. <i>Immunological Reviews</i> , <b>2005</b> , 207, 5-7	11.3	47
67	Differential requirements for endosomal reduction in the presentation of two H2-E(d)-restricted epitopes from influenza hemagglutinin. <i>Journal of Immunology</i> , <b>2004</b> , 172, 6607-14	5.3	30
66	Lipid-protein interactions: biosynthetic assembly of CD1 with lipids in the endoplasmic reticulum is evolutionarily conserved. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2004</b> , 101, 1022-6	11.5	67
65	Major histocompatibility complex class I molecules expressed with monoglucosylated N-linked glycans bind calreticulin independently of their assembly status. <i>Journal of Biological Chemistry</i> , <b>2004</b> , 279, 25112-21	5.4	38
64	Cell biology. Cutting and pasting antigenic peptides. <i>Science</i> , <b>2004</b> , 304, 525-7	33.3	17
63	Saposins facilitate CD1d-restricted presentation of an exogenous lipid antigen to T cells. <i>Nature Immunology</i> , <b>2004</b> , 5, 175-81	19.1	185
62	Cellular mechanisms governing cross-presentation of exogenous antigens. <i>Nature Immunology</i> , <b>2004</b> , 5, 678-84	19.1	313

61	Recent developments in MHC-class-I-mediated antigen presentation. <i>Current Opinion in Immunology</i> , <b>2004</b> , 16, 82-9	7.8	69
60	Regulation of MHC class I transport in human dendritic cells and the dendritic-like cell line KG-1. <i>Journal of Immunology</i> , <b>2003</b> , 170, 4178-88	5.3	90
59	Tapasin is a facilitator, not an editor, of class I MHC peptide binding. <i>Journal of Immunology</i> , <b>2003</b> , 171, 5287-95	5.3	89
58	Early phagosomes in dendritic cells form a cellular compartment sufficient for cross presentation of exogenous antigens. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2003</b> , 100, 12889-94	11.5	301
57	Regulation of intracellular trafficking of human CD1d by association with MHC class II molecules. <i>EMBO Journal</i> , <b>2002</b> , 21, 1650-60	13	100
56	Identification of specific glycoforms of major histocompatibility complex class I heavy chains suggests that class I peptide loading is an adaptation of the quality control pathway involving calreticulin and ERp57. <i>Journal of Biological Chemistry</i> , <b>2002</b> , 277, 46415-23	5.4	49
55	Role of the C-terminal propeptide in the activity and maturation of gamma -interferon-inducible lysosomal thiol reductase (GILT). <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2002</b> , 99, 12298-303	11.5	22
54	Absence of gamma-interferon-inducible lysosomal thiol reductase in melanomas disrupts T cell recognition of select immunodominant epitopes. <i>Journal of Experimental Medicine</i> , <b>2002</b> , 195, 1267-77	16.6	113
53	Calnexin, calreticulin, and ERp57 cooperate in disulfide bond formation in human CD1d heavy chain. <i>Journal of Biological Chemistry</i> , <b>2002</b> , 277, 44838-44	5.4	90
52	Disulfide bond isomerization and the assembly of MHC class I-peptide complexes. <i>Immunity</i> , <b>2002</b> , 16, 87-98	32.3	190
51	Multiple species express thiol oxidoreductases related to GILT. <i>Immunogenetics</i> , <b>2001</b> , 53, 342-6	3.2	38
50	A role for calnexin in the assembly of the MHC class I loading complex in the endoplasmic reticulum. <i>Journal of Immunology</i> , <b>2001</b> , 166, 1703-9	5.3	92
49	Defective antigen processing in GILT-free mice. <i>Science</i> , <b>2001</b> , 294, 1361-5	33.3	213
48	Glycosylation and the immune system. <i>Science</i> , <b>2001</b> , 291, 2370-6	33.3	1289
47	Intracellular surveillance: controlling the assembly of MHC class I-peptide complexes. <i>Traffic</i> , <b>2000</b> , 1, 301-5	5.7	62
46	Gamma-interferon-inducible lysosomal thiol reductase (GILT). Maturation, activity, and mechanism of action. <i>Journal of Biological Chemistry</i> , <b>2000</b> , 275, 25907-14	5.4	150
45	Cytomegalovirus US2 destroys two components of the MHC class II pathway, preventing recognition by CD4+ T cells. <i>Nature Medicine</i> , <b>1999</b> , 5, 1039-43	50.5	217
44	The nature of the MHC class I peptide loading complex. <i>Immunological Reviews</i> , <b>1999</b> , 172, 21-8	11.3	264

43	Thiol oxidation and reduction in MHC-restricted antigen processing and presentation. <i>Immunologic Research</i> , <b>1999</b> , 19, 191-200	4.3	18
42	Human epidermal Langerhans cells lack functional mannose receptors and a fully developed endosomal/lysosomal compartment for loading of HLA class II molecules. <i>European Journal of Immunology</i> , <b>1999</b> , 29, 571-80	6.1	45
41	The N-terminal region of tapasin is required to stabilize the MHC class I loading complex. <i>European Journal of Immunology</i> , <b>1999</b> , 29, 1858-70	6.1	139
40	The thiol oxidoreductase ERp57 is a component of the MHC class I peptide-loading complex. <i>Current Biology</i> , <b>1998</b> , 8, 709-12	6.3	157
39	Genomic analysis of the Tapasin gene, located close to the TAP loci in the MHC. <i>European Journal of Immunology</i> , <b>1998</b> , 28, 459-67	6.1	66
38	Calnexin expression does not enhance the generation of MHC class I-peptide complexes. <i>European Journal of Immunology</i> , <b>1998</b> , 28, 907-13	6.1	25
37	Elucidation of the genetic basis of the antigen presentation defects in the mutant cell line .220 reveals polymorphism and alternative splicing of the tapasin gene. <i>European Journal of Immunology</i> , <b>1998</b> , 28, 3783-91	6.1	43
36	Soluble tapasin restores MHC class I expression and function in the tapasin-negative cell line .220. <i>Immunity</i> , <b>1998</b> , 8, 221-31	32.3	248
35	HLA-B27-restricted antigen presentation in the absence of tapasin reveals polymorphism in mechanisms of HLA class I peptide loading. <i>Immunity</i> , <b>1998</b> , 8, 531-42	32.3	224
34	Mechanisms of MHC class I-restricted antigen processing. <i>Annual Review of Immunology</i> , <b>1998</b> , 16, 323-58	34.7	861
33	Genomic analysis of the Tapasin gene, located close to the TAP loci in the MHC <b>1998</b> , 28, 459		2
32	Calnexin expression does not enhance the generation of MHC class I-peptide complexes <b>1998</b> , 28, 907		1
31	A critical role for tapasin in the assembly and function of multimeric MHC class I-TAP complexes. <i>Science</i> , <b>1997</b> , 277, 1306-9	33.3	436
30	Negative regulation by HLA-DO of MHC class II-restricted antigen processing. <i>Science</i> , <b>1997</b> , 278, 106-9	33.3	196
29	Regulation of MHC class I heterodimer stability and interaction with TAP by tapasin. <i>Immunogenetics</i> , <b>1997</b> , 46, 477-83	3.2	73
28	Invariant chain structure and MHC class II function. <i>Cell</i> , <b>1996</b> , 84, 505-7	56.2	297
27	Roles for calreticulin and a novel glycoprotein, tapasin, in the interaction of MHC class I molecules with TAP. <i>Immunity</i> , <b>1996</b> , 5, 103-14	32.3	589
26	Processing and delivery of peptides presented by MHC class I molecules. <i>Current Opinion in Immunology</i> , <b>1996</b> , 8, 59-67	7.8	140



25	HLA-DM interactions with intermediates in HLA-DR maturation and a role for HLA-DM in stabilizing empty HLA-DR molecules. <i>Journal of Experimental Medicine</i> , <b>1996</b> , 184, 2153-65	16.6	191
24	HLA-DM induces CLIP dissociation from MHC class II alpha beta dimers and facilitates peptide loading. <i>Cell</i> , <b>1995</b> , 82, 155-65	56.2	612
23	Assembly, transport, and function of MHC class II molecules. <i>Annual Review of Immunology</i> , <b>1994</b> , 12, 259-93	34.7	689
22	MHC class I/beta 2-microglobulin complexes associate with TAP transporters before peptide binding. <i>Nature</i> , <b>1994</b> , 368, 864-7	50.4	341
21	Human transporters associated with antigen processing possess a promiscuous peptide-binding site. <i>Immunity</i> , <b>1994</b> , 1, 7-14	32.3	154
20	In vivo and in vitro formation and dissociation of HLA-DR complexes with invariant chain-derived peptides. <i>Immunity</i> , <b>1994</b> , 1, 763-74	32.3	172
19	Assembly and intracellular transport of HLA-DM and correction of the class II antigen-processing defect in T2 cells. <i>Immunity</i> , <b>1994</b> , 1, 595-606	32.3	230
18	Transport properties of free and MHC class II-associated oligomers containing different isoforms of human invariant chain. <i>International Immunology</i> , <b>1994</b> , 6, 439-51	4.9	55
17	Assembly and transport of class I MHC-peptide complexes. <i>Novartis Foundation Symposium</i> , <b>1994</b> , 187, 150-62; discussion 162-9		5
16	Presentation of viral antigen by MHC class I molecules is dependent on a putative peptide transporter heterodimer. <i>Nature</i> , <b>1992</b> , 355, 644-6	50.4	313
15	HLA-A2 molecules in an antigen-processing mutant cell contain signal sequence-derived peptides. <i>Nature</i> , <b>1992</b> , 356, 443-6	50.4	456
14	Proteasome subunits encoded in the MHC are not generally required for the processing of peptides bound by MHC class I molecules. <i>Nature</i> , <b>1992</b> , 360, 171-4	50.4	194
13	HLA-DR molecules from an antigen-processing mutant cell line are associated with invariant chain peptides. <i>Nature</i> , <b>1992</b> , 360, 474-7	50.4	343
12	Invariant chain association with HLA-DR molecules inhibits immunogenic peptide binding. <i>Nature</i> , <b>1990</b> , 345, 615-8	50.4	435
11	Co-localization of molecules involved in antigen processing and presentation in an early endocytic compartment. <i>Nature</i> , <b>1990</b> , 343, 133-9	50.4	350
10	Differential transport requirements of HLA and H-2 class I glycoproteins. <i>Immunogenetics</i> , <b>1989</b> , 29, 380-32		134
9	An epitope common to HLA class I and class II antigens, Ig light chains, and beta 2-microglobulin. <i>Immunogenetics</i> , <b>1987</b> , 25, 228-33	3.2	37
8	Immune recognition of human major histocompatibility antigens: localization by a comprehensive synthetic strategy of the continuous antigenic sites in the first domain of HLA-DR2 beta chain. <i>European Journal of Immunology</i> , <b>1987</b> , 17, 497-502	6.1	15

7	Genes regulating HLA class I antigen expression in T-B lymphoblast hybrids. <i>Immunogenetics</i> , <b>1985</b> , 21, 235-46	3.2	515
6	Expression of T-lymphoblast-encoded HLA-DR antigens on human T-B lymphoblast hybrids. <i>Immunogenetics</i> , <b>1983</b> , 17, 411-25	3.2	14
5	Expression of cell surface lectins on activated human lymphoid cells. <i>European Journal of Immunology</i> , <b>1982</b> , 12, 570-6	6.1	25
4	Modulation of cell surface iron transferrin receptors by cellular density and state of activation. <i>Journal of Supramolecular Structure</i> , <b>1979</b> , 11, 579-86		320
3	Antisera to human B-lymphocyte membrane glycoproteins block stimulation in mixed lymphocyte culture. <i>Nature</i> , <b>1975</b> , 257, 147-9	50.4	85
2	The small subunit of HL-A antigens is beta 2-microglobulin. <i>Journal of Experimental Medicine</i> , <b>1973</b> , 138, 1608-12	16.6	342
1	Antigen Processing and Presentation Mechanisms in Myeloid Cells 209-223		2