

Hideaki Miyake

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7450596/hideaki-miyake-publications-by-citations.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

313 papers	5,196 citations	36 h-index	54 g-index
330 ext. papers	5,828 ext. citations	3 avg, IF	5.5 L-index

#	Paper	IF	Citations
313	Overexpression of insulin-like growth factor binding protein-5 helps accelerate progression to androgen-independence in the human prostate LNCaP tumor model through activation of phosphatidylinositol 3-kinase pathway. <i>Endocrinology</i> , 2000 , 141, 2257-65	4.8	125
312	Alternative nonsteroidal antiandrogen therapy for advanced prostate cancer that relapsed after initial maximum androgen blockade. <i>Journal of Urology</i> , 2008 , 180, 921-7	2.5	116
311	Enhanced radiation sensitivity in prostate cancer by inhibition of the cell survival protein clusterin. <i>Clinical Cancer Research</i> , 2002 , 8, 3276-84	12.9	104
310	Overexpression of Bcl-2 in bladder cancer cells inhibits apoptosis induced by cisplatin and adenoviral-mediated p53 gene transfer. <i>Oncogene</i> , 1998 , 16, 933-43	9.2	100
309	Chemosensitization of human renal cell cancer using antisense oligonucleotides targeting the antiapoptotic gene clusterin. <i>Neoplasia</i> , 2001 , 3, 360-7	6.4	96
308	A novel antisense oligonucleotide inhibiting several antiapoptotic Bcl-2 family members induces apoptosis and enhances chemosensitivity in androgen-independent human prostate cancer PC3 cells. <i>Molecular Cancer Therapeutics</i> , 2005 , 4, 1689-98	6.1	92
307	Stress protein GRP78 prevents apoptosis induced by calcium ionophore, ionomycin, but not by glycosylation inhibitor, tunicamycin, in human prostate cancer cells. <i>Journal of Cellular Biochemistry</i> , 2000 , 77, 396-408	4.7	92
306	Overexpression of clusterin in transitional cell carcinoma of the bladder is related to disease progression and recurrence. <i>Urology</i> , 2002 , 59, 150-4	1.6	87
305	Use of antisense oligonucleotides targeting the cytoprotective gene, clusterin, to enhance androgen- and chemo-sensitivity in prostate cancer. <i>World Journal of Urology</i> , 2005 , 23, 38-46	4	84
304	Synergistic chemosensitization and inhibition of progression to androgen independence by antisense Bcl-2 oligodeoxynucleotide and paclitaxel in the LNCaP prostate tumor model. <i>International Journal of Cancer</i> , 2001 , 91, 846-50	7.5	84
303	Serum matrix metalloproteinase-2 and its density in men with prostate cancer as a new predictor of disease extension. <i>International Journal of Cancer</i> , 1998 , 79, 96-101	7.5	80
302	Castration-induced increases in insulin-like growth factor-binding protein 2 promotes proliferation of androgen-independent human prostate LNCaP tumors. <i>Cancer Research</i> , 2003 , 63, 3575-84	10.1	78
301	Comparison of quality of life following laparoscopic and open prostatectomy for prostate cancer. <i>Journal of Urology</i> , 2003 , 169, 2045-8	2.5	77
300	Basic Fibroblast Growth Factor Regulates Matrix Metalloproteinases Production and In Vitro Invasiveness in Human Bladder Cancer Cell Lines. <i>Journal of Urology</i> , 1997 , 157, 2351-2355	2.5	72
299	Risk factors for intravesical recurrence after surgical management of transitional cell carcinoma of the upper urinary tract. <i>Urology</i> , 2008 , 71, 123-7	1.6	70
298	Protection of androgen-dependent human prostate cancer cells from oxidative stress-induced DNA damage by overexpression of clusterin and its modulation by androgen. <i>Prostate</i> , 2004 , 61, 318-23	4.2	67
297	Expression of the secreted form of clusterin protein in renal cell carcinoma as a predictor of disease extension. <i>BJU International</i> , 2005 , 96, 895-9	5.6	61

296	Expression of potential molecular markers in prostate cancer: correlation with clinicopathological outcomes in patients undergoing radical prostatectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2010 , 28, 145-51	2.8	58
295	OVER EXPRESSION OF CLUSTERIN IS AN INDEPENDENT PROGNOSTIC FACTOR FOR NONPAPILLARY RENAL CELL CARCINOMA. <i>Journal of Urology</i> , 2002 , 167, 703-706	2.5	57
294	Inhibition of progression to androgen-independence by combined adjuvant treatment with antisense BCL-XL and antisense Bcl-2 oligonucleotides plus taxol after castration in the Shionogi tumor model. <i>International Journal of Cancer</i> , 2000 , 86, 855-62	7.5	57
293	Expression of epithelial-mesenchymal transition markers in renal cell carcinoma: impact on prognostic outcomes in patients undergoing radical nephrectomy. <i>BJU International</i> , 2012 , 110, E1131-7	5.6	55
292	Detection of micrometastases in pelvic lymph nodes in patients undergoing radical cystectomy for locally invasive bladder cancer by real-time reverse transcriptase-PCR for cytokeratin 19 and uroplakin II. <i>Clinical Cancer Research</i> , 2005 , 11, 3773-7	12.9	55
291	Clinical outcome of maximum androgen blockade using flutamide as second-line hormonal therapy for hormone-refractory prostate cancer. <i>BJU International</i> , 2005 , 96, 791-5	5.6	53
290	Polymeric micellar paclitaxel phosphorylates Bcl-2 and induces apoptotic regression of androgen-independent LNCaP prostate tumors. <i>Prostate</i> , 2000 , 44, 156-63	4.2	52
289	Quantitative detection of micrometastases in pelvic lymph nodes in patients with clinically localized prostate cancer by real-time reverse transcriptase-PCR. <i>Clinical Cancer Research</i> , 2007 , 13, 1192-7	12.9	51
288	Expression patterns of epithelial-mesenchymal transition markers in localized prostate cancer: significance in clinicopathological outcomes following radical prostatectomy. <i>BJU International</i> , 2013 , 111, 30-7	5.6	50
287	Antisense oligodeoxynucleotide therapy targeting clusterin gene for prostate cancer: Vancouver experience from discovery to clinic. <i>International Journal of Urology</i> , 2005 , 12, 785-94	2.3	45
286	SIGNIFICANCE OF MATRIX METALLOPROTEINASES AND TISSUE INHIBITORS OF METALLOPROTEINASE EXPRESSION IN THE RECURRENCE OF SUPERFICIAL TRANSITIONAL CELL CARCINOMA OF THE BLADDER. <i>Journal of Urology</i> , 2001 , 165, 1769-1772	2.5	43
285	Expression of major heat shock proteins in prostate cancer: correlation with clinicopathological outcomes in patients undergoing radical prostatectomy. <i>Journal of Urology</i> , 2007 , 177, 757-61	2.5	41
284	Expression profile of E-cadherin and N-cadherin in non-muscle-invasive bladder cancer as a novel predictor of intravesical recurrence following transurethral resection. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2012 , 30, 161-6	2.8	40
283	Abnormalities of thyroid function in Japanese patients with metastatic renal cell carcinoma treated with sorafenib: a prospective evaluation. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2010 , 28, 515-9	2.8	40
282	A comparison of the biological features between prostate cancers arising in the transition and peripheral zones. <i>BJU International</i> , 2005 , 96, 528-32	5.6	40
281	Comparison of the transperitoneal and retroperitoneal approach in robot-assisted partial nephrectomy in an initial case series in Japan. <i>Journal of Endourology</i> , 2013 , 27, 1384-8	2.7	39
280	Acquisition of resistance to Fas-mediated apoptosis by overexpression of clusterin in human renal-cell carcinoma cells. <i>Molecular Urology</i> , 2001 , 5, 105-11		39
279	Metastatic collecting duct carcinoma of the kidney responded to sunitinib. <i>International Journal of Clinical Oncology</i> , 2011 , 16, 153-5	4.2	38

278	Synergistic chemosensitization and inhibition of tumor growth and metastasis by adenovirus-mediated P53 gene transfer in human bladder cancer model. <i>Urology</i> , 2000 , 56, 332-6	1.6	37
277	Synergistic antitumor activity by combined treatment with gemcitabine and antisense oligodeoxynucleotide targeting clusterin gene in an intravesical administration model against human bladder cancer kotcc-1 cells. <i>Journal of Urology</i> , 2004 , 171, 2477-81	2.5	36
276	Predictors of prostate cancer on repeat transrectal ultrasound-guided systematic prostate biopsy. <i>International Journal of Urology</i> , 2003 , 10, 68-71	2.3	36
275	Beyond simple castration: targeting the molecular basis of treatment resistance in advanced prostate cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2005 , 56 Suppl 1, 47-57	3.5	36
274	Acquired resistance to sunitinib in human renal cell carcinoma cells is mediated by constitutive activation of signal transduction pathways associated with tumour cell proliferation. <i>BJU International</i> , 2013 , 112, E211-20	5.6	35
273	Introduction of clusterin gene into human renal cell carcinoma cells enhances their resistance to cytotoxic chemotherapy through inhibition of apoptosis both in vitro and in vivo. <i>Japanese Journal of Cancer Research</i> , 2001 , 92, 1220-4		35
272	Prognostic significance of oxidative DNA damage evaluated by 8-hydroxy-2'-deoxyguanosine in patients undergoing radical nephrectomy for renal cell carcinoma. <i>Urology</i> , 2004 , 64, 1057-61	1.6	34
271	Expression level of vascular endothelial growth factor receptor-2 in radical nephrectomy specimens as a prognostic predictor in patients with metastatic renal cell carcinoma treated with sunitinib. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 493-8	2.8	33
270	Resistance to cytotoxic chemotherapy-induced apoptosis in human prostate cancer cells is associated with intracellular clusterin expression. <i>Oncology Reports</i> , 2003 , 10, 469-73	3.5	33
269	Microscopic venous invasion in renal cell carcinoma as a predictor of recurrence after radical surgery. <i>International Journal of Urology</i> , 2004 , 11, 264-8	2.3	32
268	Expression of clusterin in prostate cancer correlates with Gleason score but not with prognosis in patients undergoing radical prostatectomy without neoadjuvant hormonal therapy. <i>Urology</i> , 2006 , 68, 609-14	1.6	31
267	Clinical outcome of bacillus Calmette-Guérin perfusion therapy for carcinoma in situ of the upper urinary tract. <i>International Journal of Urology</i> , 2002 , 9, 677-80	2.3	31
266	Synergistic enhancement of resistance to cisplatin in human bladder cancer cells by overexpression of mutant-type p53 and Bcl-2. <i>Journal of Urology</i> , 1999 , 162, 2176-81	2.5	31
265	Significance of Aurora-A expression in renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2007 , 25, 128-33	2.8	30
264	Novel therapeutic strategy for advanced prostate cancer using antisense oligodeoxynucleotides targeting anti-apoptotic genes upregulated after androgen withdrawal to delay androgen-independent progression and enhance chemosensitivity. <i>International Journal of Urology</i> , 2001 , 8, 337-49	2.3	30
263	Multiparametric magnetic resonance imaging: Current role in prostate cancer management. <i>International Journal of Urology</i> , 2016 , 23, 550-7	2.3	30
262	Inverse association between histologic inflammation in needle biopsy specimens and prostate cancer in men with serum PSA of 10-50 ng/mL. <i>Urology</i> , 2008 , 72, 1194-7	1.6	29
261	Enhanced expression of the secreted form of clusterin following neoadjuvant hormonal therapy as a prognostic predictor in patients undergoing radical prostatectomy for prostate cancer. <i>Oncology Reports</i> , 2005 , 14, 1371-5	3.5	29

260	Console-integrated real-time three-dimensional image overlay navigation for robot-assisted partial nephrectomy with selective arterial clamping: early single-centre experience with 17 cases. <i>International Journal of Medical Robotics and Computer Assisted Surgery</i> , 2014 , 10, 385-90	2.9	28
259	Chemosensitization of gemcitabine-resistant human bladder cancer cell line both in vitro and in vivo using antisense oligonucleotide targeting the anti-apoptotic gene, clusterin. <i>BJU International</i> , 2009 , 103, 384-90	5.6	28
258	Analysis of differences in clinicopathological features between prostate cancers located in the transition and peripheral zones. <i>International Journal of Urology</i> , 2006 , 13, 368-72	2.3	28
257	Comparative Assessment of Efficacies Between 2 Alternative Therapeutic Sequences With Novel Androgen Receptor-Axis-Targeted Agents in Patients With Chemotherapy-Naïve Metastatic Castration-Resistant Prostate Cancer. <i>Clinical Genitourinary Cancer</i> , 2017 , 15, e591-e597	3.3	27
256	Expression profile of epithelial-mesenchymal transition markers in non-muscle-invasive urothelial carcinoma of the bladder: correlation with intravesical recurrence following transurethral resection. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015 , 33, 110.e11-8	2.8	27
255	Assessment of efficacy, safety and quality of life of 110 patients treated with sunitinib as first-line therapy for metastatic renal cell carcinoma: experience in real-world clinical practice in Japan. <i>Medical Oncology</i> , 2014 , 31, 978	3.7	27
254	Prognostic prediction following radical prostatectomy for prostate cancer using conventional as well as molecular biological approaches. <i>International Journal of Urology</i> , 2013 , 20, 301-11	2.3	27
253	Significance of micrometastases in pelvic lymph nodes detected by real-time reverse transcriptase polymerase chain reaction in patients with clinically localized prostate cancer undergoing radical prostatectomy after neoadjuvant hormonal therapy. <i>BJU International</i> , 2007 , 99, 315-20	5.6	27
252	Comparative Assessment of Clinical Outcomes Between Abiraterone Acetate and Enzalutamide in Patients With Docetaxel-Naïve Metastatic Castration-Resistant Prostate Cancer: Experience in Real-World Clinical Practice in Japan. <i>Clinical Genitourinary Cancer</i> , 2017 , 15, 313-319	3.3	26
251	Long-term changes in renal function outcomes following radical cystectomy and urinary diversion. <i>International Journal of Clinical Oncology</i> , 2014 , 19, 1105-11	4.2	26
250	Expression of cell cycle-associated proteins in non-muscle-invasive bladder cancer: correlation with intravesical recurrence following transurethral resection. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2011 , 29, 495-501	2.8	26
249	Clinical outcome of combined immunotherapy with interferon-alpha and low-dose interleukine-2 for Japanese patients with metastatic renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2009 , 27, 598-603	2.8	26
248	Therapeutic efficacy of adenoviral-mediated p53 gene transfer is synergistically enhanced by combined use of antisense oligodeoxynucleotide targeting clusterin gene in a human bladder cancer model. <i>Neoplasia</i> , 2005 , 7, 171-9	6.4	26
247	Improved health-related quality of life of patients with metastatic renal cell carcinoma treated with a 2 weeks on and 1 week off schedule of sunitinib. <i>Medical Oncology</i> , 2015 , 32, 78	3.7	25
246	Serum level of clusterin and its density in men with prostate cancer as novel biomarkers reflecting disease extension. <i>Urology</i> , 2010 , 75, 454-9	1.6	25
245	Significance of micrometastases in prostate cancer. <i>Surgical Oncology</i> , 2008 , 17, 247-52	2.5	25
244	Significance of radical cystectomy for bladder cancer in patients over 80 years old. <i>International Urology and Nephrology</i> , 2007 , 39, 209-14	2.3	25
243	Enhanced expression of heat shock protein 27 following neoadjuvant hormonal therapy is associated with poor clinical outcome in patients undergoing radical prostatectomy for prostate cancer. <i>Anticancer Research</i> , 2006 , 26, 1583-7	2.3	25

242	Feasibility and usefulness of laparoscopic radical prostatectomy: Kobe University experience. <i>International Journal of Urology</i> , 2002 , 9, 635-40	2.3	24
241	Discovery of lipid biomarkers correlated with disease progression in clear cell renal cell carcinoma using desorption electrospray ionization imaging mass spectrometry. <i>Oncotarget</i> , 2019 , 10, 1688-1703	3.3	24
240	Expression of potential molecular markers in renal cell carcinoma: impact on clinicopathological outcomes in patients undergoing radical nephrectomy. <i>BJU International</i> , 2009 , 104, 942-6	5.6	23
239	Introduction of insulin-like growth factor binding protein-2 gene into human bladder cancer cells enhances their metastatic potential. <i>Oncology Reports</i> , 2005 , 13, 341-5	3.5	23
238	De Ritis (aspartate transaminase/alanine transaminase) ratio as a significant predictor of recurrence-free survival in patients with upper urinary tract urothelial carcinoma following nephroureterectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2016 , 34, 417.e9-417.e15	2.8	22
237	Surgical margin status of open versus laparoscopic radical prostatectomy specimens. <i>International Journal of Urology</i> , 2008 , 15, 704-7; discussion 708	2.3	22
236	Improved accuracy for predicting the Gleason score of prostate cancer by increasing the number of transrectal biopsy cores. <i>Urologia Internationalis</i> , 2007 , 79, 302-6	1.9	22
235	Synergistic inhibition of tumor growth and metastasis by combined treatment with TNP-470 and gemcitabine in a human bladder cancer KoTCC-1 model. <i>Journal of Urology</i> , 2004 , 172, 1485-9	2.5	22
234	Oncological outcome of docetaxel-based chemotherapy for Japanese men with metastatic castration-resistant prostate cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 733-8	2.8	21
233	Assessment of Efficacy, Safety, and Quality of Life of 124 Patients Treated With Axitinib as Second-Line Therapy for Metastatic Renal-Cell Carcinoma: Experience in Real-World Clinical Practice in Japan. <i>Clinical Genitourinary Cancer</i> , 2017 , 15, 122-128	3.3	21
232	Prognostic Factors Influencing Postoperative Development of Chronic Kidney Disease in Patients with Small Renal Tumors who Underwent Partial Nephrectomy. <i>Current Urology</i> , 2013 , 6, 129-35	1.7	21
231	The potential of clusterin inhibiting antisense oligodeoxynucleotide therapy for prostate cancer. <i>Expert Opinion on Investigational Drugs</i> , 2006 , 15, 507-17	5.9	21
230	Prediction of potentially insignificant prostate cancer in men undergoing radical prostatectomy for clinically organ-confined disease. <i>International Journal of Urology</i> , 2005 , 12, 270-4	2.3	21
229	Over expression of clusterin is an independent prognostic factor for nonpapillary renal cell carcinoma. <i>Journal of Urology</i> , 2002 , 167, 703-6	2.5	21
228	Overexpression of Bcl-2 regulates sodium butyrate- and/or docetaxel-induced apoptosis in human bladder cancer cells both in vitro and in vivo. <i>International Journal of Cancer</i> , 2001 , 93, 26-32	7.5	20
227	Avelumab plus axitinib vs sunitinib for advanced renal cell carcinoma: Japanese subgroup analysis from JAVELIN Renal 101. <i>Cancer Science</i> , 2020 , 111, 907-923	6.9	19
226	Modulation of the microenvironment by growth factors regulates the in vivo growth of skeletal myoblasts. <i>BJU International</i> , 2009 , 103, 1569-73	5.6	19
225	Significance of renal function in changes in acid-base metabolism after orthotopic bladder replacement: colon neobladder compared with ileal neobladder. <i>International Journal of Urology</i> , 2004 , 11, 83-7	2.3	19

224	Clinical outcome of conservative therapy for stage T1, grade 3 transitional cell carcinoma of the bladder. <i>International Journal of Urology</i> , 2003 , 10, 19-24	2.3	19
223	Early Tumor Shrinkage Under Treatment with First-line Tyrosine Kinase Inhibitors as a Predictor of Overall Survival in Patients with Metastatic Renal Cell Carcinoma: a Retrospective Multi-Institutional Study in Japan. <i>Targeted Oncology</i> , 2016 , 11, 175-82	5	18
222	Expression level of phosphorylated-4E-binding protein 1 in radical nephrectomy specimens as a prognostic predictor in patients with metastatic renal cell carcinoma treated with mammalian target of rapamycin inhibitors. <i>Medical Oncology</i> , 2014 , 31, 792	3.7	18
221	Expression pattern of testicular claudin-11 in infertile men. <i>Urology</i> , 2012 , 80, 1161.e13-7	1.6	18
220	Small interfering RNA targeting heat shock protein 70 enhances chemosensitivity in human bladder cancer cells. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 843-8	2.8	18
219	Retroperitoneoscopic nephroureterectomy for upper urinary tract cancer: a comparative study with conventional open retroperitoneal nephroureterectomy. <i>Journal of Endourology</i> , 2008 , 22, 1693-9	2.7	18
218	Significance of docetaxel-based chemotherapy as treatment for metastatic castration-resistant prostate cancer in Japanese men over 75 years old. <i>International Urology and Nephrology</i> , 2012 , 44, 1697-703	2.3	17
217	Analysis of factors predicting intravesical recurrence of superficial transitional cell carcinoma of the bladder without concomitant carcinoma in situ. <i>International Journal of Urology</i> , 2006 , 13, 1389-92	2.3	17
216	Improved survival in patients with metastatic renal cell carcinoma undergoing cytoreductive nephrectomy in the era of targeted therapy. <i>International Journal of Clinical Oncology</i> , 2014 , 19, 674-8	4.2	16
215	Prognostic prediction in patients with metastatic renal cell carcinoma treated with sorafenib based on expression levels of potential molecular markers in radical nephrectomy specimens. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 42-50	2.8	16
214	Expression profile of CD44s, CD44v6, and CD44v10 in localized prostate cancer: effect on prognostic outcomes following radical prostatectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 694-700	2.8	16
213	Inhibition of tumor growth and sensitization to chemotherapy by RNA interference targeting interleukin-6 in the androgen-independent human prostate cancer PC3 model. <i>Cancer Science</i> , 2011 , 102, 769-75	6.9	16
212	Significance of preoperatively observed detrusor overactivity as a predictor of continence status early after robot-assisted radical prostatectomy. <i>Asian Journal of Andrology</i> , 2014 , 16, 869-72	2.8	16
211	Prognostic Significance of Time to Castration Resistance in Patients With Metastatic Castration-sensitive Prostate Cancer. <i>Anticancer Research</i> , 2019 , 39, 1391-1396	2.3	15
210	Partial nephrectomy for hilar tumors: comparison of conventional open and robot-assisted approaches. <i>International Journal of Clinical Oncology</i> , 2015 , 20, 808-13	4.2	15
209	Expression of molecular markers associated with the mammalian target of rapamycin pathway in nonmetastatic renal cell carcinoma: Effect on prognostic outcomes following radical nephrectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 49.e15-21	2.8	15
208	Assessment of long-term quality of life in patients with orthotopic neobladder followed for more than 5 years. <i>International Urology and Nephrology</i> , 2011 , 43, 749-54	2.3	15
207	Clinical outcome of ureteroscopic lithotripsy for 2,129 patients with ureteral stones. <i>Urological Research</i> , 2007 , 35, 149-53		15

206	Papillary renal cell carcinoma: a clinicopathological study of 35 cases. <i>International Journal of Urology</i> , 2006 , 13, 1049-52	2.3	15
205	Expression pattern of autophagy-related markers in non-metastatic clear cell renal cell carcinoma: association with disease recurrence following radical nephrectomy. <i>Journal of Cancer Research and Clinical Oncology</i> , 2015 , 141, 1585-91	4.9	14
204	Adverse prognostic impact of capsular incision at radical prostatectomy for Japanese men with clinically localized prostate cancer. <i>International Urology and Nephrology</i> , 2009 , 41, 581-6	2.3	14
203	Improved sensitivity for detecting micrometastases in pelvic lymph nodes by real-time reverse transcriptase polymerase chain reaction (RT-PCR) compared with conventional RT-PCR in patients with clinically localized prostate cancer undergoing radical prostatectomy. <i>BJU International</i> , 2009 , 103, 1074-8	5.6	14
202	Health-related quality of life in patients undergoing lithotripsy for urinary stones. <i>International Urology and Nephrology</i> , 2008 , 40, 39-43	2.3	14
201	Hand-assisted laparoscopic radical nephrectomy: comparison with conventional open radical nephrectomy. <i>Journal of Endourology</i> , 2007 , 21, 429-32	2.7	14
200	Expression pattern of immune checkpoint-associated molecules in radical nephrectomy specimens as a prognosticator in patients with metastatic renal cell carcinoma treated with tyrosine kinase inhibitors. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2017 , 35, 363-369	2.8	13
199	Prospective assessment of time-dependent changes in quality of life of Japanese patients with prostate cancer following robot-assisted radical prostatectomy. <i>Journal of Robotic Surgery</i> , 2016 , 10, 201-7	2.9	13
198	Significance of erection hardness score as a diagnostic tool to assess erectile function recovery in Japanese men after robot-assisted radical prostatectomy. <i>Journal of Robotic Surgery</i> , 2016 , 10, 221-6	2.9	13
197	Significance of circulating matrix metalloproteinase-9 to tissue inhibitor of metalloproteinases-2 ratio as a predictor of disease progression in patients with metastatic renal cell carcinoma receiving sunitinib. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 584-8	2.8	13
196	Clinical outcome of microsurgical varicocelectomy in infertile men with severe oligozoospermia. <i>Urology</i> , 2014 , 83, 1071-4	1.6	13
195	Impact of sorafenib on health-related quality of life in Japanese patients with metastatic renal cell carcinoma: a prospective evaluation. <i>BJU International</i> , 2010 , 106, 1643-7	5.6	13
194	Long-term functional outcomes in patients with various types of orthotopic intestinal neobladder. <i>International Journal of Urology</i> , 2008 , 15, 612-5	2.3	13
193	Persistent expression of Aurora-A after neoadjuvant hormonal therapy as a predictor of a poor clinical outcome in patients undergoing radical prostatectomy for prostate cancer. <i>BJU International</i> , 2007 , 100, 310-4	5.6	13
192	Significance of routine transition zone biopsies in Japanese men undergoing transrectal ultrasound-guided prostate biopsies. <i>International Journal of Urology</i> , 2005 , 12, 964-8	2.3	13
191	Significant impact of R.E.N.A.L. nephrometry score on changes in postoperative renal function early after robot-assisted partial nephrectomy. <i>International Journal of Clinical Oncology</i> , 2015 , 20, 586-92	4.2	12
190	Assessment of Time-dependent Changes in Semen Parameters in Infertile Men After Microsurgical Varicocelectomy. <i>Urology</i> , 2015 , 86, 48-51	1.6	12
189	Significance of 4E-binding protein 1 as a therapeutic target for invasive urothelial carcinoma of the bladder. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015 , 33, 166.e9-15	2.8	12

188	Efficacy and safety of temsirolimus in Japanese patients with metastatic renal cell carcinoma on hemodialysis. <i>International Journal of Clinical Oncology</i> , 2013 , 18, 1054-9	4.2	12
187	Expression of integrin proteins in non-muscle-invasive bladder cancer: significance of intravesical recurrence after transurethral resection. <i>BJU International</i> , 2011 , 107, 240-6	5.6	12
186	Orthotopic sigmoid neobladder after radical cystectomy: assessment of complications, functional outcomes and quality of life in 82 Japanese patients. <i>BJU International</i> , 2010 , 106, 412-6	5.6	12
185	Characterization of prostate cancer incidentally detected in radical cystoprostatectomy specimens from Japanese men with bladder cancer. <i>International Urology and Nephrology</i> , 2010 , 42, 73-9	2.3	12
184	Additional sampling of dorsal apex on systematic prostate biopsy: impact on early detection of prostate cancer. <i>Urology</i> , 2007 , 69, 738-42	1.6	12
183	Clinical symptoms in localized renal cell carcinoma reflect its invasive potential: comparative study between incidentally detected and symptomatic diseases. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2006 , 24, 201-6	2.8	12
182	Neoadjuvant androgen withdrawal prior to external radiotherapy for locally advanced adenocarcinoma of the prostate. <i>International Journal of Urology</i> , 2002 , 9, 322-8; discussion 328	2.3	12
181	Serum cathepsin D and its density in men with prostate cancer as new predictors of disease progression. <i>Oncology Reports</i> , 2002 , 9, 1379-83	3.5	12
180	Enhanced radiosensitivity by inhibition of the anti-apoptotic gene clusterin using antisense oligodeoxynucleotide in a human bladder cancer model. <i>Oncology Reports</i> , 2005 , 13, 885-90	3.5	12
179	Comparison of Alternative Androgen Receptor-axis-targeted Agent (ARATA) and Docetaxel as Second-line Therapy for Patients With Metastatic Castration-resistant Prostate Cancer With Progression After Initial ARATA in Real-world Clinical Practice in Japan. <i>Clinical Genitourinary Cancer</i> , 2018 , 16, 210-225	3.3	11
178	Prognostic outcome in patients treated with tyrosine kinase inhibitors as first-line molecular-targeted therapy for metastatic renal cell carcinoma: Experience in real-world clinical practice in Japan. <i>Molecular and Clinical Oncology</i> , 2015 , 3, 601-606	1.6	11
177	Salvage chemotherapy with methotrexate, etoposide and actinomycin D in men with metastatic nonseminomatous germ cell tumors with a choriocarcinoma component: a preliminary report. <i>International Journal of Urology</i> , 2010 , 17, 881-5	2.3	11
176	Oncological outcome of laparoscopic prostatectomy. <i>International Journal of Urology</i> , 2007 , 14, 515-20	2.3	11
175	Limited significance of routine excretory urography in the follow-up of patients with superficial bladder cancer after transurethral resection. <i>BJU International</i> , 2006 , 97, 720-3	5.6	11
174	Long-term results of adjuvant hormonal therapy plus radiotherapy following radical prostatectomy for patients with pT3N0 or pT3N1 prostate cancer. <i>International Journal of Urology</i> , 2004 , 11, 397-401	2.3	11
173	Limited usefulness of the free-to-total prostate-specific antigen ratio for the diagnosis and staging of prostate cancer in Japanese men. <i>International Journal of Clinical Oncology</i> , 2004 , 9, 64-7	4.2	11
172	Complete response to combination therapy with nivolumab and ipilimumab for metastatic collecting duct carcinoma of the kidney. <i>International Cancer Conference Journal</i> , 2020 , 9, 32-35	0.9	11
171	Synergistic antitumor effect of combined use of adenoviral-mediated p53 gene transfer and antisense oligodeoxynucleotide targeting clusterin gene in an androgen-independent human prostate cancer model. <i>Molecular Cancer Therapeutics</i> , 2005 , 4, 187-95	6.1	11

170	Development of a Complete En-Bloc Technique with Direct Bladder Neck Incision: A Newly Modified Approach for Holmium Laser Enucleation of the Prostate. <i>Journal of Endourology</i> , 2019 , 33, 835-840	2.7	10
169	A multicenter retrospective study of nivolumab monotherapy in previously treated metastatic renal cell carcinoma patients: interim analysis of Japanese real-world data. <i>International Journal of Clinical Oncology</i> , 2020 , 25, 1533-1542	4.2	10
168	Clinical outcomes of anti-androgen withdrawal and subsequent alternative anti-androgen therapy for advanced prostate cancer following failure of initial maximum androgen blockade. <i>Molecular and Clinical Oncology</i> , 2016 , 4, 839-844	1.6	10
167	Third-line sunitinib following sequential use of cytokine therapy and sorafenib in Japanese patients with metastatic renal cell carcinoma. <i>International Journal of Clinical Oncology</i> , 2013 , 18, 81-6	4.2	10
166	Prediction of response to combined immunotherapy with interferon- γ and low-dose interleukin-2 in metastatic renal cell carcinoma: Expression patterns of potential molecular markers in radical nephrectomy specimens. <i>International Journal of Urology</i> , 2009 , 16, 465-471	2.3	10
165	Suppressed tumour growth and enhanced chemosensitivity by RNA interference targeting Aurora-A in the PC3 human prostate cancer model. <i>BJU International</i> , 2010 , 106, 121-7	5.6	10
164	Combination chemotherapy with weekly docetaxel and estramustine for hormone refractory prostate cancer in Japanese patients. <i>International Journal of Urology</i> , 2008 , 15, 106-9	2.3	10
163	Prediction of the extent of prostate cancer by the combined use of systematic biopsy and serum level of cathepsin D. <i>International Journal of Urology</i> , 2003 , 10, 196-200	2.3	10
162	Clinical outcome of high-dose chemotherapy combined with peripheral blood stem cell transplantation for male germ cell tumors. <i>Anti-Cancer Drugs</i> , 1999 , 10, 711-8	2.4	10
161	Hyponatremia as a powerful prognostic predictor for Japanese patients with clear cell renal cell carcinoma treated with a tyrosine kinase inhibitor. <i>International Journal of Clinical Oncology</i> , 2015 , 20, 351-7	4.2	9
160	Impact of Prior Use of an Androgen Receptor-Axis-Targeted (ARAT) Agent With or Without Subsequent Taxane Therapy on the Efficacy of Another ARAT Agent in Patients With Metastatic Castration-Resistant Prostate Cancer. <i>Clinical Genitourinary Cancer</i> , 2017 , 15, e217-e222	3.3	9
159	Enhanced sensitivity to sorafenib by inhibition of Akt1 expression in human renal cell carcinoma ACHN cells both in vitro and in vivo. <i>Human Cell</i> , 2015 , 28, 114-21	4.5	9
158	The antiandrogen bicalutamide activates the androgen receptor (AR) with a mutation in codon 741 through the mitogen activated protein kinase (MARK) pathway in human prostate cancer PC3 cells. <i>Oncology Reports</i> , 2010 , 24, 1395-9	3.5	9
157	Marked response to nivolumab combined with external radiation therapy for metastatic renal cell carcinoma: report of two cases. <i>International Cancer Conference Journal</i> , 2019 , 8, 29-32	0.9	9
156	Health-related quality of life in patients with testicular cancer: a comparative analysis according to therapeutic modalities. <i>Oncology Reports</i> , 2004 , 12, 867-70	3.5	9
155	Expression Profile of Autophagy-related Markers in Localized Prostate Cancer: Correlation With Biochemical Recurrence After Radical Prostatectomy. <i>Urology</i> , 2015 , 85, 1424-30	1.6	8
154	Risk for intravesical recurrence of bladder cancer stratified by the results on two consecutive UroVysion fluorescence in situ hybridization tests: a prospective follow-up study in Japan. <i>International Journal of Clinical Oncology</i> , 2020 , 25, 1163-1169	4.2	8
153	Renal Functional and Perioperative Outcomes of Selective Versus Complete Renal Arterial Clamping During Robot-Assisted Partial Nephrectomy: Early Single-Center Experience With 39 Cases. <i>Surgical Innovation</i> , 2016 , 23, 242-8	2	8

152	Significance of De Ritis (Aspartate Transaminase/Alanine Transaminase) Ratio as a Significant Prognostic But Not Predictive Biomarker in Japanese Patients with Metastatic Castration-resistant Prostate Cancer Treated with Cabazitaxel. <i>Anticancer Research</i> , 2018 , 38, 4179-4185	2.3	8
151	Orthotopic sigmoid vs. ileal neobladders in Japanese patients: a comparative assessment of complications, functional outcomes, and quality of life. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 1155-60	2.8	8
150	Prognostic significance of primary Gleason pattern in Japanese men with Gleason score 7 prostate cancer treated with radical prostatectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 1511-6	2.8	8
149	Orthotopic bladder substitution following radical cystectomy in women: comparative study between sigmoid and ileal neobladders. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2012 , 30, 38-43	2.8	8
148	Age at diagnosis as a powerful predictor for disease recurrence after radical nephrectomy in Japanese patients with pT1 renal cell carcinoma. <i>International Journal of Urology</i> , 2011 , 18, 121-5	2.3	8
147	Impact of body mass index on perioperative outcomes of laparoscopic radical nephrectomy in Japanese patients with clinically localized renal cell carcinoma. <i>International Journal of Urology</i> , 2010 , 17, 522-6	2.3	8
146	Assessment of voiding function of orthotopic neobladders in elderly patients with long-term survival. <i>BJU International</i> , 2009 , 103, 927-30	5.6	8
145	Clinical outcome of radical cystectomy for patients with pT4 bladder cancer. <i>International Journal of Urology</i> , 2008 , 15, 58-61	2.3	8
144	First-line high-dose chemotherapy combined with peripheral blood stem cell transplantation for patients with advanced extragonadal germ cell tumors. <i>International Journal of Urology</i> , 2007 , 14, 336-8	2.3	8
143	1088: Novel Bispecific Antisense Oligonucleotides Inhibiting both BCL-2 and BCL-XL Expression Induce Apoptosis and Enhance Chemosensitivity in Human Androgen-Independent Prostate Cancer Cells. <i>Journal of Urology</i> , 2004 , 171, 287-287	2.5	8
142	Clinical outcome of transrectal ultrasound-guided prostate biopsy, targeting eight cores, for detecting prostate cancer in Japanese men. <i>International Journal of Clinical Oncology</i> , 2004 , 9, 47-50	4.2	8
141	Prognostic significance of insulin-like growth factor (IGF) binding protein-2 to IGF-binding protein-3 ratio in patients undergoing radical cystectomy for invasive transitional cell carcinoma of the bladder. <i>BJU International</i> , 2005 , 95, 987-91	5.6	8
140	Non-significant impact of proteinuria on renal function in Japanese patients with metastatic renal cell carcinoma treated with axitinib. <i>International Journal of Clinical Oncology</i> , 2015 , 20, 796-801	4.2	7
139	Significance of multiple preoperative laboratory abnormalities as prognostic indicators in patients with urothelial carcinoma of the upper urinary tract following radical nephroureterectomy. <i>International Journal of Clinical Oncology</i> , 2018 , 23, 151-157	4.2	7
138	Prognostic Significance of Early Tumor Shrinkage Under Second-Line Targeted Therapy for Metastatic Renal Cell Carcinoma: A Retrospective Multi-Institutional Study in Japan. <i>Molecular Diagnosis and Therapy</i> , 2016 , 20, 385-92	4.5	7
137	Changes in Renal Function of Patients with Metastatic Renal Cell Carcinoma During Treatment with Molecular-Targeted Agents. <i>Targeted Oncology</i> , 2016 , 11, 329-35	5	7
136	Sequential use of mammalian target of rapamycin inhibitors in patients with metastatic renal cell carcinoma following failure of tyrosine kinase inhibitors. <i>Medical Oncology</i> , 2013 , 30, 745	3.7	7
135	No significant impact of prior treatment profile with docetaxel on the efficacy of cabazitaxel in Japanese patients with metastatic castration-resistant prostate cancer. <i>Medical Oncology</i> , 2017 , 34, 141	3.7	7

134	Growth inhibition and enhanced chemosensitivity induced by down-regulation of Aurora-A in human renal cell carcinoma Caki-2 cells using short hairpin RNA. <i>Oncology Letters</i> , 2011 , 2, 713-717	2.6	7
133	Increased detection of clinically significant prostate cancer by additional sampling from the anterior lateral horns of the peripheral zone in combination with the standard sextant biopsy. <i>International Journal of Urology</i> , 2004 , 11, 402-6	2.3	7
132	Clinicopathological features of recurrence after radical surgery for nonmetastatic renal cell carcinoma. <i>International Journal of Clinical Oncology</i> , 2004 , 9, 369-72	4.2	7
131	Antisense oligodeoxynucleotide therapy for bladder cancer: recent advances and future prospects. <i>Expert Review of Anticancer Therapy</i> , 2005 , 5, 1001-9	3.5	7
130	Impact of penile rehabilitation with low-dose vardenafil on recovery of erectile function in Japanese men following nerve-sparing radical prostatectomy. <i>Asian Journal of Andrology</i> , 2014 , 16, 892-6	2.8	7
129	GLI2 expression levels in radical nephrectomy specimens as a predictor of disease progression in patients with metastatic clear cell renal cell carcinoma following treatment with sunitinib. <i>Molecular and Clinical Oncology</i> , 2016 , 5, 186-192	1.6	7
128	Urinary levels of vascular endothelial growth factor in patients with prostate cancer as a predictor of disease progression. <i>Anticancer Research</i> , 2005 , 25, 3645-9	2.3	7
127	Comparative assessment of prognostic outcomes between first-generation antiandrogens and novel androgen-receptor-axis-targeted agents in patients with non-metastatic castration-resistant prostate cancer. <i>International Journal of Clinical Oncology</i> , 2019 , 24, 842-847	4.2	6
126	Oncological outcomes after cytoreductive nephrectomy for patients with metastatic renal cell carcinoma with inferior vena caval tumor thrombus. <i>International Journal of Clinical Oncology</i> , 2018 , 23, 553-558	4.2	6
125	Efficacy and safety of axitinib in elderly patients with metastatic renal cell carcinoma. <i>Medical Oncology</i> , 2016 , 33, 95	3.7	6
124	Objective assessment of residual nerve tissues in radical prostatectomy specimens by immunohistochemical staining of neuronal nitric oxide synthase-positive nerves and its impact on postoperative erectile function. <i>Urology</i> , 2014 , 84, 1395-401	1.6	6
123	Prediction of response to combined immunotherapy with interferon-alpha and low-dose interleukin-2 in metastatic renal cell carcinoma: expression patterns of potential molecular markers in radical nephrectomy specimens. <i>International Journal of Urology</i> , 2009 , 16, 465-71	2.3	6
122	Prognostic significance of adjuvant cisplatin-based combination chemotherapy following radical cystectomy in patients with invasive bladder cancer. <i>International Journal of Urology</i> , 2008 , 15, 314-8	2.3	6
121	Clinical outcome of orthotopic neobladder replacement in patients with a solitary functioning kidney. <i>International Journal of Urology</i> , 2007 , 14, 398-401	2.3	6
120	Health-related quality of life after chemotherapy for advanced germ cell tumors: a comparison of standard-dose and high-dose chemotherapy. <i>International Journal of Urology</i> , 2004 , 11, 542-6	2.3	6
119	An evaluation of quality of life in patients who underwent orthotopic bladder replacement after cystectomy: comparison of ileal neobladder versus colon neobladder. <i>Urologia Internationalis</i> , 2002 , 69, 195-9	1.9	6
118	Characterization of urocortin as an anti-apoptotic protein in experimental ischemia-reperfusion model of the rat testis. <i>Biochemical and Biophysical Research Communications</i> , 2016 , 479, 387-392	3.4	6
117	Feasibility of the ACL (albumin, C-reactive protein and lactate dehydrogenase) model as a novel prognostic tool in patients with metastatic renal cell carcinoma previously receiving first-line targeted therapy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2020 , 38, 6.e9-6.e16	2.8	6

116	Clinical outcome of oral uracil/tegafur (UFT) therapy for patients with hormone refractory prostate cancer. <i>Oncology Reports</i> , 2005 , 14, 673-6	3.5	6
115	No Significant Correlation of Clinical Outcomes Between First- and Second-line Tyrosine Kinase Inhibitors in Patients with Metastatic Renal Cell Carcinoma. <i>Anticancer Research</i> , 2015 , 35, 3067-73	2.3	6
114	Value of three-dimensional T2-weighted turbo spin-echo imaging with tissue-specific variable refocusing flip angle for 3-T magnetic resonance imaging of prostate cancer: comparison with conventional two- and three-dimensional T2-weighted turbo spin-echo imaging. <i>Japanese Journal of Radiology</i> , 2017 , 35, 707-717	2.9	5
113	Enhanced Sensitivity to Sunitinib by Inhibition of Akt1 Expression in Human Castration-resistant Prostate Cancer PC3 Cells Both In Vitro and In Vivo. <i>Urology</i> , 2015 , 85, 1215.e1-1215.e7	1.6	5
112	Significance of introduction of alternative dosing schedule for sunitinib during first-line treatment of patients with metastatic renal cell carcinoma. <i>Medical Oncology</i> , 2018 , 35, 133	3.7	5
111	Assessment of optimal target genes for detecting micrometastases in pelvic lymph nodes in patients with prostate cancer undergoing radical prostatectomy by real-time reverse transcriptase-polymerase chain reaction. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2018 , 36, 615-621	2.8	5
110	Health-related quality of life in Japanese patients with metastatic renal cell carcinoma treated with sunitinib. <i>International Journal of Clinical Oncology</i> , 2013 , 18, 220-5	4.2	5
109	Elevated urinary cytokine levels in patients undergoing ileal neobladder replacement compared with sigmoid neobladder replacement. <i>International Urology and Nephrology</i> , 2007 , 39, 467-72	2.3	5
108	Screening for prostate cancer using prostate-specific antigen testing in Japanese men on hemodialysis. <i>International Urology and Nephrology</i> , 2008 , 40, 345-9	2.3	5
107	Clinical outcome of combined immunotherapy with low-dose interleukin-2 and interferon-alpha for Japanese patients with metastatic renal cell carcinoma who had undergone radical nephrectomy: a preliminary report. <i>International Journal of Clinical Oncology</i> , 2005 , 10, 338-41	4.2	5
106	Prognostic significance of the tumor volume in radical prostatectomy specimens after neoadjuvant hormonal therapy. <i>Urologia Internationalis</i> , 2005 , 74, 27-31	1.9	5
105	Differential involvement of the Fas receptor/ligand system in p53-dependent apoptosis in human prostate cancer cells. <i>Prostate</i> , 2000 , 45, 341-9	4.2	5
104	Significance of urethral fibrosis evaluated by preoperative magnetic resonance imaging as a predictor of continence status after robot-assisted radical prostatectomy. <i>International Journal of Medical Robotics and Computer Assisted Surgery</i> , 2016 , 12, 496-501	2.9	5
103	Initial learning curve for robot-assisted partial nephrectomy performed by a single experienced robotic surgeon. <i>Asian Journal of Endoscopic Surgery</i> , 2020 , 13, 59-64	1.4	5
102	Irinotecan and nedaplatin as salvage therapy for patients with advanced germ cell tumors following intensive treatment with cisplatin-based combination chemotherapies. <i>International Journal of Clinical Oncology</i> , 2016 , 21, 162-7	4.2	4
101	Independent association between time to prostate-specific antigen (PSA) nadir and PSA progression-free survival in patients with docetaxel-naïve, metastatic castration-resistant prostate cancer receiving abiraterone acetate, but not enzalutamide. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2017 , 35, 432-437	2.8	4
100	Contralateral pneumothorax during retroperitoneal laparoscopic donor nephrectomy: A case report. <i>Asian Journal of Endoscopic Surgery</i> , 2017 , 10, 202-204	1.4	4
99	Early Single-Center Experience with Robotic Partial Nephrectomy Using the da Vinci Xi: Comparative Assessment with Conventional Open Partial Nephrectomy. <i>Current Urology</i> , 2019 , 13, 13-18	1.7	4

98	Preoperative differentiation between benign and malignant renal masses smaller than 4 cm treated with partial nephrectomy. <i>International Journal of Clinical Oncology</i> , 2015 , 20, 150-5	4.2	4
97	Assessment of health-related quality of life in Japanese patients with metastatic renal cell carcinoma during treatment with tyrosine kinase inhibitors. <i>Medical Oncology</i> , 2014 , 31, 190	3.7	4
96	Assessment of efficacy, safety and quality of life of 55 patients with metastatic renal cell carcinoma treated with temsirolimus: a single-center experience in Japan. <i>International Journal of Clinical Oncology</i> , 2014 , 19, 679-85	4.2	4
95	Assessment of postoperative quality of life: comparative study between laparoscopic and minimum incision endoscopic radical prostatectomies. <i>International Journal of Clinical Oncology</i> , 2014 , 19, 1092-7	4.2	4
94	Characterization of the anatomical extension pattern of localized prostate cancer arising in the peripheral zone. <i>BJU International</i> , 2010 , 105, 1514-8	5.6	4
93	Clinicopathological features of recurrence after radical cystectomy for patients with transitional cell carcinoma of the bladder. <i>International Urology and Nephrology</i> , 2006 , 38, 49-55	2.3	4
92	Increased serum carcinoembryonic antigen level in patients undergoing colon neobladder replacement compared with ileal neobladder replacement. <i>Urology</i> , 2002 , 60, 363-7	1.6	4
91	Significant impact of three-dimensional volumetry of perinephric fat on the console time during robot-assisted partial nephrectomy. <i>BMC Urology</i> , 2019 , 19, 132	2.2	4
90	Improved perioperative outcomes by early unclamping prior to renorrhaphy compared with conventional clamping during robot-assisted partial nephrectomy: a propensity score matching analysis. <i>Journal of Robotic Surgery</i> , 2020 , 14, 47-53	2.9	4
89	Absence of Significant Correlation of Adverse Events Between First- and Second-Line Tyrosine Kinase Inhibitors in Patients With Metastatic Renal Cell Carcinoma. <i>Clinical Genitourinary Cancer</i> , 2016 , 14, e19-24	3.3	3
88	Postoperative bacteriuria, pyuria and urinary tract infection in patients with an orthotopic sigmoid colon neobladder replacement. <i>Journal of Antibiotics</i> , 2014 , 67, 143-5	3.7	3
87	Efficacy and tolerability of TIP (paclitaxel, ifosfamide and cisplatin) incorporated into induction chemotherapy for patients with intermediate- or poor-risk metastatic germ cell tumors. <i>Medical Oncology</i> , 2014 , 31, 296	3.7	3
86	Significant impact of age at diagnosis on the prognosis of Japanese patients with pT1 renal cell carcinoma following surgical resection. <i>BJU International</i> , 2012 , 109, 695-9	5.6	3
85	UNC-51-like kinase 1 expression in radical nephrectomy specimens as a predicting factor of progression-free survival in patients with metastatic renal cell carcinoma treated with mammalian target of rapamycin inhibitors. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015 , 33, 506.e1-7	2.8	3
84	Downregulation of lectin-like oxidized low-density lipoprotein receptor-1 after ischemic preconditioning in ischemia-reperfused rat kidneys. <i>Urology</i> , 2009 , 73, 906-10	1.6	3
83	A comparison of clinical features between chlamydial and non-chlamydial urethritis in men negative for gonococcal infection who attended a urological outpatient clinic in Japan. <i>International Urology and Nephrology</i> , 2007 , 39, 809-13	2.3	3
82	The limited significance of a longer duration of neoadjuvant hormonal therapy prior to radical prostatectomy for high-risk prostate cancer in Japanese men. <i>Urologia Internationalis</i> , 2006 , 77, 122-6	1.9	3
81	Pathological findings of radical prostatectomy specimens in Japanese men diagnosed on single core positive prostate biopsy in eight with a Gleason score less than 4. <i>International Journal of Urology</i> , 2003 , 10, 383-6	2.3	3

80	Is a limited lymphadenectomy targeting obturator nodes alone an adequate procedure for Japanese men undergoing radical prostatectomy?. <i>International Journal of Urology</i> , 2005 , 12, 739-44	2.3	3
79	Clinicopathological features of prostate cancer in Japanese men diagnosed on repeat transrectal ultrasound-guided biopsy. <i>International Journal of Clinical Oncology</i> , 2005 , 10, 30-4	4.2	3
78	Predicting the extent of prostate cancer using a combination of serum prostate-specific antigen-alpha(1)-antichymotrypsin complex and systematic biopsy. <i>Urologia Internationalis</i> , 2002 , 68, 232-6	1.9	3
77	Impact of Periurethral Inflammation on Continence Status Early After Robot-Assisted Radical Prostatectomy. <i>Journal of Endourology</i> , 2016 , 30, 1207-1213	2.7	3
76	Development of novel ACN (albumin, C-reactive protein and neutrophil-to-lymphocyte ratio) prognostication model for patients with metastatic renal cell carcinoma receiving first-line molecular-targeted therapy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2021 , 39, 78.e1-78.e8	2.8	3
75	Efficacy and safety of avelumab plus axitinib (A + Ax) versus sunitinib (S) in elderly patients with advanced renal cell carcinoma (aRCC): Extended follow-up results from JAVELIN Renal 101.. <i>Journal of Clinical Oncology</i> , 2021 , 39, 301-301	2.2	3
74	Role of the serotonergic system in urethral continence reflexes during sneezing in rats. <i>American Journal of Physiology - Renal Physiology</i> , 2018 , 315, F79-F85	4.3	3
73	Significance of Age in Japanese Patients Receiving Sunitinib as First-line Systemic Therapy for Metastatic Renal Cell Carcinoma: Comparative Assessment of Efficacy and Safety between Patients Aged . <i>Anticancer Research</i> , 2018 , 38, 3593-3599	2.3	3
72	Clinical Outcomes of 42 Renal Cell Carcinoma Patients With Metastases Solely to the Lung Who Received Sorafenib as Second-line Systemic Therapy. <i>Anticancer Research</i> , 2019 , 39, 1067-1072	2.3	2
71	Optimal Timing of Cabazitaxel Introduction for Japanese Patients With Metastatic Castration-resistant Prostate Cancer. <i>Anticancer Research</i> , 2019 , 39, 3089-3094	2.3	2
70	Promising response to axitinib rechallenge for metastatic renal cell carcinoma after progression on prior axitinib: report of two cases. <i>International Cancer Conference Journal</i> , 2018 , 7, 30-34	0.9	2
69	No significant impact of response to prior androgen receptor-axis-targeted agents on the efficacy of subsequent docetaxel in patients with metastatic castration-resistant prostate cancer. <i>International Journal of Clinical Oncology</i> , 2018 , 23, 576-583	4.2	2
68	Erectile function and its impact on quality of life in Japanese men on hemodialysis. <i>Cogent Medicine</i> , 2016 , 3, 1211594	1.4	2
67	Prostate-specific antigen response patterns during cabazitaxel therapy in patients with metastatic castration-resistant prostate cancer. <i>Japanese Journal of Clinical Oncology</i> , 2019 , 49, 1043-1048	2.8	2
66	Inhibition of Tumor Growth and Sensitization to Sunitinib by RNA Interference Targeting Programmed Death-ligand 1 in Mouse Renal Cell Carcinoma RenCa Model. <i>Anticancer Research</i> , 2019 , 39, 4737-4742	2.3	2
65	Impact of prior androgen receptor-axis-targeted agents on the clinical activity of subsequent docetaxel in patients with metastatic castration-resistant prostate cancer: comparative assessment between abiraterone acetate and enzalutamide. <i>Medical Oncology</i> , 2017 , 34, 200	3.7	2
64	Phase II study of palonosetron, aprepitant and dexamethasone to prevent nausea and vomiting induced by multiple-day emetogenic chemotherapy. <i>Supportive Care in Cancer</i> , 2018 , 26, 1419-1423	3.9	2
63	Sigmoid neobladder as an ideal form of orthotopic urinary reconstruction. <i>International Journal of Urology</i> , 2012 , 19, 184-6	2.3	2

62	Long-term complete response to very-low-dose interleukin-2 therapy in patients with metastatic renal cell carcinoma: report of two cases. <i>Clinical and Experimental Nephrology</i> , 2011 , 15, 966-9	2.5	2
61	Experience with various types of orthotopic neobladder in Japanese men: long-term follow-up. <i>Urologia Internationalis</i> , 2010 , 84, 34-9	1.9	2
60	Predictors of Cancer-Specific Survival following Radical Cystectomy in Patients with Node-Positive Bladder Cancer. <i>Current Urology</i> , 2010 , 4, 188-192	1.7	2
59	Clinicopathological features of patients with biochemical recurrence after radical prostatectomy without progressive rise in serum prostate-specific antigen. <i>Urologia Internationalis</i> , 2006 , 77, 200-4	1.9	2
58	Enhanced expression of the secreted form of clusterin following neoadjuvant hormonal therapy as a prognostic predictor in patients undergoing radical prostatectomy for prostate cancer. <i>Oncology Reports</i> , 2005 , 14, 1371	3.5	2
57	Predictive factors for persistence of preoperative overactive bladder symptoms after transvaginal mesh surgery in women with pelvic organ prolapse. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2020 , 12, 167-172	1.9	2
56	Enhanced Sensitivity to NVP-BEZ235 by Inhibition of p62/SQSTM1 in Human Bladder Cancer KoTCC-1 Cells Both <i>In Vivo</i> and <i>In Vitro</i> . <i>In Vivo</i> , 2020 , 34, 1001-1008	2.3	2
55	Prognostic significance of third-line treatment for patients with metastatic castration-resistant prostate cancer: comparative assessments between cabazitaxel and other agents. <i>International Journal of Clinical Oncology</i> , 2021 , 26, 1745-1751	4.2	2
54	The Intrarenal Renin-angiotensin System Is Activated Immediately after Kidney Donation in Kidney Transplant Donors. <i>Internal Medicine</i> , 2019 , 58, 643-648	1.1	2
53	Acceleration of Proteinuria without Significant Impact on Renal Function and Its Protection by Angiotensin II Receptor Blocker in Rats Treated with Axitinib. <i>Targeted Oncology</i> , 2016 , 11, 309-15	5	1
52	Detection of Significant Prostate Cancer According to Anatomical Areas of Sampling Cores Obtained with Transrectal Systematic 12-Core Biopsy. <i>Current Urology</i> , 2015 , 8, 91-5	1.7	1
51	Assessment of Factors Predicting Disease Progression in Japanese Patients With Non-Metastatic Castration-resistant Prostate Cancer. <i>Anticancer Research</i> , 2020 , 40, 1101-1106	2.3	1
50	Second transurethral resection for high-risk non-muscle invasive bladder cancer patients: a propensity score matched analysis. <i>Japanese Journal of Clinical Oncology</i> , 2020 , 50, 718-721	2.8	1
49	Isosamidin, an extract of <i>Peucedanum japonicum</i> , inhibits phenylephrine-mediated contractions of the human prostate in vitro. <i>Phytotherapy Research</i> , 2018 , 32, 949-952	6.7	1
48	Clinical characteristics of self-reported nocturia in patients with interstitial cystitis, and effects of bladder hydrodistention (with fulguration of Hunner lesions) on nocturia. <i>LUTS: Lower Urinary Tract Symptoms</i> , 2019 , 11, O141-O146	1.9	1
47	Effect of Targeting Clusterin Using OGX-011 on Antitumor Activity of Temsirolimus in a Human Renal Cell Carcinoma Model. <i>Targeted Oncology</i> , 2017 , 12, 69-79	5	1
46	The Inverse Association between the Baseline Renal Function and Overall Survival in Patients with Metastatic Renal Cell Carcinoma Treated with Molecular-Targeted Agents. <i>Current Urology</i> , 2017 , 10, 186-192	1.7	1
45	Significance of uracil/tegafur for preventing intravesical recurrence of non-muscle invasive urothelial carcinoma of the bladder. <i>Current Urology</i> , 2012 , 6, 27-32	1.7	1

44	Prognostic Significance of Capsular Invasion and Capsular Penetration in Japanese Men with Prostate Cancer Undergoing Radical Retropubic Prostatectomy. <i>UroOncology</i> , 2004 , 4, 27-30		1
43	Long-term Outcome of Adjuvant Chemotherapy with MVP-CAB Regimen (Methotrexate, Vincristine, Cisplatin, Cyclophosphamide, Adriamycin and Bleomycin) for Locally Advanced Bladder Cancer. <i>UroOncology</i> , 2004 , 4, 151-156		1
42	Clinicopathological Features of Incidentally Detected Renal Cell Carcinoma in Japan. <i>UroOncology</i> , 2003 , 3, 95-98		1
41	Preoperative prediction of final pathological features is not improved by the free-to-total prostate-specific antigen ratio in Japanese men with clinically localized prostate cancer. <i>International Journal of Urology</i> , 2005 , 12, 182-6	2.3	1
40	Significance of the percentage of prostate needle biopsy cores with cancer as a predictor of disease extension in radical prostatectomy specimens in Japanese men. <i>International Urology and Nephrology</i> , 2005 , 37, 305-10	2.3	1
39	Significance of 5-Aminolevulinic Acid-mediated Photodynamic Diagnosis Following Standard Transurethral Resection in Non-muscle Invasive Bladder Cancer.. <i>Cancer Diagnosis & Prognosis</i> , 2021 , 1, 201-205		1
38	Comparison of clinical outcomes between androgen deprivation therapy with up-front abiraterone and bicalutamide for Japanese patients with LATITUDE high-risk prostate cancer in a real-world retrospective analysis. <i>International Journal of Clinical Oncology</i> , 2021 , 1	4.2	1
37	Assessment of prognostic factors in previously treated Japanese patients with metastatic renal cell carcinoma who received nivolumab: An observational multi-institute study. <i>Molecular and Clinical Oncology</i> , 2022 , 16, 17	1.6	1
36	Comprehensive assessment of oncological outcomes in 186 patients with high-risk non-muscle-invasive bladder cancer: A single institution retrospective study. <i>Molecular and Clinical Oncology</i> , 2020 , 13, 16	1.6	1
35	Comparison of Tyrosine Kinase Inhibitor Mammalian Target of Rapamycin Inhibitor as Second-line Molecular-targeted Therapy for Patients with Poor-risk Metastatic Renal Cell Carcinoma. <i>Anticancer Research</i> , 2017 , 37, 1523-1528	2.3	1
34	Discontinuation of first-line molecular-targeted therapy and prognosis in patients with metastatic renal cell carcinoma: Impact of disease progression vs. adverse events. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2020 , 38, 937.e19-937.e25	2.8	1
33	Prognostic Stratification of the IMDC Intermediate Risk Group After Treatment With First-line Molecular-targeted Therapy for Metastatic Renal Cell Carcinoma. <i>Anticancer Research</i> , 2020 , 40, 4395-4400	2.3	1
32	Risk Factors for Totally Implantable Central Venous Access Port-related Infection in Patients With Malignancy. <i>Anticancer Research</i> , 2021 , 41, 1547-1553	2.3	1
31	Health-Related Quality of Life in Patients with Small Renal Mass Who Underwent Robot-Assisted Partial Nephrectomy: A Prospective Evaluation. <i>Journal of Endourology</i> , 2021 , 35, 1644-1649	2.7	1
30	Prognostic assessments in patients with upper tract urothelial carcinoma undergoing radical nephroureterectomy and systematic regional lymph node dissection. <i>Urologia</i> , 2021 , 3915603211034943	1.2	1
29	Assessment of Blood Plasma Free-amino Acid Levels in Infertile Men. <i>In Vivo</i> , 2021 , 35, 1843-1847	2.3	1
28	Japanese Expert Panel Meeting on the Management of Prostate Cancer with Bone Metastases. <i>Oncology and Therapy</i> , 2018 , 6, 157-171	2.7	1
27	Molecular Mechanism Mediating Cytotoxic Activity of Cabazitaxel in Docetaxel-resistant Human Prostate Cancer Cells. <i>Anticancer Research</i> , 2021 , 41, 3753-3758	2.3	1

26	Stress protein GRP78 prevents apoptosis induced by calcium ionophore, ionomycin, but not by glycosylation inhibitor, tunicamycin, in human prostate cancer cells 2000 , 77, 396		1
25	Clinicopathological outcome of radical retropubic prostatectomy for 200 men with prostate cancer in a single institution in Japan. <i>Acta Urologica Japonica</i> , 2004 , 50, 151-6		1
24	A DNA adductome analysis revealed a reduction in the global level of C5-hydroxymethyl-2'-deoxycytidine in the non-tumoral upper urinary tract mucosa of urothelial carcinoma patients. <i>Genes and Environment</i> , 2021 , 43, 52	2.8	o
23	Expression Profile of S100A2 and its Clinicopathological Significance in Renal Cell Carcinoma. <i>Anticancer Research</i> , 2020 , 40, 6337-6343	2.3	o
22	Therapeutic Endoscopic Treatment Plus Maintenance Dimethyl Sulfoxide Therapy Prolongs Recurrence-Free Time in Patients With Hunner Type Interstitial Cystitis: A Pilot Study. <i>International Neurourology Journal</i> , 2019 , 23, 327-333	2.6	o
21	Perioperative outcomes in patients undergoing robot-assisted partial nephrectomy: Comparative assessments between complex and non-complex renal tumors. <i>Asian Journal of Endoscopic Surgery</i> , 2021 , 14, 379-385	1.4	o
20	Finite element method for nerve root decompression in minimally invasive endoscopic spinal surgery. <i>Asian Journal of Endoscopic Surgery</i> , 2021 , 14, 628-635	1.4	o
19	Frequency of pre-treatment may not increase the immune-related adverse events of RCC patients treated with nivolumab. <i>Medicine (United States)</i> , 2021 , 100, e25402	1.8	o
18	Renal allograft rejection after treatment with nivolumab in patients with metastatic renal cell carcinoma. <i>International Cancer Conference Journal</i> , 2021 , 10, 116-118	0.9	o
17	Pembrolizumab Combined Chemotherapy With Gemcitabine and Paclitaxel: A Comparative Assessment of Clinical Outcomes in Patients With Platinum-refractory Advanced Urothelial Cancer. <i>In Vivo</i> , 2021 , 35, 1889-1894	2.3	o
16	Avelumab (Ave) first-line (1L) maintenance plus best supportive care (BSC) versus BSC alone for advanced urothelial carcinoma (UC): JAVELIN Bladder 100 Japanese subgroup analysis.. <i>Journal of Clinical Oncology</i> , 2021 , 39, 425-425	2.2	o
15	Impacts of cachexia progression in addition to serum IgG and blood lymphocytes on serum nivolumab in advanced cancer patients. <i>European Journal of Clinical Pharmacology</i> , 2021 , 1	2.8	o
14	Impact of Bladder Neck Angle Measured by Postoperative Magnetic Resonance Imaging on Midterm Recovery of Urinary Continence in Prostate Cancer Patients Undergoing Robot-Assisted Radical Prostatectomy. <i>Journal of Endourology</i> , 2021 , 35, 1610-1615	2.7	o
13	Concurrent occurrence of three renal cell carcinomas with different histological subtypes in the same kidney. <i>International Journal of Urology</i> , 2014 , 21, 222-3	2.3	
12	Biomarkers of Potential Therapeutic Value 2013 , 181-187		
11	Oncological Outcome Following Radical Prostatectomy in Patients with pT3 Prostate Cancer. <i>Current Urology</i> , 2008 , 2, 87-91	1.7	
10	Oncological Outcome following Radical Prostatectomy in Japanese Men with Clinical Stage T1c Prostate Cancer. <i>Current Urology</i> , 2009 , 3, 62-66	1.7	
9	Antisense Oligodeoxynucleotide Therapy for Prostate Cancer Targeting Antiapoptotic Genes Involved in the Mechanism Mediating Progression to Androgen Independence. <i>Current Signal Transduction Therapy</i> , 2006 , 1, 219-226	0.8	

8	Prognostic Significance of Bone Metastasis Patterns in Patients with Prostate Cancer. <i>UroOncology</i> , 2004 , 4, 135-138	
7	Global analysis of gene expression profiles in ileum in a rat bladder augmentation model using cDNA microarrays. <i>International Journal of Urology</i> , 2004 , 11, 1009-12	2.3
6	Impact of carcinoma in situ on the outcome of intravesical Bacillus Calmette-Guérin therapy for non-muscle-invasive bladder cancer: a comparative analysis of large real-world data.. <i>International Journal of Clinical Oncology</i> , 2022 , 1	4.2
5	Corticosteroids alleviate adverse events associated with enzalutamide in patients with metastatic castration-resistant prostate cancer. <i>Molecular and Clinical Oncology</i> , 2020 , 12, 495-500	1.6
4	Robot-Assisted Partial Nephrectomy for Patients with Multifocal Renal Tumors Arising in a Solitary Kidney: Report of Three Cases. <i>Journal of Endourology Case Reports</i> , 2020 , 6, 370-373	0.3
3	Phase I/II study to evaluate the efficacy of TAS0313, a cancer peptide vaccine, combined with pembrolizumab for locally advanced or metastatic urothelial carcinoma.. <i>Journal of Clinical Oncology</i> , 2021 , 39, 4522-4522	2.2
2	External validation of the albumin, C-reactive protein and lactate dehydrogenase model in patients with metastatic renal cell carcinoma receiving second-line axitinib therapy in a Japanese multi-center cohort. <i>Japanese Journal of Clinical Oncology</i> , 2021 , 51, 810-818	2.8
1	Reduction of residual tumors by photodynamic diagnosis-assisted TURBT using 5-aminolevulinic acid for high-risk nonmuscle-invasive bladder cancer (BRIGHT study).. <i>Journal of Clinical Oncology</i> , 2022 , 40, 489-489	2.2