

Ulrich John

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7462065/ulrich-john-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

306
papers

15,203
citations

56
h-index

114
g-index

316
ext. papers

17,014
ext. citations

4.7
avg, IF

5.78
L-index

| # | Paper | IF | Citations |
|-----|---|------|-----------|
| 306 | Social Equity in the Efficacy of Computer-Based and In-Person Brief Alcohol Interventions Among General Hospital Patients With At-Risk Alcohol Use: A Randomized Controlled Trial.. <i>JMIR Mental Health</i> , 2022 , 9, e31712 | 6 | |
| 305 | Severity of alcohol dependence and mortality after 20 years in an adult general population sample.. <i>International Journal of Methods in Psychiatric Research</i> , 2022 , e1915 | 4.3 | 0 |
| 304 | The Role of Tobacco Smoking in the Efficacy of Brief Alcohol Intervention: Results from a Randomized Controlled Trial. <i>International Journal of Environmental Research and Public Health</i> , 2022 , 19, 5847 | 4.6 | |
| 303 | Alcohol abstinence and mortality in a general population sample of adults in Germany: A cohort study. <i>PLoS Medicine</i> , 2021 , 18, e1003819 | 11.6 | 1 |
| 302 | Effect of a 1-year short message service in detoxified alcohol-dependent patients: a multi-center, open-label randomized controlled trial. <i>Addiction</i> , 2021 , 116, 1431-1442 | 4.6 | 2 |
| 301 | Brief alcohol intervention at a municipal registry office: reach and retention. <i>European Journal of Public Health</i> , 2021 , 31, 418-423 | 2.1 | 2 |
| 300 | Latent alcohol use patterns and their link to depressive symptomatology in medical care patients. <i>Addiction</i> , 2021 , 116, 1063-1073 | 4.6 | 2 |
| 299 | Effects of a brief alcohol intervention addressing the full spectrum of drinking in an adult general population sample: a randomized controlled trial. <i>Addiction</i> , 2021 , 116, 2056-2066 | 4.6 | 1 |
| 298 | Alkoholkonsum [Bedeutung für die Prävention und Gesundheitsförderung. <i>The Springer Reference Pflege/Therapie, Gesundheit</i> , 2021 , 441-457 | 0.2 | |
| 297 | Sick days in general hospital patients two years after brief alcohol intervention: Secondary outcomes from a randomized controlled trial. <i>Preventive Medicine</i> , 2020 , 139, 106106 | 4.3 | 2 |
| 296 | Mental disorders and total mortality after 20 years in an adult general population sample. <i>European Psychiatry</i> , 2020 , 63, e30 | 6 | 4 |
| 295 | Untreated pathological gamblers: who recovers and who does not?. <i>International Gambling Studies</i> , 2020 , 20, 200-213 | 1.8 | 1 |
| 294 | Stability of At-risk Alcohol Use Screening Results in a General Population Sample. <i>Alcoholism: Clinical and Experimental Research</i> , 2020 , 44, 1312-1320 | 3.7 | 2 |
| 293 | Diagnostic performance of the Alcohol Use Disorders Identification Test (AUDIT) in detecting DSM-5 alcohol use disorders in the General population. <i>Drug and Alcohol Dependence</i> , 2019 , 204, 107530 | 4.9 | 19 |
| 292 | Proactive multipurpose health risk screening in health care settings: Methods, design, and reach. <i>International Journal of Methods in Psychiatric Research</i> , 2019 , 28, e1760 | 4.3 | 9 |
| 291 | Does prior recall of past week alcohol use affect screening results for at-risk drinking? Findings from a randomized study. <i>PLoS ONE</i> , 2019 , 14, e0217595 | 3.7 | 4 |
| 290 | Feasibility and Acceptability of an Intervention Providing Computer-Generated Tailored Feedback to Target Alcohol Consumption and Depressive Symptoms in Proactively Recruited Health Care Patients and Reactively Recruited Media Volunteers: Results of a Pilot Study. <i>European Addiction Research</i> , 2019 , 25, 119-131 | 4.6 | 6 |

| | | | |
|-----|---|-----|----|
| 289 | Alkoholkonsum Bedeutung für Prävention und Gesundheitsförderung. <i>The Springer Reference Pflege, Gesundheit, 2019</i> , 1-17 | 0.2 | |
| 288 | Behavioral health risk factor profiles in general hospital patients: identifying the need for screening and brief intervention. <i>BMC Public Health, 2019</i> , 19, 1594 | 4.1 | 0 |
| 287 | Can brief alcohol interventions in general hospital inpatients improve mental and general health over 2 years? Results from a randomized controlled trial. <i>Psychological Medicine, 2019</i> , 49, 1722-1730 | 6.9 | 9 |
| 286 | How alcohol use problem severity affects the outcome of brief intervention delivered in-person versus through computer-generated feedback letters. <i>Drug and Alcohol Dependence, 2018</i> , 183, 82-88 | 4.9 | 14 |
| 285 | Effect of blood pressure and total cholesterol measurement on risk prediction using the Systematic COronary Risk Evaluation (SCORE). <i>BMC Cardiovascular Disorders, 2018</i> , 18, 84 | 2.3 | 1 |
| 284 | Proactive expert system intervention to prevent or quit at-risk alcohol use (PRINT): study protocol of a randomized controlled trial. <i>BMC Public Health, 2018</i> , 18, 851 | 4.1 | 3 |
| 283 | Do brief alcohol interventions among unemployed at-risk drinkers increase re-employment after 15 month?. <i>European Journal of Public Health, 2018</i> , 28, 510-515 | 2.1 | 1 |
| 282 | Health Risk Behavior Patterns in a National Adult Population Survey. <i>International Journal of Environmental Research and Public Health, 2018</i> , 15, | 4.6 | 16 |
| 281 | Short-term stability of different drinking patterns over the course of four weeks among adults. A latent transition analysis. <i>Drug and Alcohol Dependence, 2018</i> , 191, 181-186 | 4.9 | 9 |
| 280 | Measurement invariance of the alcohol use disorders identification test: Establishing its factor structure in different settings and across gender. <i>Drug and Alcohol Dependence, 2018</i> , 189, 55-61 | 4.9 | 13 |
| 279 | Novel genetic associations for blood pressure identified via gene-alcohol interaction in up to 570K individuals across multiple ancestries. <i>PLoS ONE, 2018</i> , 13, e0198166 | 3.7 | 31 |
| 278 | In-person alcohol counseling versus computer-generated feedback: Results from a randomized controlled trial. <i>Health Psychology, 2018</i> , 37, 70-80 | 5 | 19 |
| 277 | Socioeconomic factors and heavy drinking over the course of the week among job seekers with at-risk alcohol use. <i>Archives of Environmental and Occupational Health, 2018</i> , 73, 154-161 | 2 | |
| 276 | Who benefits from computer-based brief alcohol intervention? Day-to-day drinking patterns as a moderator of intervention efficacy. <i>Drug and Alcohol Dependence, 2017</i> , 175, 119-126 | 4.9 | 7 |
| 275 | Reach of Individuals at Risk for Cardiovascular Disease by Proactive Recruitment Strategies in General Practices, Job Centers, and Health Insurance. <i>International Journal of Behavioral Medicine, 2017</i> , 24, 153-160 | 2.6 | 12 |
| 274 | Does impaired mental health interfere with the outcome of brief alcohol intervention at general hospitals?. <i>Journal of Consulting and Clinical Psychology, 2017</i> , 85, 562-573 | 6.5 | 7 |
| 273 | Motivating smokers to quit using computer-generated letters that target either reduction or cessation: A population-based randomized controlled trial among smokers who do not intend to quit. <i>Drug and Alcohol Dependence, 2016</i> , 166, 177-86 | 4.9 | 11 |
| 272 | In-person and computer-based alcohol interventions at general hospitals: reach and retention. <i>European Journal of Public Health, 2016</i> , 26, 844-849 | 2.1 | 18 |

| | | | |
|-----|--|-----|----|
| 271 | Sex- and age-specific associations between major depressive disorder and metabolic syndrome in two general population samples in Germany. <i>Nordic Journal of Psychiatry</i> , 2016 , 70, 611-20 | 2.3 | 15 |
| 270 | The inverse link between genetic risk for schizophrenia and migraine through NMDA (N-methyl-D-aspartate) receptor activation via D-serine. <i>European Neuropsychopharmacology</i> , 2016 , 26, 1507-1515 | 1.2 | 9 |
| 269 | Effect of the interaction between childhood abuse and rs1360780 of the FKBP5 gene on gray matter volume in a general population sample. <i>Human Brain Mapping</i> , 2016 , 37, 1602-13 | 5.9 | 47 |
| 268 | Belief incongruence and the intention-behavior gap in persons with at-risk alcohol use. <i>Addictive Behaviors</i> , 2015 , 48, 5-11 | 4.2 | 7 |
| 267 | Factors associated with secondhand smoke exposure prevalence and secondhand smoke level of children living with parental smokers: a cross sectional study. <i>Journal of Community Health</i> , 2015 , 40, 501-7 | 4 | 5 |
| 266 | Sex differences in urine metabolites related with risk of diabetes using NMR spectroscopy: results of the study of health in pomerania. <i>Metabolomics</i> , 2015 , 11, 1405-1415 | 4.7 | 14 |
| 265 | Patterns of health risk behaviors among job-seekers: a latent class analysis. <i>International Journal of Public Health</i> , 2015 , 60, 111-9 | 4 | 10 |
| 264 | Liver cirrhosis mortality, alcohol consumption and tobacco consumption over a 62 year period in a high alcohol consumption country: a trend analysis. <i>BMC Research Notes</i> , 2015 , 8, 822 | 2.3 | 12 |
| 263 | The Stability of Social Desirability: A Latent Change Analysis. <i>Journal of Personality</i> , 2015 , 83, 404-12 | 4.4 | 9 |
| 262 | Social support and the serotonin transporter genotype (5-HTTLPR) moderate levels of resilience, sense of coherence, and depression. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2015 , 168B, 383-91 | 3.5 | 16 |
| 261 | A pilot study on the feasibility and acceptability of a text message-based aftercare treatment programme among alcohol outpatients. <i>Alcohol and Alcoholism</i> , 2015 , 50, 188-94 | 3.5 | 24 |
| 260 | The impact of a stage tailored intervention on alcohol use trajectories among those who do not intend to change. <i>Drug and Alcohol Dependence</i> , 2015 , 147, 167-74 | 4.9 | 2 |
| 259 | Prevalence trends in lifestyle-related risk factors. <i>Deutsches A&#x0308;rztblatt International</i> , 2015 , 112, 185-92 | 2.5 | 32 |
| 258 | Persönliche vs. computerbasierte Alkoholintervention für Krankenhauspatienten: Studiendesign. <i>Sucht</i> , 2015 , 61, 347-355 | 0.5 | 4 |
| 257 | Occupational exposure to asbestos is associated with increased mortality in men recruited for a population-based study in Germany. <i>International Journal of Occupational Medicine and Environmental Health</i> , 2015 , 28, 849-62 | 1.5 | |
| 256 | Does stage tailoring matter in brief alcohol interventions for job-seekers? A randomized controlled trial. <i>Addiction</i> , 2014 , 109, 1845-56 | 4.6 | 19 |
| 255 | Reducing tobacco smoke exposure in children aged below 4 years - a randomized controlled trial. <i>Preventive Medicine</i> , 2014 , 69, 208-13 | 4.3 | 2 |
| 254 | The impact of childhood trauma on depression: does resilience matter? Population-based results from the Study of Health in Pomerania. <i>Journal of Psychosomatic Research</i> , 2014 , 77, 97-103 | 4.1 | 53 |

| | | | |
|-----|---|------|----|
| 253 | Predictors of indoor smoking at young children's homes--a cross-sectional study. <i>European Journal of Pediatrics</i> , 2014 , 173, 1187-91 | 4.1 | 8 |
| 252 | The phenomenology of the first panic attack in clinical and community-based samples. <i>Journal of Anxiety Disorders</i> , 2014 , 28, 522-9 | 10.9 | 13 |
| 251 | Alexithymia and brain gray matter volumes in a general population sample. <i>Human Brain Mapping</i> , 2014 , 35, 5932-45 | 5.9 | 47 |
| 250 | A surveillance tool using mobile phone short message service to reduce alcohol consumption among alcohol-dependent patients. <i>Alcoholism: Clinical and Experimental Research</i> , 2014 , 38, 1728-36 | 3.7 | 29 |
| 249 | High social support buffers the effects of 5-HTTLPR genotypes within social anxiety disorder. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2014 , 264, 433-9 | 5.1 | 20 |
| 248 | Hilfen zu problematischem Alkoholkonsum [Noch kommen wir zu sp]. <i>Sucht</i> , 2014 , 60, 107-113 | 0.5 | 3 |
| 247 | Reaching families at their homes for an intervention to reduce tobacco smoke exposure among infants. <i>Journal of Community Health</i> , 2013 , 38, 215-20 | 4 | 4 |
| 246 | Predictors of hazardous drinking, tobacco smoking and physical inactivity in vocational school students. <i>BMC Public Health</i> , 2013 , 13, 475 | 4.1 | 27 |
| 245 | Gender-specific predictors of risky alcohol use among general hospital inpatients. <i>General Hospital Psychiatry</i> , 2013 , 35, 9-15 | 5.6 | 9 |
| 244 | Excess mortality of alcohol-dependent individuals after 14 years and mortality predictors based on treatment participation and severity of alcohol dependence. <i>Alcoholism: Clinical and Experimental Research</i> , 2013 , 37, 156-63 | 3.7 | 46 |
| 243 | Associations between the oxytocin receptor gene (OXTR) and "mind-reading" in humans--an exploratory study. <i>Nordic Journal of Psychiatry</i> , 2013 , 67, 15-21 | 2.3 | 45 |
| 242 | A genomewide association study of smoking relapse in four European population-based samples. <i>Psychiatric Genetics</i> , 2013 , 23, 143-52 | 2.9 | 4 |
| 241 | Periodontitis is related to lung volumes and airflow limitation: a cross-sectional study. <i>European Respiratory Journal</i> , 2013 , 42, 1524-35 | 13.6 | 20 |
| 240 | Testing the diathesis-stress model: 5-HTTLPR, childhood emotional maltreatment, and vulnerability to social anxiety disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2013 , 162B, 253-61 | 3.5 | 18 |
| 239 | Relationship of smoking and/or passive exposure to tobacco smoke on the association between serum thyrotropin and body mass index in large groups of adolescents and children. <i>Thyroid</i> , 2013 , 23, 262-8 | 6.2 | 26 |
| 238 | Association of socioeconomic status with iodine supply and thyroid disorders in northeast Germany. <i>Thyroid</i> , 2013 , 23, 346-53 | 6.2 | 26 |
| 237 | How well do TTM measures work among a sample of individuals with unhealthy alcohol use that is characterized by low readiness to change?. <i>Psychology of Addictive Behaviors</i> , 2013 , 27, 573-83 | 3.4 | 9 |
| 236 | Correlates of adverse outcomes in abdominally obese individuals: findings from the five-year followup of the population-based study of health in Pomerania. <i>Journal of Obesity</i> , 2013 , 2013, 762012 | 3.7 | 6 |

| | | | |
|-----|--|------|-----|
| 235 | Efficacy of a text message-based smoking cessation intervention for young people: a cluster randomized controlled trial. <i>Journal of Medical Internet Research</i> , 2013 , 15, e171 | 7.6 | 85 |
| 234 | A pre-post study on the appropriateness and effectiveness of a Web- and text messaging-based intervention to reduce problem drinking in emerging adults. <i>Journal of Medical Internet Research</i> , 2013 , 15, e196 | 7.6 | 39 |
| 233 | Alexithymie, traumatischer Stress und posttraumatische Belastungsstörung [Befunde aus der Allgemeinbevölkerung. <i>Zeitschrift Fur Psychiatrie, Psychologie Und Psychotherapie</i> , 2013 , 61, 27-36 | 0.3 | 2 |
| 232 | Age-specific reference ranges for serum testosterone and androstenedione concentrations in women measured by liquid chromatography-tandem mass spectrometry. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012 , 97, 408-15 | 5.6 | 126 |
| 231 | Genetic epistasis between the brain-derived neurotrophic factor Val66Met polymorphism and the 5-HTT promoter polymorphism moderates the susceptibility to depressive disorders after childhood abuse. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2012 , 36, 264-70 | 5.5 | 63 |
| 230 | Adoption, reach and effectiveness of computer-based, practitioner delivered and combined smoking interventions in general medical practices: a three-arm cluster randomized trial. <i>Drug and Alcohol Dependence</i> , 2012 , 121, 124-32 | 4.9 | 17 |
| 229 | FTO genotype is associated with phenotypic variability of body mass index. <i>Nature</i> , 2012 , 490, 267-72 | 50.4 | 304 |
| 228 | Low testosterone concentrations in men contribute to the gender gap in cardiovascular morbidity and mortality. <i>Gender Medicine</i> , 2012 , 9, 557-68 | | 25 |
| 227 | Efficacy of a text messaging (SMS) based smoking cessation intervention for adolescents and young adults: study protocol of a cluster randomised controlled trial. <i>BMC Public Health</i> , 2012 , 12, 51 | 4.1 | 48 |
| 226 | Moderation of adult depression by the serotonin transporter promoter variant (5-HTTLPR), childhood abuse and adult traumatic events in a general population sample. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2012 , 159B, 298-309 | 3.5 | 43 |
| 225 | The association between fatty liver disease and blood pressure in a population-based cohort study. <i>Journal of Hypertension</i> , 2012 , 30, 1260-1 | 1.9 | 4 |
| 224 | Efficacy of an internet program for smoking cessation during and after inpatient rehabilitation treatment: a quasi-randomized controlled trial. <i>Addictive Behaviors</i> , 2011 , 36, 1369-72 | 4.2 | 14 |
| 223 | Overestimation of drinking norms and its association with alcohol consumption in apprentices. <i>Alcohol and Alcoholism</i> , 2011 , 46, 204-9 | 3.5 | 25 |
| 222 | Moderation of adult depression by a polymorphism in the FKBP5 gene and childhood physical abuse in the general population. <i>Neuropsychopharmacology</i> , 2011 , 36, 1982-91 | 8.7 | 191 |
| 221 | Multiple dimensions of health locus of control in a representative population sample: ordinal factor analysis and cross-validation of an existing three and a new four factor model. <i>BMC Medical Research Methodology</i> , 2011 , 11, 114 | 4.7 | 12 |
| 220 | Health risk factors and self-rated health among job-seekers. <i>BMC Public Health</i> , 2011 , 11, 659 | 4.1 | 40 |
| 219 | Development and psychometric properties of a short situational urge to smoke scale (SUS) for smoking adolescents. <i>Substance Use and Misuse</i> , 2011 , 46, 959-68 | 2.2 | 3 |
| 218 | Smokers' expectations toward the engagement of their general practitioner in discussing lifestyle behaviors. <i>Journal of Health Communication</i> , 2011 , 16, 135-47 | 2.5 | 9 |

| | | |
|-----|--|-----------|
| 217 | Cohort profile: the study of health in Pomerania. <i>International Journal of Epidemiology</i> , 2011 , 40, 294-307.8 | 704 |
| 216 | Kommentare zu Rumpf und Kiefer (2011). DSM-5: Die Aufhebung der Unterscheidung von Abhängigkeit und Missbrauch und die Öffnung für Verhaltensstile. <i>Sucht</i> , 2011 , 57, 52-53 | 0.5 1 |
| 215 | Meta-analysis and imputation refines the association of 15q25 with smoking quantity. <i>Nature Genetics</i> , 2010 , 42, 436-40 | 36.3 521 |
| 214 | Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. <i>Nature Genetics</i> , 2010 , 42, 937-48 | 36.3 2267 |
| 213 | Measuring readiness to change for problematic consumption of prescription drugs: Development of an adapted and shortened Readiness to Change Questionnaire. <i>Addiction Research and Theory</i> , 2010 , 18, 110-118 | 2.6 2 |
| 212 | Predictive modeling of health care costs: do cardiovascular risk markers improve prediction?. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2010 , 17, 355-62 | 4 |
| 211 | The predictive value of different measures of obesity for incident cardiovascular events and mortality. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, 1777-85 | 5.6 268 |
| 210 | Predicting the urge to smoke in a population-based sample of adolescent smokers. <i>Substance Use and Misuse</i> , 2010 , 45, 1636-51 | 2.2 1 |
| 209 | Loci from a genome-wide analysis of bilirubin levels are associated with gallstone risk and composition. <i>Gastroenterology</i> , 2010 , 139, 1942-1951.e2 | 13.3 74 |
| 208 | Population-based smoking cessation in women post partum: adherence to motivational interviewing in relation to client characteristics and behavioural outcomes. <i>Midwifery</i> , 2010 , 26, 202-10 | 2.8 12 |
| 207 | Alexithymia, hypertension, and subclinical atherosclerosis in the general population. <i>Journal of Psychosomatic Research</i> , 2010 , 68, 139-47 | 4.1 70 |
| 206 | Effects of practitioner-delivered brief counseling and computer-generated tailored letters on cigarettes per day among smokers who do not quit—a quasi-randomized controlled trial. <i>Drug and Alcohol Dependence</i> , 2010 , 112, 81-9 | 4.9 5 |
| 205 | Problematischer Alkoholkonsum: Prävalenz und Frühintervention. <i>Public Health Forum</i> , 2010 , 18, 29-30 | 0.1 |
| 204 | Alcohol dependent inpatients who receive general hospital care vs. detoxification in psychiatric care and alcohol problem 1 year later. <i>Addictive Behaviors</i> , 2010 , 35, 756-63 | 4.2 7 |
| 203 | Heart valve sclerosis predicts all-cause and cardiovascular mortality. <i>Atherosclerosis</i> , 2010 , 209, 606-10 | 3.1 79 |
| 202 | The impact of having a loved one with alcohol consumption-related problems on subjective health status and health-risk behaviors in a general hospital sample. <i>Substance Use and Misuse</i> , 2010 , 45, 2470-80 ² | 2.2 3 |
| 201 | The association between fatty liver disease and blood pressure in a population-based prospective longitudinal study. <i>Journal of Hypertension</i> , 2010 , 28, 1829-35 | 1.9 47 |
| 200 | Need for cognition as a predictor and a moderator of outcome in a tailored letters smoking cessation intervention. <i>Health Psychology</i> , 2010 , 29, 367-73 | 5 16 |

| | | | |
|-----|--|------|-----|
| 199 | Workplace smoking restrictions: smoking behavior and the intention to change among continuing smokers. <i>International Journal of Public Health</i> , 2010 , 55, 599-608 | 4 | 7 |
| 198 | Severity of unhealthy alcohol consumption in medical inpatients and the general population: is the general hospital a suitable place for brief interventions?. <i>International Journal of Public Health</i> , 2010 , 55, 637-43 | 4 | 8 |
| 197 | Association of posttraumatic stress disorder with low-grade elevation of C-reactive protein: evidence from the general population. <i>Journal of Psychiatric Research</i> , 2010 , 44, 15-21 | 5.2 | 153 |
| 196 | A descriptive analysis of relations between parents' self-reported smoking behavior and infants' daily exposure to environmental tobacco smoke. <i>BMC Public Health</i> , 2010 , 10, 424 | 4.1 | 6 |
| 195 | Motivation to change risky drinking and motivation to seek help for alcohol risk drinking among general hospital inpatients with problem drinking and alcohol-related diseases. <i>General Hospital Psychiatry</i> , 2010 , 32, 86-93 | 5.6 | 22 |
| 194 | Childhood maltreatment, the corticotropin-releasing hormone receptor gene and adult depression in the general population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2010 , 153B, 1483-93 | 3.5 | 84 |
| 193 | Predictors and moderators of outcome in different brief interventions for smoking cessation in general medical practice. <i>Patient Education and Counseling</i> , 2010 , 78, 57-64 | 3.1 | 26 |
| 192 | Internetbasierte Förderung der Tabakabstinenz in der Rehabilitation. <i>Sucht</i> , 2010 , 56, 117-124 | 0.5 | |
| 191 | Smoking cessation counselling for pregnant and postpartum women among midwives, gynaecologists and paediatricians in Germany. <i>International Journal of Environmental Research and Public Health</i> , 2009 , 6, 96-107 | 4.6 | 12 |
| 190 | At-risk and heavy episodic drinking, motivation to change, and the development of alcohol dependence among men. <i>Journal of Studies on Alcohol and Drugs</i> , 2009 , 70, 937-42 | 1.9 | 2 |
| 189 | Remission ohne formelle Hilfe bei Alkoholabhängigkeit: Der Stand der Forschung. <i>Sucht</i> , 2009 , 55, 75-85 | 0.5 | 9 |
| 188 | Serotonin transporter gene (SLC6A4) promoter polymorphisms and the susceptibility to posttraumatic stress disorder in the general population. <i>American Journal of Psychiatry</i> , 2009 , 166, 926-33 | 11.9 | 142 |
| 187 | Continuous individual support of smoking cessation using text messaging: a pilot experimental study. <i>Nicotine and Tobacco Research</i> , 2009 , 11, 915-23 | 4.9 | 99 |
| 186 | Reported beverage consumed and alcohol-related diseases among male hospital inpatients with problem drinking. <i>Alcohol and Alcoholism</i> , 2009 , 44, 216-21 | 3.5 | 1 |
| 185 | Intention to change smoking in pregnant and postpartum women according to number of pregnancies. <i>European Journal of Public Health</i> , 2009 , 19, 218-21 | 2.1 | 9 |
| 184 | Subclinical hyperthyroidism and blood pressure in a population-based prospective cohort study. <i>European Journal of Endocrinology</i> , 2009 , 161, 615-21 | 6.5 | 33 |
| 183 | Early onset of obsessive-compulsive disorder and associated comorbidity. <i>Depression and Anxiety</i> , 2009 , 26, 1012-7 | 8.4 | 36 |
| 182 | Mental health and readiness to change smoking behavior in daily smoking primary care patients. <i>International Journal of Behavioral Medicine</i> , 2009 , 16, 347-54 | 2.6 | 3 |

| | | | |
|-----|---|------|-----|
| 181 | Support for smoke-free policies in a pro-smoking culture: findings from the European survey on tobacco control attitudes and knowledge. <i>International Journal of Public Health</i> , 2009 , 54, 403-8 | | 20 |
| 180 | Association of mental distress with health care utilization and costs: a 5-year observation in a general population. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2009 , 44, 835-44 | 4.5 | 55 |
| 179 | Randomized controlled trial of a brief intervention for problematic prescription drug use in non-treatment-seeking patients. <i>Addiction</i> , 2009 , 104, 109-17 | 4.6 | 52 |
| 178 | Tooth loss and cognitive impairment. <i>Journal of Clinical Periodontology</i> , 2009 , 36, 550-7 | 7.7 | 62 |
| 177 | Nicotine dependence criteria and nicotine withdrawal symptoms in relation to pain among an adult general population sample. <i>European Journal of Pain</i> , 2009 , 13, 82-8 | 3.7 | 14 |
| 176 | Does the smoking status of general practitioners affect the efficacy of smoking cessation counselling?. <i>Patient Education and Counseling</i> , 2009 , 74, 23-8 | 3.1 | 16 |
| 175 | Brief intervention in general hospital for problematic prescription drug use: 12-month outcome. <i>Drug and Alcohol Dependence</i> , 2009 , 105, 221-6 | 4.9 | 32 |
| 174 | Extended recruitment efforts minimize attrition but not necessarily bias. <i>Journal of Clinical Epidemiology</i> , 2009 , 62, 252-60 | 5.7 | 72 |
| 173 | Smoking cessation and relapse prevention for postpartum women: results from a randomized controlled trial at 6, 12, 18 and 24 months. <i>Addictive Behaviors</i> , 2009 , 34, 1-8 | 4.2 | 36 |
| 172 | Trauma, posttraumatic stress disorder, and physical illness: findings from the general population. <i>Psychosomatic Medicine</i> , 2009 , 71, 1012-7 | 3.7 | 189 |
| 171 | The performance of two motivation measures and outcome after alcohol detoxification. <i>Alcohol and Alcoholism</i> , 2009 , 44, 77-83 | 3.5 | 5 |
| 170 | A genome-wide association study identifies colorectal cancer susceptibility loci on chromosomes 10p14 and 8q23.3. <i>Nature Genetics</i> , 2008 , 40, 623-30 | 36.3 | 463 |
| 169 | Proactive interventions for smoking cessation in general medical practice: a quasi-randomized controlled trial to examine the efficacy of computer-tailored letters and physician-delivered brief advice. <i>Addiction</i> , 2008 , 103, 294-304 | 4.6 | 60 |
| 168 | [Commentary] SUPPORTING GPS IN THE PROVISION OF SMOKING CESSATION INTERVENTION FROM RESEARCH TO DAILY ROUTINE. <i>Addiction</i> , 2008 , 103, 309-309 | 4.6 | |
| 167 | Reasons for not using smoking cessation aids. <i>BMC Public Health</i> , 2008 , 8, 129 | 4.1 | 27 |
| 166 | The relationship between smokers' motivation to quit and intensity of tobacco control at the population level: a comparison of five European countries. <i>BMC Public Health</i> , 2008 , 8, 2 | 4.1 | 31 |
| 165 | Alcohol problem drinking among general hospital inpatients in northeastern Germany. <i>General Hospital Psychiatry</i> , 2008 , 30, 147-54 | 5.6 | 17 |
| 164 | Computer-generated tailored feedback letters for smoking cessation: theoretical and empirical variability of tailoring. <i>International Journal of Medical Informatics</i> , 2008 , 77, 715-22 | 5.3 | 15 |

| | | | |
|-----|---|------|-----|
| 163 | Harm avoidance in subjects with obsessive-compulsive disorder and their families. <i>Journal of Affective Disorders</i> , 2008 , 107, 265-9 | 6.6 | 48 |
| 162 | Impact of fatty liver disease on health care utilization and costs in a general population: a 5-year observation. <i>Gastroenterology</i> , 2008 , 134, 85-94 | 13.3 | 155 |
| 161 | Brief alcohol intervention for general hospital inpatients: a randomized controlled trial. <i>Drug and Alcohol Dependence</i> , 2008 , 93, 233-43 | 4.9 | 50 |
| 160 | Evaluation of a telephone-based stepped care intervention for alcohol-related disorders: a randomized controlled trial. <i>Drug and Alcohol Dependence</i> , 2008 , 93, 244-51 | 4.9 | 75 |
| 159 | Effectiveness of brief alcohol interventions for general practice patients with problematic drinking behavior and comorbid anxiety or depressive disorders. <i>Drug and Alcohol Dependence</i> , 2008 , 94, 214-20 | 4.9 | 32 |
| 158 | Relation of self-reported sleep duration with carotid intima-media thickness in a general population sample. <i>Atherosclerosis</i> , 2008 , 196, 727-32 | 3.1 | 75 |
| 157 | The relation of exposure to shift work with atherosclerosis and myocardial infarction in a general population. <i>Atherosclerosis</i> , 2008 , 201, 205-11 | 3.1 | 89 |
| 156 | Association between parity and autoimmune thyroiditis in a general female population. <i>Autoimmunity</i> , 2008 , 41, 174-80 | 3 | 36 |
| 155 | Computer-tailored smoking cessation intervention in a general population setting in Germany: outcome of a randomized controlled trial. <i>Nicotine and Tobacco Research</i> , 2008 , 10, 371-9 | 4.9 | 13 |
| 154 | Smoking as a risk factor for thyroid volume progression and incident goiter in a region with improved iodine supply. <i>European Journal of Endocrinology</i> , 2008 , 159, 761-6 | 6.5 | 19 |
| 153 | Obsessive-compulsive disorder and posttraumatic stress disorder. <i>Psychopathology</i> , 2008 , 41, 129-34 | 3.4 | 28 |
| 152 | Does response bias influence population studies of thyroid disorders?. <i>Thyroid</i> , 2008 , 18, 873-8 | 6.2 | 5 |
| 151 | Smoking during pregnancy and postpartum: smoking rates and intention to quit smoking or resume after pregnancy. <i>Journal of Women's Health</i> , 2008 , 17, 631-40 | 3 | 26 |
| 150 | Gender differences in the efficacy of brief interventions with a stepped care approach in general practice patients with alcohol-related disorders. <i>Alcohol and Alcoholism</i> , 2008 , 43, 334-40 | 3.5 | 21 |
| 149 | No association of seropositivity for anti-Borrelia IgG antibody with mental and physical complaints. <i>Nordic Journal of Psychiatry</i> , 2008 , 62, 386-91 | 2.3 | 5 |
| 148 | Does precontemplation represent a homogeneous stage category? A latent class analysis on German smokers. <i>Journal of Consulting and Clinical Psychology</i> , 2008 , 76, 840-51 | 6.5 | 9 |
| 147 | Prevalence and determinants of increased serum lipase levels in a general population. <i>Pancreas</i> , 2008 , 37, 411-7 | 2.6 | 5 |
| 146 | Male at-risk drinkers with heavy episodic drinking: a subclinical diagnosis?. <i>Journal of Studies on Alcohol and Drugs</i> , 2008 , 69, 85-90 | 1.9 | 1 |

| | | | |
|-----|--|------|----|
| 145 | Kurzinterventionen bei Alkoholproblemen im Allgemeinkrankenhaus. <i>Psychosomatik Und Konsiliarpsychiatrie</i> , 2008 , 2, 8-14 | | 3 |
| 144 | Predicting utilization of formal and informal help among general hospital inpatients with alcohol use disorders. <i>International Journal of Methods in Psychiatric Research</i> , 2008 , 17 Suppl 1, S70-3 | 4.3 | 4 |
| 143 | Differences in help seeking rates after brief intervention for alcohol use disorders in general practice patients with and without comorbid anxiety or depressive disorders. <i>International Journal of Methods in Psychiatric Research</i> , 2008 , 17 Suppl 1, S74-7 | 4.3 | 14 |
| 142 | Dose-response relation between volume of drinking and alcohol-related diseases in male general hospital inpatients. <i>Alcohol and Alcoholism</i> , 2008 , 43, 34-8 | 3.5 | 18 |
| 141 | Trauma and posttraumatic stress disorder in the elderly: findings from a German community study. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 693-700 | 4.6 | 70 |
| 140 | Patterns and predictors of change in a smoking intervention study: latent growth analysis of a multivariate outcome model. <i>Health Psychology</i> , 2008 , 27, S233-42 | 5 | 6 |
| 139 | Menopausal status and hepatic steatosis in a general female population. <i>Gut</i> , 2007 , 56, 594-5 | 19.2 | 99 |
| 138 | Alcohol consumption is associated with an interaction between DRD2 exon 8 A/A genotype and self-directedness in males. <i>Neuropsychobiology</i> , 2007 , 56, 24-31 | 4 | 12 |
| 137 | The spectrum of thyroid disorders in adult type 1 diabetes mellitus. <i>Diabetes/Metabolism Research and Reviews</i> , 2007 , 23, 227-33 | 7.5 | 19 |
| 136 | Stability of subtypes of natural recovery from alcohol dependence after two years. <i>Addiction</i> , 2007 , 102, 904-8 | 4.6 | 26 |
| 135 | Sedative, hypnotic, anxiolytic and opioid medicament use and its co-occurrence with tobacco smoking and alcohol risk drinking in a community sample. <i>BMC Public Health</i> , 2007 , 7, 337 | 4.1 | 14 |
| 134 | The glu298asp polymorphism in the nitric oxide synthase 3 gene is associated with the risk of ischemic stroke in two large independent case-control studies. <i>Human Genetics</i> , 2007 , 121, 169-78 | 6.3 | 62 |
| 133 | Development and evaluation of a screening instrument for alcohol-use disorders and at-risk drinking: the brief alcohol screening instrument for medical care (BASIC). <i>Journal of Studies on Alcohol and Drugs</i> , 2007 , 68, 607-14 | 1.9 | 23 |
| 132 | Menopause and determinants of quality of life in women at midlife and beyond: the study of health in pomerania (SHIP). <i>Menopause</i> , 2007 , 14, 123-34 | 2.5 | 38 |
| 131 | Inverse association between number of teeth and left ventricular mass in women. <i>Journal of Hypertension</i> , 2007 , 25, 2035-43 | 1.9 | 18 |
| 130 | Adherence to the principles of Motivational Interviewing, clients characteristics and behavior outcome in a smoking cessation and relapse prevention trial in women postpartum. <i>Addictive Behaviors</i> , 2007 , 32, 2297-303 | 4.2 | 40 |
| 129 | Variability of tailoring of a smoking cessation intervention based on the transtheoretical model. <i>Addictive Behaviors</i> , 2007 , 32, 3083-7 | 4.2 | 9 |
| 128 | Intention to utilize formal help in a sample with alcohol problems: a prospective study. <i>Drug and Alcohol Dependence</i> , 2007 , 87, 210-6 | 4.9 | 18 |

| | | | |
|-----|--|-----|----|
| 127 | Effects of smoking cessation on health care use: is elevated risk of hospitalization among former smokers attributable to smoking-related morbidity?. <i>Drug and Alcohol Dependence</i> , 2007 , 88, 197-203 | 4.9 | 9 |
| 126 | Population impact--definition, calculation and its use in prevention science in the example of tobacco smoking reduction. <i>Health Policy</i> , 2007 , 82, 348-56 | 3.2 | 15 |
| 125 | Outcome after coronary artery bypass graft surgery, coronary angioplasty and stenting. <i>International Journal of Cardiology</i> , 2007 , 116, 46-52 | 3.2 | 12 |
| 124 | Skalen zur Erfassung der Konstrukte des Transtheoretischen Modells zur Förderung des Rauchverhaltens. <i>Zeitschrift Fur Gesundheitspsychologie</i> , 2007 , 15, 67-77 | | 6 |
| 123 | Remission without Formal Help: New Directions in Studies Using Survey Data 2007 , 73-85 | | 4 |
| 122 | Subjective health complaints, health-related quality of life and physician visits: results of the study of health in Pomerania (SHIP). <i>International Journal of Public Health</i> , 2006 , 51, 162-73 | | 8 |
| 121 | Provision of smoking cessation counseling by general practitioners assisted by training and screening procedure. <i>Patient Education and Counseling</i> , 2006 , 63, 232-8 | 3.1 | 31 |
| 120 | An intervention to support postpartum women to quit smoking or remain smoke-free. <i>Journal of Midwifery and Women's Health</i> , 2006 , 51, 45-50 | 2.2 | 24 |
| 119 | Relation between awareness of circulatory disorders and smoking in a general population health examination. <i>BMC Public Health</i> , 2006 , 6, 48 | 4.1 | 14 |
| 118 | Negative perceived paternal parenting is associated with dopamine D2 receptor exon 8 and GABA(A) alpha 6 receptor variants: an explorative study. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2006 , 141B, 167-72 | 3.5 | 27 |
| 117 | Performance of the pictorial representation of illness and self measure in individuals with alcohol dependence, alcohol abuse or at-risk drinking. <i>Psychotherapy and Psychosomatics</i> , 2006 , 75, 249-56 | 9.4 | 11 |
| 116 | Attitudes towards smoking policies and tobacco control measures in relation to smoking status and smoking behaviour. <i>European Journal of Public Health</i> , 2006 , 16, 513-9 | 2.1 | 37 |
| 115 | Low serum thyrotropin is associated with high plasma fibrinogen. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2006 , 91, 530-4 | 5.6 | 57 |
| 114 | Thyroid function and serum ferritin levels: the study of health in Pomerania. <i>Thyroid</i> , 2006 , 16, 681-6 | 6.2 | 4 |
| 113 | Tele-electrocardiography in the epidemiological study of Health in Pomerania (SHIP). <i>Journal of Telemedicine and Telecare</i> , 2006 , 12, 103-7 | 6.8 | 7 |
| 112 | Alcohol consumption and out-patient services utilization by abstainers and drinkers. <i>Addiction</i> , 2006 , 101, 1285-91 | 4.6 | 15 |
| 111 | Seropositivity for anti-Borrelia IgG antibody is independently associated with carotid atherosclerosis. <i>Atherosclerosis</i> , 2006 , 184, 108-12 | 3.1 | 14 |
| 110 | Alexithymia in obsessive-compulsive disorder - results from a family study. <i>Psychotherapy and Psychosomatics</i> , 2006 , 75, 312-8 | 9.4 | 42 |

| | | | |
|-----|---|------|-----|
| 109 | Psychometric properties of the Processes of Change scale for alcohol misuse and its short form (POC-20). <i>Addictive Behaviors</i> , 2006 , 31, 821-32 | 4.2 | 14 |
| 108 | Postpartum return to smoking: identifying different groups to tailor interventions. <i>Addictive Behaviors</i> , 2006 , 31, 1785-96 | 4.2 | 16 |
| 107 | Tobacco smoking in relation to analgesic drug use in a national adult population sample. <i>Drug and Alcohol Dependence</i> , 2006 , 85, 49-55 | 4.9 | 35 |
| 106 | Midwives' attitudes to counselling women about their smoking behaviour during pregnancy and postpartum. <i>Midwifery</i> , 2006 , 22, 32-9 | 2.8 | 20 |
| 105 | Psychiatric comorbidity including nicotine dependence among individuals with eating disorder criteria in an adult general population sample. <i>Psychiatry Research</i> , 2006 , 141, 71-9 | 9.9 | 14 |
| 104 | Changes in the "stages of change" as outcome measures of a smoking cessation intervention: a randomized controlled trial. <i>Preventive Medicine</i> , 2006 , 43, 101-6 | 4.3 | 21 |
| 103 | Tobacco smoking in relation to pain in a national general population survey. <i>Preventive Medicine</i> , 2006 , 43, 477-81 | 4.3 | 87 |
| 102 | Alcohol consumption and health-services utilization in Germany. <i>Journal of Studies on Alcohol and Drugs</i> , 2006 , 67, 429-35 | | 15 |
| 101 | Stability of remission from alcohol dependence without formal help. <i>Alcohol and Alcoholism</i> , 2006 , 41, 311-4 | 3.5 | 20 |
| 100 | Gender differences in the relation between number of teeth and systolic blood pressure. <i>Journal of Hypertension</i> , 2006 , 24, 1257-63 | 1.9 | 42 |
| 99 | Familiality of obsessive-compulsive disorder in nonclinical and clinical subjects. <i>American Journal of Psychiatry</i> , 2006 , 163, 1986-92 | 11.9 | 74 |
| 98 | The association between subclinical hyperthyroidism and blood pressure in a population-based study. <i>Journal of Hypertension</i> , 2006 , 24, 1947-53 | 1.9 | 36 |
| 97 | The reliability and validity of the Alcohol Use Disorders Identification Test (AUDIT) in a German general practice population sample. <i>Journal of Studies on Alcohol and Drugs</i> , 2006 , 67, 473-81 | | 88 |
| 96 | Measuring activities in tobacco control across the EU. The MATOC. <i>Substance Abuse Treatment, Prevention, and Policy</i> , 2006 , 1, 9 | 3.4 | 10 |
| 95 | Predictors of increased body mass index following cessation of smoking. <i>American Journal on Addictions</i> , 2006 , 15, 192-7 | 3.7 | 16 |
| 94 | Supporting the intention to change health risk behaviors. <i>Zeitschrift Fur Gesundheitswissenschaften</i> , 2006 , 14, 377-383 | 1.4 | 1 |
| 93 | Post-traumatic stress disorder: the role of trauma, pre-existing psychiatric disorders, and gender. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2006 , 256, 299-306 | 5.1 | 102 |
| 92 | Association between alcohol consumption and subclinical carotid atherosclerosis: the Study of Health in Pomerania. <i>Stroke</i> , 2005 , 36, 1746-52 | 6.7 | 50 |

| | | | |
|----|--|-----|-----|
| 91 | Stages of change and other factors in light cigarette smokers. <i>European Journal of Public Health</i> , 2005 , 15, 146-51 | 2.1 | 6 |
| 90 | Self-rated general health and psychiatric disorders in a general population sample. <i>European Psychiatry</i> , 2005 , 20, 223-8 | 6 | 7 |
| 89 | Comparison of a population-based sample of "risky drinking" smokers and groups consuming just one substance. <i>Substance Use and Misuse</i> , 2005 , 40, 1721-32 | 2.2 | |
| 88 | Occupational exposure to ionizing radiation is associated with autoimmune thyroid disease. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005 , 90, 4587-92 | 5.6 | 38 |
| 87 | Tooth loss is independently associated with the risk of acquired aortic valve sclerosis. <i>American Heart Journal</i> , 2005 , 150, 1198-203 | 4.9 | 38 |
| 86 | Reference intervals of serum thyroid function tests in a previously iodine-deficient area. <i>Thyroid</i> , 2005 , 15, 279-85 | 6.2 | 155 |
| 85 | Psychiatric disorders among at-risk consumers of alcohol in the general population. <i>Journal of Studies on Alcohol and Drugs</i> , 2005 , 66, 246-53 | | 31 |
| 84 | Association of smoking and nicotine dependence with trauma and posttraumatic stress disorder in a general population sample. <i>Journal of Nervous and Mental Disease</i> , 2005 , 193, 843-6 | 1.8 | 110 |
| 83 | Endothelial nitric oxide synthase Glu298Asp gene polymorphism, blood pressure and hypertension in a general population sample. <i>Journal of Hypertension</i> , 2005 , 23, 1361-6 | 1.9 | 30 |
| 82 | Relation of parity with common carotid intima-media thickness among women of the Study of Health in Pomerania. <i>Stroke</i> , 2005 , 36, 938-43 | 6.7 | 45 |
| 81 | Identification and functional analysis of genetic variants of the human beta-glucuronidase in a German population sample. <i>Pharmacogenetics and Genomics</i> , 2005 , 15, 875-81 | 1.9 | 10 |
| 80 | Stage of change transitions and processes of change, decisional balance, and self-efficacy in smokers: a transtheoretical model validation using longitudinal data. <i>Psychology of Addictive Behaviors</i> , 2005 , 19, 3-9 | 3.4 | 28 |
| 79 | Endothelial nitric oxide synthase Glu(298)-->Asp polymorphism, carotid atherosclerosis and intima-media thickness in a general population sample. <i>Clinical Science</i> , 2005 , 109, 475-81 | 6.5 | 21 |
| 78 | Longitudinal relationships among transtheoretical model constructs for smokers in the precontemplation and contemplation stages of change. <i>Annals of Behavioral Medicine</i> , 2005 , 30, 12-20 | 4.5 | 5 |
| 77 | Calcium antagonists and deep gingival pockets in the population-based SHIP study. <i>British Journal of Clinical Pharmacology</i> , 2005 , 60, 552-9 | 3.8 | 16 |
| 76 | Relationships of psychiatric disorders with overweight and obesity in an adult general population. <i>Obesity</i> , 2005 , 13, 101-9 | | 68 |
| 75 | Relationships of psychiatric disorders with sleep duration in an adult general population sample. <i>Journal of Psychiatric Research</i> , 2005 , 39, 577-83 | 5.2 | 41 |
| 74 | Gender and age differences among current smokers in a general population survey. <i>BMC Public Health</i> , 2005 , 5, 57 | 4.1 | 12 |

| | | | |
|----|---|-----|-----|
| 73 | Association between thyroid function and gallstone disease. <i>World Journal of Gastroenterology</i> , 2005 , 11, 5530-4 | 5.6 | 25 |
| 72 | Intention to change drinking behaviour in general practice patients with problematic drinking and comorbid depression or anxiety. <i>Alcohol and Alcoholism</i> , 2005 , 40, 394-400 | 3.5 | 20 |
| 71 | Asymptomatic carotid atherosclerosis is associated with circulating chlamydia pneumoniae DNA in younger normotensive subjects in a general population survey. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005 , 25, 386-91 | 9.4 | 11 |
| 70 | The association of thyroid function with cardiac mass and left ventricular hypertrophy. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005 , 90, 673-7 | 5.6 | 106 |
| 69 | Consistency or change in nicotine dependence according to the Fagerström Test for Nicotine Dependence over three years in a population sample. <i>Journal of Addictive Diseases</i> , 2005 , 24, 85-100 | 1.7 | 11 |
| 68 | Readiness for change and readiness for help-seeking: a composite assessment of client motivation. <i>Alcohol and Alcoholism</i> , 2005 , 40, 540-4 | 3.5 | 42 |
| 67 | Putative association between anti-Borrelia IgG and autoimmune thyroid disease?. <i>Thyroid</i> , 2005 , 15, 1273-7 | 3.7 | 7 |
| 66 | Independent risk factors for gallstone formation in a region with high cholelithiasis prevalence. <i>Digestion</i> , 2005 , 71, 97-105 | 3.6 | 141 |
| 65 | Hepatic steatosis is associated with an increased risk of carotid atherosclerosis. <i>World Journal of Gastroenterology</i> , 2005 , 11, 1848-53 | 5.6 | 197 |
| 64 | Autoevaluación de la salud general y trastornos psiquiátricos en una muestra de la población general. <i>European Psychiatry (Ed Española)</i> , 2005 , 12, 349-354 | | |
| 63 | Periodontal disease, but not edentulism, is independently associated with increased plasma fibrinogen levels. Results from a population-based study. <i>Thrombosis and Haemostasis</i> , 2004 , 92, 244-52 | 7 | 40 |
| 62 | Thyroid function and carotid wall thickness. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004 , 89, 2145-9 | 5.6 | 120 |
| 61 | Sex and the risk of restless legs syndrome in the general population. <i>Archives of Internal Medicine</i> , 2004 , 164, 196-202 | | 511 |
| 60 | Prevalence, characteristics, associated mental disorders and predictors of DSM-IV nicotine dependence. <i>European Addiction Research</i> , 2004 , 10, 29-34 | 4.6 | 18 |
| 59 | The relation of a family history of alcoholism, obstetric complications and family environment to behavioral problems among 154 adolescents in Germany: results from the children of alcoholics study in Pomerania. <i>European Addiction Research</i> , 2004 , 10, 8-14 | 4.6 | 10 |
| 58 | Subtyping general population smokers not intending to quit by stages to reduce smoking. <i>Nicotine and Tobacco Research</i> , 2004 , 6, 1043-50 | 4.9 | 7 |
| 57 | Short report: Paediatricians' attitude towards counselling parents postpartum about their smoking behaviour. <i>European Journal of Public Health</i> , 2004 , 14, 199-200 | 2.1 | 8 |
| 56 | Association between high serum ferritin levels and carotid atherosclerosis in the study of health in Pomerania (SHIP). <i>Stroke</i> , 2004 , 35, 453-7 | 6.7 | 84 |

| | | | |
|----|--|-----|-----|
| 55 | Gender differences in the relationship between periodontal disease, tooth loss, and atherosclerosis. <i>Stroke</i> , 2004 , 35, 2029-35 | 6.7 | 122 |
| 54 | Laboratory markers carbohydrate-deficient transferrin, gamma-glutamyltransferase, and mean corpuscular volume are not useful as screening tools for high-risk drinking in the general population: results from the Study of Health in Pomerania (SHIP). <i>Alcoholism: Clinical and Experimental Research</i> , 2004 , 28, 224-28 | 3.7 | 37 |
| 53 | Malocclusions and clinical signs or subjective symptoms of temporomandibular disorders (TMD) in adults. Results of the population-based Study of Health in Pomerania (SHIP). <i>Journal of Orofacial Orthopedics</i> , 2004 , 65, 88-103 | 2.9 | 23 |
| 52 | Impact of psychiatric disorders in the general population: satisfaction with life and the influence of comorbidity and disorder duration. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2004 , 39, 435-41 | 4.5 | 34 |
| 51 | Caries and periodontal disease of the elderly in Pomerania, Germany: results of the Study of Health in Pomerania. <i>Gerodontology</i> , 2004 , 21, 27-36 | 2.8 | 52 |
| 50 | Smoking, nicotine dependence and psychiatric comorbidity--a population-based study including smoking cessation after three years. <i>Drug and Alcohol Dependence</i> , 2004 , 76, 287-95 | 4.9 | 175 |
| 49 | Nicotine dependence, quit attempts, and quitting among smokers in a regional population sample from a country with a high prevalence of tobacco smoking. <i>Preventive Medicine</i> , 2004 , 38, 350-8 | 4.3 | 67 |
| 48 | Hepatitis B and C virus infection and the risk of atherosclerosis in a general population. <i>Atherosclerosis</i> , 2004 , 174, 99-103 | 3.1 | 102 |
| 47 | Nicotine dependence and lifetime amount of smoking in a population sample. <i>European Journal of Public Health</i> , 2004 , 14, 182-5 | 2.1 | 19 |
| 46 | Plasma fibrinogen levels are associated with a strong family history of myocardial infarction. <i>Blood Coagulation and Fibrinolysis</i> , 2004 , 15, 497-502 | 1 | 5 |
| 45 | Readiness to change versus readiness to seek help for alcohol problems: the development of the Treatment Readiness Tool (TReaT). <i>Journal of Studies on Alcohol and Drugs</i> , 2004 , 65, 801-9 | | 25 |
| 44 | Depressive disorders are related to nicotine dependence in the population but do not necessarily hamper smoking cessation. <i>Journal of Clinical Psychiatry</i> , 2004 , 65, 169-76 | 4.6 | 53 |
| 43 | Verhaltensspezifische Determinanten der Veränderungs motivation bei multiplem Risikoverhalten. <i>Zeitschrift Fur Gesundheitspsychologie</i> , 2004 , 12, 56-64 | | 1 |
| 42 | Die Prozesse der Verhaltensänderung im Kontext des Lebensalters am Beispiel des Tabakrauchens. <i>Zeitschrift Fur Gesundheitspsychologie</i> , 2004 , 12, 39-47 | | 0 |
| 41 | Estimation of tobacco- or alcohol-attributable disease rates in national hospital care: an approach based on routine in-patient disease register data and systematic diagnosis of alcohol use disorders. <i>Alcohol and Alcoholism</i> , 2003 , 38, 339-46 | 3.5 | 4 |
| 40 | Strength of the relationship between tobacco smoking, nicotine dependence and the severity of alcohol dependence syndrome criteria in a population-based sample. <i>Alcohol and Alcoholism</i> , 2003 , 38, 606-12 | 3.5 | 82 |
| 39 | Functional significance of a hereditary adenine insertion variant in the 5QTR of the endothelin-1 gene. <i>Pharmacogenetics and Genomics</i> , 2003 , 13, 445-51 | | 34 |
| 38 | Somatoform pain disorder in the general population. <i>Psychotherapy and Psychosomatics</i> , 2003 , 72, 88-94 | 9.4 | 37 |

| | | | |
|----|--|-----|-----|
| 37 | Distribution and dose response of laboratory markers to alcohol consumption in a general population: results of the study of health in Pomerania (SHIP). <i>Journal of Studies on Alcohol and Drugs</i> , 2003 , 64, 75-82 | | 33 |
| 36 | Measuring sense of coherence with only three items: a useful tool for population surveys. <i>British Journal of Health Psychology</i> , 2003 , 8, 409-21 | 8.3 | 43 |
| 35 | Probabilities of alcohol high-risk drinking, abuse or dependence estimated on grounds of tobacco smoking and nicotine dependence. <i>Addiction</i> , 2003 , 98, 805-14 | 4.6 | 83 |
| 34 | Intentionally reduced smoking among untreated general population smokers: prevalence, stability, prediction of smoking behaviour change and differences between subjects choosing either reduction or abstinence. <i>Addiction</i> , 2003 , 98, 1101-10 | 4.6 | 31 |
| 33 | Late onset alcoholism. <i>European Psychiatry</i> , 2003 , 18, 112-8 | 6 | 35 |
| 32 | The Fagerström test for nicotine dependence in two adult population samples-potential influence of lifetime amount of tobacco smoked on the degree of dependence. <i>Drug and Alcohol Dependence</i> , 2003 , 71, 1-6 | 4.9 | 27 |
| 31 | Relation among stage of change, demographic characteristics, smoking history, and nicotine dependence in an adult German population. <i>Preventive Medicine</i> , 2003 , 37, 368-74 | 4.3 | 40 |
| 30 | Alcohol high risk drinking, abuse and dependence among tobacco smoking medical care patients and the general population. <i>Drug and Alcohol Dependence</i> , 2003 , 69, 189-95 | 4.9 | 43 |
| 29 | Specific somatoform disorder in the general population. <i>Psychosomatics</i> , 2003 , 44, 304-11 | 2.6 | 40 |
| 28 | The prevalence of undiagnosed thyroid disorders in a previously iodine-deficient area. <i>Thyroid</i> , 2003 , 13, 803-10 | 6.2 | 184 |
| 27 | Tobacco- and alcohol-attributable mortality and years of potential life lost in Germany. <i>European Journal of Public Health</i> , 2003 , 13, 275-7 | 2.1 | 17 |
| 26 | Skalen zu Kernkonstrukten des Transtheoretischen Modells (TTM) im Verhaltensbereich Tabakkonsum. <i>Zeitschrift Fur Gesundheitspsychologie</i> , 2003 , 11, 57-66 | | 7 |
| 25 | Study of Health in Pomerania (SHIP): a health survey in an East German region. Objectives and design of the oral health section. <i>Quintessence International</i> , 2003 , 34, 370-8 | | 30 |
| 24 | Prosthodontic status among old adults in Pomerania, related to income, education level, and general health (results of the Study of Health in Pomerania, SHIP). <i>International Journal of Prosthodontics</i> , 2003 , 16, 313-8 | 1.9 | 11 |
| 23 | The role of family and partnership in recovery from alcohol dependence: comparison of individuals remitting with and without formal help. <i>European Addiction Research</i> , 2002 , 8, 122-7 | 4.6 | 24 |
| 22 | The readiness to change questionnaire in subjects with hazardous alcohol consumption, alcohol misuse and dependence in a general population survey. <i>Alcohol and Alcoholism</i> , 2002 , 37, 362-9 | 3.5 | 28 |
| 21 | Screening for alcohol use disorders and at-risk drinking in the general population: psychometric performance of three questionnaires. <i>Alcohol and Alcoholism</i> , 2002 , 37, 261-8 | 3.5 | 198 |
| 20 | Association between behavior-dependent cardiovascular risk factors and asymptomatic carotid atherosclerosis in a general population. <i>Stroke</i> , 2002 , 33, 2929-35 | 6.7 | 78 |

| | | | |
|----|--|-----|-------|
| 19 | Alcohol-attributable mortality in a high per capita consumption country -- Germany. <i>Alcohol and Alcoholism</i> , 2002 , 37, 581-5 | 3.5 | 32 |
| 18 | Remission from alcohol dependence without help: how restrictive should our definition of treatment be?. <i>Journal of Studies on Alcohol and Drugs</i> , 2002 , 63, 229-36 | | 20 |
| 17 | The importance of a positive family history of alcoholism, parental rejection and emotional warmth, behavioral problems and peer substance use for alcohol problems in teenagers: a path analysis. <i>Journal of Studies on Alcohol and Drugs</i> , 2002 , 63, 305-15 | | 105 |
| 16 | Naturalistic changes in the readiness to quit tobacco smoking in a German general population sample. <i>Preventive Medicine</i> , 2002 , 35, 326-33 | 4.3 | 21 |
| 15 | Study of Health In Pomerania (SHIP): a health examination survey in an east German region: objectives and design. <i>International Journal of Public Health</i> , 2001 , 46, 186-94 | | 395 |
| 14 | Lifetime-comorbidity of obsessive-compulsive disorder and subclinical obsessive-compulsive disorder in Northern Germany. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2001 , 251, 130-5 | 5.1 | 62 |
| 13 | Screening for mental health: validity of the MHI-5 using DSM-IV Axis I psychiatric disorders as gold standard. <i>Psychiatry Research</i> , 2001 , 105, 243-53 | 9.9 | 394 |
| 12 | The Composite International Diagnostic Interview: feasibility and necessity of editing and interviewer training in general population surveys. <i>International Journal of Methods in Psychiatric Research</i> , 2000 , 9, 32-42 | 4.3 | 19 |
| 11 | Psychological problems of partnership and profit or: who has the higher population impact?. <i>Addiction</i> , 2000 , 95, 1436-7 | 4.6 | 1 |
| 10 | Maintenance Factors of Recovery From Alcohol Dependence in Treated and Untreated Individuals. <i>Alcoholism: Clinical and Experimental Research</i> , 2000 , 24, 1773-1777 | 3.7 | 24 |
| 9 | Inanspruchnahme suchtspezifischer Hilfen von Alkoholabhängigen und -mißbrauchern: Ergebnisse der TACOS-Bevölkerungsstudie. <i>Sucht</i> , 2000 , 46, 9-17 | 0.5 | 31 |
| 8 | Inanspruchnahme von Hilfen zur Erlangung der Nikotin-Abstinenz. <i>Sucht</i> , 2000 , 46, 398-407 | 0.5 | 10 |
| 7 | Estimating prevalence of alcohol abuse and dependence in one general hospital: an approach to reduce sample selection bias. <i>Alcohol and Alcoholism</i> , 1999 , 34, 786-94 | 3.5 | 21 |
| 6 | Motivation to change drinking behavior: comparison of alcohol-dependent individuals in a general hospital and a general population sample. <i>General Hospital Psychiatry</i> , 1999 , 21, 348-53 | 5.6 | 38 |
| 5 | Prevalence of Alcohol Dependence and Abuse in General Practice. <i>Alcoholism: Clinical and Experimental Research</i> , 1998 , 22, 935-940 | 3.7 | 18 |
| 4 | Previous help seeking and motivation to change drinking behavior in alcohol-dependent general hospital patients. <i>General Hospital Psychiatry</i> , 1998 , 20, 115-9 | 5.6 | 16 |
| 3 | Development of a Screening Questionnaire for the General Hospital and General Practices. <i>Alcoholism: Clinical and Experimental Research</i> , 1997 , 21, 894-898 | 3.7 | 74 |
| 2 | Förderung der Tabakabstinenz durch neue Kommunikationsmedien und Expertensysteme | | 39-49 |

- 1 Die Adhärenz Erwachsener an der Bewegungsaufzeichnung über 7 Tage mittels Akzelerometer.
Pravention Und Gesundheitsforderung,1 0.5