

Antonios Stavropoulos-Kalinoglou

List of Publications by Year in Descending Order

Source:

<https://exaly.com/author-pdf/7462924/antonios-stavropoulos-kalinoglou-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

50
papers

2,630
citations

29
h-index

51
g-index

52
ext. papers

2,951
ext. citations

3.3
avg, IF

4.61
L-index

#	Paper	IF	Citations
50	Position Statement on Exercise Dosage in Rheumatic and Musculoskeletal Diseases: The Role of the IMPACT-RMD Toolkit. <i>Mediterranean Journal of Rheumatology</i> , 2021 , 32, 378-385	1.4	2
49	Participation in physical activity decreased more in people with rheumatoid arthritis than the general population during the COVID-19 lockdown: a cross-sectional study. <i>Rheumatology International</i> , 2021 , 1	3.6	2
48	The effects of training with high-speed interval running on muscle performance are modulated by slope. <i>Physiological Reports</i> , 2021 , 9, e14656	2.6	1
47	Smartphone pedometers in adults with asthma: a practical approach to physical activity assessment? A pilot validation study. <i>Journal of Asthma</i> , 2021 , 1-9	1.9	0
46	Omega-3 polyunsaturated fatty acid supplementation versus placebo on vascular health, glycaemic control, and metabolic parameters in people with type 1 diabetes: a randomised controlled preliminary trial. <i>Cardiovascular Diabetology</i> , 2020 , 19, 127	8.7	9
45	ACSM Preparticipation Health Screening Guidelines: A UK University Cohort Perspective. <i>Medicine and Science in Sports and Exercise</i> , 2019 , 51, 1047-1054	1.2	6
44	Could IL-6 inhibition prevent exercise-induced fat loss in RA?. <i>Nature Reviews Rheumatology</i> , 2019 , 15, 192-194	8.1	
43	Comparison of the effects of exercise and anti-TNF treatment on cardiovascular health in rheumatoid arthritis: results from two controlled trials. <i>Rheumatology International</i> , 2019 , 39, 219-225	3.6	10
42	The role of exercise in the management of rheumatoid arthritis. <i>Expert Review of Clinical Immunology</i> , 2015 , 11, 1121-30	5.1	42
41	Cardiorespiratory fitness levels and their association with cardiovascular profile in patients with rheumatoid arthritis: a cross-sectional study. <i>Rheumatology</i> , 2015 , 54, 2215-20	3.9	27
40	Adiponectin, Resistin, and Visfatin in Childhood Obesity and Exercise. <i>Pediatric Exercise Science</i> , 2015 , 27, 454-62	2	17
39	Prolonged QTc interval predicts all-cause mortality in patients with rheumatoid arthritis: an association driven by high inflammatory burden. <i>Rheumatology</i> , 2014 , 53, 131-7	3.9	59
38	Three months of moderate-intensity exercise reduced plasma 3-nitrotyrosine in rheumatoid arthritis patients. <i>European Journal of Applied Physiology</i> , 2014 , 114, 1483-92	3.4	29
37	Individualised exercise improves endothelial function in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 748-51	2.4	80
36	Muscle wasting in rheumatoid arthritis: The role of oxidative stress. <i>World Journal of Rheumatology</i> , 2014 , 4, 44	0.5	4
35	Predictors of asymmetric dimethylarginine levels in patients with rheumatoid arthritis: the role of insulin resistance. <i>Scandinavian Journal of Rheumatology</i> , 2013 , 42, 176-81	1.9	16
34	Individualised aerobic and resistance exercise training improves cardiorespiratory fitness and reduces cardiovascular risk in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 1819-25	2.4	155

33	Anti-tumour necrosis factor alpha therapy improves insulin sensitivity in normal-weight but not in obese patients with rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2012 , 14, R160	5.7	46
32	Obesity and Arthritis 2012 , 355-380		
31	Disease activity and low physical activity associate with number of hospital admissions and length of hospitalisation in patients with rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2011 , 13, R108	5.7	37
30	Platelet function in rheumatoid arthritis: arthritic and cardiovascular implications. <i>Rheumatology International</i> , 2011 , 31, 153-64	3.6	105
29	Inverted BMI rather than BMI is a better proxy for percentage of body fat. <i>Annals of Human Biology</i> , 2011 , 38, 681-4	1.7	25
28	Obesity in rheumatoid arthritis. <i>Rheumatology</i> , 2011 , 50, 450-62	3.9	144
27	Rheumatoid arthritis susceptibility genes associate with lipid levels in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 1025-32	2.4	44
26	Anti-TNF α therapy may lead to blood pressure reductions through improved endothelium-dependent microvascular function in patients with rheumatoid arthritis. <i>Journal of Human Hypertension</i> , 2011 , 25, 699-702	2.6	32
25	What predicts obesity in patients with rheumatoid arthritis? An investigation of the interactions between lifestyle and inflammation. <i>International Journal of Obesity</i> , 2010 , 34, 295-301	5.5	27
24	The rationale for comparative studies of accelerated atherosclerosis in rheumatic diseases. <i>Current Vascular Pharmacology</i> , 2010 , 8, 437-49	3.3	75
23	Association of mean platelet volume with hypertension in rheumatoid arthritis. <i>Inflammation and Allergy: Drug Targets</i> , 2010 , 9, 45-50		61
22	Vascular function and inflammation in rheumatoid arthritis: the role of physical activity. <i>Open Cardiovascular Medicine Journal</i> , 2010 , 4, 89-96	0.7	32
21	Target organ damage in patients with rheumatoid arthritis: the role of blood pressure and heart rate. <i>Atherosclerosis</i> , 2010 , 209, 255-60	3.1	27
20	Rheumatoid cachexia and cardiovascular disease. <i>Nature Reviews Rheumatology</i> , 2010 , 6, 445-51	8.1	98
19	Mean platelet volume in patients with rheumatoid arthritis: the effect of anti-TNF α therapy. <i>Rheumatology International</i> , 2010 , 30, 1125-9	3.6	131
18	The Rationale for Comparative Studies of Accelerated Atherosclerosis in Rheumatic Diseases. <i>Current Vascular Pharmacology</i> , 2010 , 999, 1-13	3.3	3
17	Association of physical inactivity with increased cardiovascular risk in patients with rheumatoid arthritis. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2009 , 16, 188-94		125
16	Associations of obesity with modifiable risk factors for the development of cardiovascular disease in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2009 , 68, 242-5	2.4	68

15	Underweight and obese states both associate with worse disease activity and physical function in patients with established rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2009 , 28, 439-44	3.9	66
14	Lack of an association of GNB3 C825T polymorphism and blood pressure in patients with rheumatoid arthritis. <i>Clinical and Experimental Hypertension</i> , 2009 , 31, 428-39	2.2	2
13	Association of interleukin-6 (IL-6)-174G/C gene polymorphism with cardiovascular disease in patients with rheumatoid arthritis: the role of obesity and smoking. <i>Atherosclerosis</i> , 2009 , 204, 178-83	3.1	82
12	Transforming growth factor-beta1 869T/C, but not interleukin-6 -174G/C, polymorphism associates with hypertension in rheumatoid arthritis. <i>Rheumatology</i> , 2009 , 48, 113-8	3.9	35
11	Cigarette smoking associates with body weight and muscle mass of patients with rheumatoid arthritis: a cross-sectional, observational study. <i>Arthritis Research and Therapy</i> , 2008 , 10, R59	5.7	28
10	Cigarette smoking significantly increases basal metabolic rate in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2008 , 67, 70-3	2.4	38
9	Rheumatoid arthritis, cardiovascular disease and physical exercise: a systematic review. <i>Rheumatology</i> , 2008 , 47, 239-48	3.9	152
8	Polymorphisms of the endothelin-1 gene associate with hypertension in patients with rheumatoid arthritis. <i>Endothelium: Journal of Endothelial Cell Research</i> , 2008 , 15, 203-12		45
7	Serum uric acid is independently associated with hypertension in patients with rheumatoid arthritis. <i>Journal of Human Hypertension</i> , 2008 , 22, 177-82	2.6	30
6	Metabolism in Patients with Rheumatoid Arthritis: Resting Energy Expenditure, Physical Activity and Diet-Induced Thermogenesis. Invited Review. <i>Recent Patents on Endocrine, Metabolic & Immune Drug Discovery</i> , 2008 , 2, 97-102		2
5	Long-term exposure to medium-dose glucocorticoid therapy associates with hypertension in patients with rheumatoid arthritis. <i>Rheumatology</i> , 2008 , 47, 72-5	3.9	134
4	New resting energy expenditure prediction equations for patients with rheumatoid arthritis. <i>Rheumatology</i> , 2008 , 47, 500-6	3.9	33
3	Blockade of tumour necrosis factor-alpha in rheumatoid arthritis: effects on components of rheumatoid cachexia. <i>Rheumatology</i> , 2007 , 46, 1824-7	3.9	116
2	Prevalence and associations of hypertension and its control in patients with rheumatoid arthritis. <i>Rheumatology</i> , 2007 , 46, 1477-82	3.9	211
1	Redefining overweight and obesity in rheumatoid arthritis patients. <i>Annals of the Rheumatic Diseases</i> , 2007 , 66, 1316-21	2.4	117