

# Laura Leyva

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7478143/publications.pdf>

Version: 2024-02-01

71  
papers

2,488  
citations

172207

29  
h-index

205818

48  
g-index

71  
all docs

71  
docs citations

71  
times ranked

3921  
citing authors

#	ARTICLE	IF	CITATIONS
1	Identification of the genetic mechanism that associates <i>L3MBTL3</i> to multiple sclerosis. <i>Human Molecular Genetics</i> , 2022, 31, 2155-2163.	1.4	4
2	Treatment of faecal incontinence with autologous expanded mesenchymal stem cells: results of a pilot study. <i>Colorectal Disease</i> , 2021, 23, 698-709.	0.7	11
3	Soluble Receptor Isoform of IFN-Beta (sIFNAR2) in Multiple Sclerosis Patients and Their Association With the Clinical Response to IFN-Beta Treatment. <i>Frontiers in Immunology</i> , 2021, 12, 778204.	2.2	5
4	A New Risk Variant for Multiple Sclerosis at 11q23.3 Locus Is Associated with Expansion of CXCR5+ Circulating Regulatory T Cells. <i>Journal of Clinical Medicine</i> , 2020, 9, 625.	1.0	5
5	Antiviral, Immunomodulatory and Antiproliferative Activities of Recombinant Soluble IFNAR2 without IFN- $\gamma$ Mediation. <i>Journal of Clinical Medicine</i> , 2020, 9, 959.	1.0	4
6	Exome sequencing in multiple sclerosis families identifies 12 candidate genes and nominates biological pathways for the genesis of disease. <i>PLoS Genetics</i> , 2019, 15, e1008180.	1.5	46
7	Adipose-derived mesenchymal stem cells (AdMSC) for the treatment of secondary-progressive multiple sclerosis: A triple blinded, placebo controlled, randomized phase I/II safety and feasibility study. <i>PLoS ONE</i> , 2018, 13, e0195891.	1.1	112
8	Global methylation correlates with clinical status in multiple sclerosis patients in the first year of IFNbeta treatment. <i>Scientific Reports</i> , 2017, 7, 8727.	1.6	17
9	Cross-reactivity of antibodies against interferon beta in multiple sclerosis patients and interference of the JAK-STAT signaling pathway. <i>Scientific Reports</i> , 2017, 7, 16585.	1.6	7
10	Gene therapy with mesenchymal stem cells expressing IFN- $\gamma$ ameliorates neuroinflammation in experimental models of multiple sclerosis. <i>British Journal of Pharmacology</i> , 2017, 174, 238-253.	2.7	34
11	Decreased soluble IFN- $\gamma$ receptor (sIFNAR2) in multiple sclerosis patients: A potential serum diagnostic biomarker. <i>Multiple Sclerosis Journal</i> , 2017, 23, 937-945.	1.4	12
12	Activation of the JAK-STAT Signaling Pathway after In Vitro Stimulation with IFN- $\gamma$ in Multiple Sclerosis Patients According to the Therapeutic Response to IFN- $\gamma$ . <i>PLoS ONE</i> , 2017, 12, e0170031.	1.1	15
13	The HLA DRB1*03:01 allele is associated with NMO regardless of the NMO-IgG status in Brazilian patients from Rio de Janeiro. <i>Journal of Neuroimmunology</i> , 2017, 310, 1-7.	1.1	22
14	TRAIL and TRAIL receptors splice variants during long-term interferon $\gamma$ treatment of patients with multiple sclerosis: evaluation as biomarkers for therapeutic response. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2016, 87, jnnp-2014-309932.	0.9	6
15	Analysis of Plasminogen Genetic Variants in Multiple Sclerosis Patients. <i>G3: Genes, Genomes, Genetics</i> , 2016, 6, 2073-2079.	0.8	13
16	Cell-based product classification procedure: What can be done differently to improve decisions on borderline products?. <i>Cytherapy</i> , 2016, 18, 809-815.	0.3	12
17	Recombinant soluble IFN receptor (sIFNAR2) exhibits intrinsic therapeutic efficacy in a murine model of Multiple Sclerosis. <i>Neuropharmacology</i> , 2016, 110, 480-492.	2.0	5
18	Development and validation of an ELISA for quantification of soluble IFN- $\gamma$ receptor: assessment in multiple sclerosis. <i>Bioanalysis</i> , 2015, 7, 2869-2880.	0.6	6

#	ARTICLE	IF	CITATIONS
19	Pharmacogenomic study in patients with multiple sclerosis. <i>Neurology: Neuroimmunology and NeuroInflammation</i> , 2015, 2, e154.	3.1	19
20	Lipid-specific immunoglobulin G bands in cerebrospinal fluid are associated with a reduced risk of developing progressive multifocal leukoencephalopathy during treatment with natalizumab. <i>Annals of Neurology</i> , 2015, 77, 447-457.	2.8	48
21	A functional variant that affects exon-skipping and protein expression of <i>SP140</i> as genetic mechanism predisposing to multiple sclerosis. <i>Human Molecular Genetics</i> , 2015, 24, 5619-5627.	1.4	43
22	Genome-wide significant association with seven novel multiple sclerosis risk loci. <i>Journal of Medical Genetics</i> , 2015, 52, 848-855.	1.5	34
23	A new risk variant for multiple sclerosis at the immunoglobulin heavy chain locus associates with intrathecal IgG, IgM index and oligoclonal bands. <i>Multiple Sclerosis Journal</i> , 2015, 21, 1104-1111.	1.4	12
24	Mesenchymal properties of SJL mice-stem cells and their efficacy as autologous therapy in a relapsing-remitting multiple sclerosis model. <i>Stem Cell Research and Therapy</i> , 2014, 5, 134.	2.4	12
25	HLA alleles as biomarkers of high-titre neutralising antibodies to interferon- $\beta$ therapy in multiple sclerosis. <i>Journal of Medical Genetics</i> , 2014, 51, 395-400.	1.5	19
26	Killer-Cell Immunoglobulin-Like Receptor Expression on Lymphocyte Subsets in Multiple Sclerosis Patients Treated with Interferon- $\beta$ : Evaluation as Biomarkers for Clinical Response. <i>CNS Drugs</i> , 2014, 28, 559-570.	2.7	2
27	Human Endogenous Retrovirus HERV-Fc1 Association with Multiple Sclerosis Susceptibility: A Meta-Analysis. <i>PLoS ONE</i> , 2014, 9, e90182.	1.1	29
28	Early development of anti-natalizumab antibodies in MS patients. <i>Journal of Neurology</i> , 2013, 260, 2343-2347.	1.8	11
29	Identification of a functional variant in the <i>KIF5A-CYP27B1-METTL1-FAM119B</i> locus associated with multiple sclerosis. <i>Journal of Medical Genetics</i> , 2013, 50, 25-33.	1.5	59
30	Candidate Gene Study of TRAIL and TRAIL Receptors: Association with Response to Interferon Beta Therapy in Multiple Sclerosis Patients. <i>PLoS ONE</i> , 2013, 8, e62540.	1.1	18
31	ANKRD55 and DHCR7 are novel multiple sclerosis risk loci. <i>Genes and Immunity</i> , 2012, 13, 253-257.	2.2	44
32	The CD4+ T-cell subset lacking expression of the CD28 costimulatory molecule is expanded and shows a higher activation state in multiple sclerosis. <i>Journal of Neuroimmunology</i> , 2012, 243, 1-11.	1.1	22
33	DRB1*03:01 Haplotypes: Differential Contribution to Multiple Sclerosis Risk and Specific Association with the Presence of Intrathecal IgM Bands. <i>PLoS ONE</i> , 2012, 7, e31018.	1.1	11
34	Genome-Wide Association Study of Multiple Sclerosis Confirms a Novel Locus at 5p13.1. <i>PLoS ONE</i> , 2012, 7, e36140.	1.1	46
35	Predictors of Fatigue Severity in Early Systemic Sclerosis: A Prospective Longitudinal Study of the GENISOS Cohort. <i>PLoS ONE</i> , 2011, 6, e26061.	1.1	30
36	Replication of top markers of a genome-wide association study in multiple sclerosis in Spain. <i>Genes and Immunity</i> , 2011, 12, 110-115.	2.2	36

#	ARTICLE	IF	CITATIONS
37	Killer cell immunoglobulin-like receptor genes in Spanish multiple sclerosis patients. <i>Molecular Immunology</i> , 2011, 48, 1896-1902.	1.0	33
38	Gene expression in IFN $\gamma$ signalling pathway differs between monocytes, CD4 and CD8 T cells from MS patients. <i>Journal of Neuroimmunology</i> , 2011, 230, 153-159.	1.1	15
39	IL28B polymorphisms are not associated with the response to interferon-beta in multiple sclerosis. <i>Journal of Neuroimmunology</i> , 2011, 239, 101-104.	1.1	18
40	The efficacy of natalizumab in patients with multiple sclerosis according to level of disability: results of an observational study. <i>Multiple Sclerosis Journal</i> , 2011, 17, 192-197.	1.4	26
41	Kinetics and incidence of anti-natalizumab antibodies in multiple sclerosis patients on treatment for 18 months. <i>Multiple Sclerosis Journal</i> , 2011, 17, 368-371.	1.4	36
42	TRAIL/TRAIL Receptor System and Susceptibility to Multiple Sclerosis. <i>PLoS ONE</i> , 2011, 6, e21766.	1.1	16
43	Hexose-6-phosphate dehydrogenase: a new risk gene for multiple sclerosis. <i>European Journal of Human Genetics</i> , 2010, 18, 618-620.	1.4	9
44	Tag-SNP analysis of the GF11-EVI5-RPL5-FAM69 risk locus for multiple sclerosis. <i>European Journal of Human Genetics</i> , 2010, 18, 827-831.	1.4	25
45	The autoimmune disease-associated KIF5A, CD226 and SH2B3 gene variants confer susceptibility for multiple sclerosis. <i>Genes and Immunity</i> , 2010, 11, 439-445.	2.2	79
46	Does the DRB1 $\ast$ 1501 allele confer more severe and faster progression in primary progressive multiple sclerosis patients? HLA in primary progressive multiple sclerosis. <i>Journal of Neuroimmunology</i> , 2009, 214, 101-103.	1.1	22
47	Neutralizing antibodies against IFN beta in patients with multiple sclerosis: A comparative study of two cytopathic effect tests (CPE) for their detection. <i>Journal of Immunological Methods</i> , 2009, 351, 41-45.	0.6	11
48	HLA class II alleles in patients with multiple sclerosis in the Biscay province (Basque Country, Spain). <i>Journal of Neurology</i> , 2009, 256, 1977-1988.	1.8	25
49	Multiple sclerosis association study with the <i>TENR</i> region in a Spanish population. <i>Tissue Antigens</i> , 2009, 74, 244-247.	1.0	20
50	IL2RA/CD25 Gene Polymorphisms: Uneven Association with Multiple Sclerosis (MS) and Type 1 Diabetes (T1D). <i>PLoS ONE</i> , 2009, 4, e4137.	1.1	65
51	Multiple sclerosis in Gypsies from southern Spain: prevalence, mitochondrial DNA haplogroups and HLA class II association. <i>Tissue Antigens</i> , 2008, 71, 426-433.	1.0	16
52	The T244I variant of the interleukin $\gamma$ receptor $\alpha$ gene and multiple sclerosis. <i>Tissue Antigens</i> , 2008, 72, 158-161.	1.0	30
53	The high producer variant of the Fc-receptor like-3 (FCRL3) gene is involved in protection against multiple sclerosis. <i>Journal of Neuroimmunology</i> , 2008, 195, 146-150.	1.1	37
54	Interferon regulatory factor 5 (IRF5) gene variants are associated with multiple sclerosis in three distinct populations. <i>Journal of Medical Genetics</i> , 2008, 45, 362-369.	1.5	128

#	ARTICLE	IF	CITATIONS
55	Interferon receptor expression in multiple sclerosis patients. <i>Journal of Neuroimmunology</i> , 2007, 183, 225-231.	1.1	22
56	HLA class II and response to interferon-beta in multiple sclerosis. <i>Acta Neurologica Scandinavica</i> , 2005, 112, 391-394.	1.0	39
57	IFNAR1 and IFNAR2 polymorphisms confer susceptibility to multiple sclerosis but not to interferon-beta treatment response. <i>Journal of Neuroimmunology</i> , 2005, 163, 165-171.	1.1	85
58	Protein tyrosine phosphatase gene (PTPN22) polymorphism in multiple sclerosis. <i>Journal of Neurology</i> , 2005, 252, 994-995.	1.8	38
59	Effects of the multiple sclerosis associated $\gamma$ 330 promoter polymorphism in IL2 allelic expression. <i>Journal of Neuroimmunology</i> , 2004, 148, 212-217.	1.1	76
60	Differences in the immunological responses in drug- and virus-induced cutaneous reactions in children. <i>Blood Cells, Molecules, and Diseases</i> , 2003, 30, 124-131.	0.6	36
61	Memory to the hapten in non-immediate cutaneous allergic reactions to betalactams resides in a lymphocyte subpopulation expressing both CD45RO and CLA markers. <i>Blood Cells, Molecules, and Diseases</i> , 2003, 31, 75-79.	0.6	6
62	Delayed reactions to drugs show levels of perforin, granzyme B, and Fas-L to be related to disease severity. <i>Journal of Allergy and Clinical Immunology</i> , 2002, 109, 155-161.	1.5	201
63	Characterization of specific IgE response in vitro against protein and drug allergens using atopic and normal donors. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2002, 57, 193-200.	2.7	4
64	Controlled administration of penicillin to patients with a positive history but negative skin and specific serum IgE tests. <i>Clinical and Experimental Allergy</i> , 2002, 32, 270-276.	1.4	77
65	Study of binding and neutralising antibodies to interferon- $\gamma$ in two groups of relapsing-remitting multiple sclerosis patients. <i>Journal of Neurology</i> , 2001, 248, 383-388.	1.8	32
66	In vitro T-cell responses to beta-lactam drugs in immediate and nonimmediate allergic reactions. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2001, 56, 611-618.	2.7	163
67	Expression of the skin-homing receptor in peripheral blood lymphocytes from subjects with nonimmediate cutaneous allergic drug reactions. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2000, 55, 998-1004.	2.7	67
68	Association of hypersensitivity to the nematode <i>Anisakis simplex</i> with HLA class II DRB1 $\alpha$ -1502-DQB1 $\alpha$ -0601 haplotype. <i>Human Immunology</i> , 2000, 61, 314-319.	1.2	24
69	Subjects with allergic reactions to drugs show in vivo polarized patterns of cytokine expression depending on the chronology of the clinical reaction. <i>Journal of Allergy and Clinical Immunology</i> , 2000, 106, 769-776.	1.5	77
70	Anticonvulsant-induced toxic epidermal necrolysis: Monitoring the immunologic response. <i>Journal of Allergy and Clinical Immunology</i> , 2000, 105, 157-165.	1.5	94
71	Preferential Expression of the Skin-Homing Receptor CLA in Peripheral T Lymphocytes from Patients with Drug-Induced Allergic Reactions. <i>International Archives of Allergy and Immunology</i> , 1999, 118, 355-357.	0.9	5