

Setsuya Aiba

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7483489/setsuya-aiba-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

61
papers

1,265
citations

18
h-index

35
g-index

61
ext. papers

1,451
ext. citations

3.4
avg, IF

4.27
L-index

#	Paper	IF	Citations
61	GWAS Identified IL4R and the Major Histocompatibility Complex Region as the Associated Loci of Total Serum IgE Levels in 9,260 Japanese Individuals. <i>Journal of Investigative Dermatology</i> , 2021 , 141, 2749-2752	4.3	1
60	Successful treatment of BRAF/MEK inhibitor-resistant advanced cutaneous melanoma with nivolumab plus ipilimumab combination therapy followed by intensity-modulated radiotherapy. <i>Journal of Dermatology</i> , 2021 , 48, 1419-1422	1.6	0
59	Optimization of the IL-2 Luc assay for immunosuppressive drugs: a novel in vitro immunotoxicity test with high sensitivity and predictivity. <i>Archives of Toxicology</i> , 2021 , 95, 2755-2768	5.8	1
58	Case series of BRAF-mutated advanced melanoma treated with encorafenib plus binimetinib combination therapy. <i>Journal of Dermatology</i> , 2021 , 48, 397-400	1.6	2
57	The modified IL-8 Luc assay, an in vitro skin sensitisation test, can significantly improve the false-negative judgment of lipophilic sensitizers with logK values > 3.5. <i>Archives of Toxicology</i> , 2021 , 95, 749-758	5.8	3
56	The IL-1 β promoter-driven luciferase reporter cell line THP-G1b can efficiently predict skin-sensitising chemicals. <i>Archives of Toxicology</i> , 2021 , 95, 1647-1657	5.8	
55	Ehlers-Danlos syndrome type IV with a novel COL3A1 exon 14 skipping variation confirmed by Tohoku Medical Megabank Organization genomic database. <i>Journal of Dermatology</i> , 2021 , 48, 1918-1922 ^{1.6}		
54	An international validation study of the IL-2 Luc assay for evaluating the potential immunotoxic effects of chemicals on T cells and a proposal for reference data for immunotoxic chemicals. <i>Toxicology in Vitro</i> , 2020 , 66, 104832	3.6	3
53	Metabolic and pathologic profiles of human LSS deficiency recapitulated in mice. <i>PLoS Genetics</i> , 2020 , 16, e1008628	6	5
52	Development of a novel in vitro assay to evaluate environmental water using an IL-8 reporter cell line. <i>EXCLI Journal</i> , 2020 , 19, 1054-1063	2.4	
51	Numerical Simulation of Effects of Bioheat Transfer Characteristics of Malignant Melanoma on Thermal Conductivity Measurements. <i>Critical Reviews in Biomedical Engineering</i> , 2020 , 48, 95-109	1.1	
50	TLR3 augments glucocorticoid-synthetic enzymes expression in epidermal keratinocytes; Implications of glucocorticoid metabolism in rosacea epidermis. <i>Journal of Dermatological Science</i> , 2020 , 100, 58-66	4.3	3
49	Severe pyrexia from nivolumab-resistant advanced melanoma after successful combined therapy with encorafenib plus binimetinib. <i>Journal of Dermatology</i> , 2020 , 47, 654-657	1.6	4
48	Metabolic and pathologic profiles of human LSS deficiency recapitulated in mice 2020 , 16, e1008628		
47	Metabolic and pathologic profiles of human LSS deficiency recapitulated in mice 2020 , 16, e1008628		
46	Metabolic and pathologic profiles of human LSS deficiency recapitulated in mice 2020 , 16, e1008628		
45	Metabolic and pathologic profiles of human LSS deficiency recapitulated in mice 2020 , 16, e1008628		

44	Metabolic and pathologic profiles of human LSS deficiency recapitulated in mice 2020 , 16, e1008628		
43	Metabolic and pathologic profiles of human LSS deficiency recapitulated in mice 2020 , 16, e1008628		
42	Significance of BRAF Kinase Inhibitors for Melanoma Treatment: From Bench to Bedside. <i>Cancers</i> , 2019 , 11,	6.6	15
41	Severe rhabdomyolysis developing in an advanced melanoma patient treated by pembrolizumab followed by dabrafenib trametinib combined therapy. <i>Journal of Dermatology</i> , 2019 , 46, e256-e258	1.6	5
40	First-in-human clinical study of novel technique to diagnose malignant melanoma via thermal conductivity measurements. <i>Scientific Reports</i> , 2019 , 9, 3853	4.9	9
39	A novel technique to diagnose non-melanoma skin cancer by thermal conductivity measurements: Correlations with cancer stromal factors. <i>Experimental Dermatology</i> , 2019 , 28, 1029-1035	4	5
38	TLR3 stimulation induces melanosome endo/phagocytosis through RHOA and CDC42 in human epidermal keratinocyte. <i>Journal of Dermatological Science</i> , 2019 , 96, 168-177	4.3	7
37	BRAF kinase inhibitors for treatment of melanoma: developments from early-stage animal studies to Phase II clinical trials. <i>Expert Opinion on Investigational Drugs</i> , 2019 , 28, 143-148	5.9	12
36	Eruptive xanthomas in a patient with soft-drink diabetic ketosis and apolipoprotein E4/2. <i>Endocrine Journal</i> , 2019 , 66, 107-114	2.9	2
35	HLA-DRB1*04:05 in two cases of Vogt-Koyanagi-Harada disease-like uveitis developing from an advanced melanoma patient treated by sequential administration of nivolumab and dabrafenib/trametinib therapy. <i>Journal of Dermatology</i> , 2018 , 45, 735-737	1.6	38
34	Profiling the immunotoxicity of chemicals based on in vitro evaluation by a combination of the Multi-ImmunoTox assay and the IL-8 Luc assay. <i>Archives of Toxicology</i> , 2018 , 92, 2043-2054	5.8	8
33	Toll-like receptors 2 and 3 enhance melanogenesis and melanosome transport in human melanocytes. <i>Pigment Cell and Melanoma Research</i> , 2018 , 31, 570-584	4.5	16
32	Tumor-Associated Macrophages: Therapeutic Targets for Skin Cancer. <i>Frontiers in Oncology</i> , 2018 , 8, 3	5.3	73
31	Pharmacokinetic Study of Bioactive Flavonoids in the Traditional Japanese Medicine Keigairengyoto Exerting Antibacterial Effects against Staphylococcus aureus. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	13
30	The performance of an in vitro skin sensitisation test, IL-8 Luc assay (OECD442E), and the integrated approach with direct peptide reactive assay (DPRA). <i>Journal of Toxicological Sciences</i> , 2018 , 43, 741-749	1.9	6
29	Keratoacanthoma, palmoplantar keratoderma developing in an advanced melanoma patient treated with vemurafenib regressed by blockade of mitogen-activated protein kinase signaling. <i>Journal of Dermatology</i> , 2017 , 44, e226-e227	1.6	10
28	Glucuronides of phytoestrogen flavonoid enhance macrophage function via conversion to aglycones by β glucuronidase in macrophages. <i>Immunity, Inflammation and Disease</i> , 2017 , 5, 265-279	2.4	18
27	In vitro test methods to evaluate the effects of chemicals on innate and adaptive immune responses. <i>Current Opinion in Toxicology</i> , 2017 , 5, 6-12	4.4	2

26	Epidermal iron metabolism for iron salvage. <i>Journal of Dermatological Science</i> , 2017 , 87, 101-109	4.3	5
25	Inhibition of Human Kallikrein 5 Protease by Triterpenoids from Natural Sources. <i>Molecules</i> , 2017 , 22,	4.8	11
24	Keigairengyoto, a traditional Japanese medicine, promotes bacterial clearance by activating innate immune cells in mouse cutaneous infection models 2017 , 1, 35		2
23	The possible interaction between periostin expressed by cancer stroma and tumor-associated macrophages in developing mycosis fungoides. <i>Experimental Dermatology</i> , 2016 , 25, 107-12	4	54
22	Tumor-associated macrophages in skin: How to treat their heterogeneity and plasticity. <i>Journal of Dermatological Science</i> , 2016 , 83, 167-73	4.3	38
21	Optimization of the IL-8 Luc assay as an in vitro test for skin sensitization. <i>Toxicology in Vitro</i> , 2015 , 29, 1816-30	3.6	37
20	Variation in the Effect of Particulate Matter on Pulmonary Function in Schoolchildren in Western Japan and Its Relation with Interleukin-8. <i>International Journal of Environmental Research and Public Health</i> , 2015 , 12, 14229-43	4.6	5
19	Decreased pulmonary function in school children in Western Japan after exposures to Asian desert dusts and its association with interleukin-8. <i>BioMed Research International</i> , 2015 , 2015, 583293	3	17
18	Suppression of Propionibacterium acnes-Induced Dermatitis by a Traditional Japanese Medicine, Jumihaidokuto, Modifying Macrophage Functions. <i>Evidence-based Complementary and Alternative Medicine</i> , 2015 , 2015, 439258	2.3	8
17	Alarmin function of cathelicidin antimicrobial peptide LL37 through IL-36 β induction in human epidermal keratinocytes. <i>Journal of Immunology</i> , 2014 , 193, 5140-8	5.3	99
16	Skin sensitization risk assessment model using artificial neural network analysis of data from multiple in vitro assays. <i>Toxicology in Vitro</i> , 2014 , 28, 626-39	3.6	45
15	Nonmetal haptens induce ATP release from keratinocytes through opening of pannexin hemichannels by reactive oxygen species. <i>Journal of Investigative Dermatology</i> , 2014 , 134, 1951-1960	4.3	27
14	Evaluation of the Multi-ImmunoTox Assay composed of 3 human cytokine reporter cells by examining immunological effects of drugs. <i>Toxicology in Vitro</i> , 2014 , 28, 759-68	3.6	13
13	In vivo imaging demonstrates ATP release from murine keratinocytes and its involvement in cutaneous inflammation after tape stripping. <i>Journal of Investigative Dermatology</i> , 2013 , 133, 2407-2415	4.3	19
12	Nickel differentially regulates NFAT and NF- κ B activation in T cell signaling. <i>Toxicology and Applied Pharmacology</i> , 2011 , 254, 245-55	4.6	18
11	Dual oxidase 1 induced by Th2 cytokines promotes STAT6 phosphorylation via oxidative inactivation of protein tyrosine phosphatase 1B in human epidermal keratinocytes. <i>Journal of Immunology</i> , 2011 , 186, 4762-70	5.3	41
10	An in vitro test to screen skin sensitizers using a stable THP-1-derived IL-8 reporter cell line, THP-G8. <i>Toxicological Sciences</i> , 2011 , 124, 359-69	4.4	57
9	Bach1-dependent and -independent regulation of heme oxygenase-1 in keratinocytes. <i>Journal of Biological Chemistry</i> , 2010 , 285, 23581-9	5.4	19

8	Gene expression profiling defines the role of ATP-exposed keratinocytes in skin inflammation. <i>Journal of Dermatological Science</i> , 2010 , 58, 143-51	4-3	13
7	Functional expression of heme oxygenase-1 in human differentiated epidermis and its regulation by cytokines. <i>Journal of Investigative Dermatology</i> , 2009 , 129, 2594-603	4-3	26
6	Miniaturization of Micro Implantable Devices With Thermosensitive Ferrite for Soft-Heating Hyperthermia. <i>IEEE Transactions on Magnetics</i> , 2007 , 43, 2454-2456	2	2
5	Redox imbalance induced by contact sensitizers triggers the maturation of dendritic cells. <i>Journal of Investigative Dermatology</i> , 2005 , 124, 579-86	4-3	89
4	p38 Mitogen-activated protein kinase and extracellular signal-regulated kinases play distinct roles in the activation of dendritic cells by two representative haptens, NiCl ₂ and 2,4-dinitrochlorobenzene. <i>Journal of Investigative Dermatology</i> , 2003 , 120, 390-9	4-3	127
3	Thermolysin improves mutation analysis in skin epidermis from ultraviolet light-irradiated Muta Mouse. <i>Environmental and Molecular Mutagenesis</i> , 2001 , 38, 55-8	3-2	13
2	Dendritic cells differently respond to haptens and irritants by their production of cytokines and expression of co-stimulatory molecules. <i>European Journal of Immunology</i> , 1997 , 27, 3031-8	6-1	216
1	Cutaneous metastasis of invasive lobular carcinoma of the breast. <i>Journal of Dermatology</i> , 1987 , 14, 270-36		3