
## Silvia de la Barrera

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7489155/publications.pdf Version: 2024-02-01



| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Patients with Multidrug-Resistant Tuberculosis Display Impaired Th1 Responses and Enhanced<br>Regulatory T-Cell Levels in Response to an Outbreak of Multidrug-Resistant <i>Mycobacterium<br/>tuberculosis</i> M and Ra Strains. Infection and Immunity, 2009, 77, 5025-5034. | 2.2 | 67        |
| 2  | <i>Mycobacterium tuberculosis</i> impairs dendritic cell response by altering CD1b, DCâ€&IGN and MR profile. Immunology and Cell Biology, 2010, 88, 716-726.  | 2.3 | 45        |
| 3  | NK cell activity in tuberculosis is associated with impaired CD11a and ICAM-1 expression: a regulatory role of monocytes in NK activation. Immunology, 2005, 116, 051025020346008.  | 4.4 | 42        |
| 4  | Role of neutrophils in CVB3 infection and viral myocarditis. Journal of Molecular and Cellular<br>Cardiology, 2018, 125, 149-161.   | 1.9 | 42        |
| 5  | Mycobacterium tuberculosis Multidrug Resistant Strain M Induces an Altered Activation of Cytotoxic<br>CD8+ T Cells. PLoS ONE, 2014, 9, e97837.  | 2.5 | 12        |
| 6  | <i>Mycobacterium tuberculosis</i> Multidrug-Resistant Strain M Induces Low IL-8 and Inhibits<br>TNF- <i>α</i> Secretion by Bronchial Epithelial Cells Altering Neutrophil Effector Functions. Mediators<br>of Inflammation, 2017, 2017, 1-13.                                 | 3.0 | 11        |
| 7  | Single nucleotide polymorphisms may explain the contrasting phenotypes of two variants of a multidrug-resistant Mycobacterium tuberculosis strain. Tuberculosis, 2017, 103, 28-36.  | 1.9 | 10        |
| 8  | Two genetically-related multidrug-resistant Mycobacterium tuberculosis strains induce divergent<br>outcomes of infection in two human macrophage models. Infection, Genetics and Evolution, 2013, 16,<br>151-156.   | 2.3 | 9         |
| 9  | C5aR contributes to the weak Th1 profile induced by an outbreak strain of Mycobacterium tuberculosis. Tuberculosis, 2017, 103, 16-23.   | 1.9 | 7         |
| 10 | Differential induction of macrophage cell death by antigens of a clustered and a non-clustered<br>multidrug-resistantMycobacterium tuberculosisstrain from Haarlem family. FEMS Immunology and<br>Medical Microbiology, 2012, 66, 363-371.                                    | 2.7 | 6         |
| 11 | Effect of lymphokines on natural killer cytotoxicity in patients with high risk of developing the acquired immune deficiency syndrome. Immunology Letters, 1986, 13, 307-311.   | 2.5 | 5         |
| 12 | A Phenotypic Characterization of Two Isolates of a Multidrug-Resistant Outbreak Strain of<br><i>Mycobacterium tuberculosis</i> with Opposite Epidemiological Fitness. BioMed Research<br>International, 2020, 2020, 1-9.  | 1.9 | 2         |