
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7489855/publications.pdf Version: 2024-02-01



Μλελτεμομ Εμλ

#	Article	IF	CITATIONS
1	A New Model for Specific Visualization of Skin Graft Neoangiogenesis Using Flt1-tdsRed BAC Transgenic Mice. Plastic and Reconstructive Surgery, 2021, 148, 89-99.	1.4	2
2	Chromosomal-scale de novo genome assemblies of Cynomolgus Macaque and Common Marmoset. Scientific Data, 2021, 8, 159.	5.3	9
3	The X chromosome dosage compensation program during the development of cynomolgus monkeys. Science, 2021, 374, eabd8887.	12.6	33
4	Intersection of regulatory pathways controlling hemostasis and hemochorial placentation. Proceedings of the National Academy of Sciences of the United States of America, 2021, 118, .	7.1	19
5	Hepatocyte ELOVL Fatty Acid Elongase 6 Determines Ceramide Acylâ€Chain Length and Hepatic Insulin Sensitivity in Mice. Hepatology, 2020, 71, 1609-1625.	7.3	44
6	Induction of the germ cell fate from pluripotent stem cells in cynomolgus monkeysâ€. Biology of Reproduction, 2020, 102, 620-638.	2.7	40
7	Neuron-derived VEGF contributes to cortical and hippocampal development independently of VEGFR1/2-mediated neurotrophism. Developmental Biology, 2020, 459, 65-71.	2.0	10
8	Aging of the Vascular System and Neural Diseases. Frontiers in Aging Neuroscience, 2020, 12, 557384.	3.4	21
9	Simultaneous fluorescence imaging of distinct nerve and blood vessel patterns in dual Thy1-YFP and Flt1-DsRed transgenic mice. Angiogenesis, 2020, 23, 459-477.	7.2	7
10	Macrophages fine-tune pupil shape during development. Developmental Biology, 2020, 464, 137-144.	2.0	1
11	Establishment of macaque trophoblast stem cell lines derived from cynomolgus monkey blastocysts. Scientific Reports, 2020, 10, 6827.	3.3	10
12	Regulation of ERK signalling pathway in the developing mouse blastocyst. Development (Cambridge), 2019, 146, .	2.5	23
13	Interaction of the nervous system and vascular system is required for the proper assembly of the neocortex. Neurochemistry International, 2019, 129, 104481.	3.8	12
14	Highly efficient induction of primate iPS cells by combining RNA transfection and chemical compounds. Genes To Cells, 2019, 24, 473-484.	1.2	19
15	Generation of an OCT3/4 reporter cynomolgus monkey ES cell line using CRISPR/Cas9. Stem Cell Research, 2019, 37, 101439.	0.7	4
16	Comprehensive evaluation of ubiquitous promoters suitable for the generation of transgenic cynomolgus monkeysâ€. Biology of Reproduction, 2019, 100, 1440-1452.	2.7	12
17	Quantification of Angiogenesis and Lymphangiogenesis in the Dual ex vivo Aortic and Thoracic Duct Assay. Protein and Peptide Letters, 2019, 27, 30-40.	0.9	4
18	Monkeys mutant for PKD1 recapitulate human autosomal dominant polycystic kidney disease. Nature Communications, 2019, 10, 5517.	12.8	33

#	Article	IF	CITATIONS
19	The dynamics of revascularization after white matter infarction monitored in Flt1-tdsRed and Flk1-GFP mice. Neuroscience Letters, 2019, 692, 70-76.	2.1	5
20	Generating Vegfr3 reporter transgenic mouse expressing membrane-tagged Venus for visualization of VEGFR3 expression in vascular and lymphatic endothelial cells. PLoS ONE, 2019, 14, e0210060.	2.5	11
21	The transcription factor Klf5 is essential for intrahepatic biliary epithelial tissue remodeling after cholestatic liver injury. Journal of Biological Chemistry, 2018, 293, 6214-6229.	3.4	14
22	Distinct expression patterns of Flk1 and Flt1 in the coronary vascular system during development and after myocardial infarction. Biochemical and Biophysical Research Communications, 2018, 495, 884-891.	2.1	18
23	Klf5 suppresses ERK signaling in mouse pluripotent stem cells. PLoS ONE, 2018, 13, e0207321.	2.5	17
24	Fluorescent reporter transgenic mice for in vivo live imaging of angiogenesis and lymphangiogenesis. Angiogenesis, 2018, 21, 677-698.	7.2	15
25	Prox1-GFP/Flt1-DsRed transgenic mice: an animal model for simultaneous live imaging of angiogenesis and lymphangiogenesis. Angiogenesis, 2017, 20, 581-598.	7.2	28
26	<i>Klf5</i> maintains the balance of primitive endoderm to epiblast specification during mouse embryonic development by suppression of <i>Fgf4</i> . Development (Cambridge), 2017, 144, 3706-3718.	2.5	24
27	Comprehensive Identification of Krüppel-Like Factor Family Members Contributing to the Self-Renewal of Mouse Embryonic Stem Cells and Cellular Reprogramming. PLoS ONE, 2016, 11, e0150715.	2.5	29
28	Visualization of the Epiblast and Visceral Endodermal Cells Using Fgf5-P2A-Venus BAC Transgenic Mice and Epiblast Stem Cells. PLoS ONE, 2016, 11, e0159246.	2.5	14
29	Developmental regression of hyaloid vasculature is triggered by neurons. Journal of Experimental Medicine, 2016, 213, 1175-1183.	8.5	43
30	Forced Expression of Nanog or Esrrb Preserves the ESC Status in the Absence of Nucleostemin Expression. Stem Cells, 2015, 33, 1089-1101.	3.2	6
31	Functional Compensation Between Myc and PI3K Signaling Supports Self-Renewal of Embryonic Stem Cells, 2015, 33, 713-725.	3.2	13
32	Platelet demand modulates the type of intravascular protrusion of megakaryocytes in bone marrow. Thrombosis and Haemostasis, 2014, 112, 743-756.	3.4	35
33	Roles of VEGF-A signalling in development, regeneration, and tumours. Journal of Biochemistry, 2014, 156, 1-10.	1.7	159
34	Neurons Limit Angiogenesis by Titrating VEGF in Retina. Cell, 2014, 159, 584-596.	28.9	232
35	Generation and Characterization of Ins1-cre-driver C57BL/6N for Exclusive Pancreatic Beta Cell-specific Cre-loxP Recombination. Experimental Animals, 2014, 63, 183-191.	1.1	24
36	Klf9 is necessary and sufficient for Purkinje cell survival in organotypic culture. Molecular and Cellular Neurosciences, 2013, 54, 9-21.	2.2	22

#	Article	IF	CITATIONS
37	Novel ROSA26 Cre-reporter Knock-in C57BL/6N Mice Exhibiting Green Emission before and Red Emission after Cre-mediated Recombination. Experimental Animals, 2013, 62, 295-304.	1.1	53
38	In Vivo Function and Evolution of the Eutherian-Specific Pluripotency Marker UTF1. PLoS ONE, 2013, 8, e68119.	2.5	17
39	Bioluminescence Imaging of \hat{l}^2 Cells and Intrahepatic Insulin Gene Activity under Normal and Pathological Conditions. PLoS ONE, 2013, 8, e60411.	2.5	13
40	Thyroid hormone triggers the developmental loss of axonal regenerative capacity via thyroid hormone receptor α1 and krüppel-like factor 9 in Purkinje cells. Proceedings of the National Academy of Sciences of the United States of America, 2012, 109, 14206-14211.	7.1	56
41	Delayed cutaneous wound healing in Fam129b/Minerva-deficient mice. Journal of Biochemistry, 2012, 152, 549-555.	1.7	10
42	Flt1 and Flk1 mediate regulation of intraocular pressure and their double heterozygosity causes the buphthalmia in mice. Biochemical and Biophysical Research Communications, 2012, 420, 422-427.	2.1	6
43	Indefinite Self-Renewal of ESCs through Myc/Max Transcriptional Complex-Independent Mechanisms. Cell Stem Cell, 2011, 9, 37-49.	11.1	64
44	The E3 Ubiquitin Ligase Activity of Trip12 Is Essential for Mouse Embryogenesis. PLoS ONE, 2011, 6, e25871.	2.5	22
45	RNA content in the nucleolus alters p53 acetylation via MYBBP1A. EMBO Journal, 2011, 30, 1054-1066.	7.8	62
46	Isolation and function of mouse tissue resident vascular precursors marked by myelin protein zero. Journal of Experimental Medicine, 2011, 208, 949-960.	8.5	34
47	Chondroitin Sulfate N-Acetylgalactosaminyltransferase 1 Is Necessary for Normal Endochondral Ossification and Aggrecan Metabolism. Journal of Biological Chemistry, 2011, 286, 5803-5812.	3.4	60
48	Inhibition of Ubiquitin Ligase F-box and WD Repeat Domain-containing 7α (Fbw7α) Causes Hepatosteatosis through Krüppel-like Factor 5 (KLF5)/Peroxisome Proliferator-activated Receptor γ2 (PPARγ2) Pathway but Not SREBP-1c Protein in Mice*. Journal of Biological Chemistry, 2011, 286, 40835-40846.	3.4	24
49	Flk1-GFP BAC Tg Mice: An Animal Model for the Study of Blood Vessel Development. Experimental Animals, 2010, 59, 615-622.	1.1	42
50	Subventricular Zone-Derived Neural Progenitor Cells Migrate Along a Blood Vessel Scaffold Toward The Post-stroke Striatum. Stem Cells, 2010, 28, 545-554.	3.2	261
51	Flt-1 haploinsufficiency ameliorates muscular dystrophy phenotype by developmentally increased vasculature in mdx mice. Human Molecular Genetics, 2010, 19, 4145-4159.	2.9	49
52	Antitumor effects of 2â€oxoglutarate through inhibition of angiogenesis in a murine tumor model. Cancer Science, 2009, 100, 1639-1647.	3.9	41
53	Poor vessel formation in embryos from knock-in mice expressing ALK5 with L45 loop mutation defective in Smad activation. Laboratory Investigation, 2009, 89, 800-810.	3.7	19
54	Genetic evidence of PEBP2β-independent activation of Runx1 in the murine embryo. International Journal of Hematology, 2008, 88, 134-138.	1.6	5

#	Article	IF	CITATIONS
55	Krüppel-like factor 5 Is Essential for Blastocyst Development and the Normal Self-Renewal of Mouse ESCs. Cell Stem Cell, 2008, 3, 555-567.	11.1	177
56	Runx1 is involved in primitive erythropoiesis in the mouse. Blood, 2008, 111, 4075-4080.	1.4	59
57	Compensatory signalling induced in the yolk sac vasculature by deletion of TGFÎ ² receptors in mice. Journal of Cell Science, 2007, 120, 4269-4277.	2.0	104
58	Characterization of GATA-1+ hemangioblastic cells in the mouse embryo. EMBO Journal, 2007, 26, 184-196.	7.8	48
59	Deletion of the selection cassette, but not cis-acting elements, in targeted Flk1-lacZ allele reveals Flk1 expression in multipotent mesodermal progenitors. Blood, 2006, 107, 111-117.	1.4	259
60	Vascular Endothelial Growth Factor A Signaling in the Podocyte-Endothelial Compartment Is Required for Mesangial Cell Migration and Survival. Journal of the American Society of Nephrology: JASN, 2006, 17, 724-735.	6.1	217
61	Vascular Endothelial Growth Factor Directly Inhibits Primitive Neural Stem Cell Survival But Promotes Definitive Neural Stem Cell Survival. Journal of Neuroscience, 2006, 26, 6803-6812.	3.6	95
62	Rapid Isolation of Glomeruli Coupled with Gene Expression Profiling Identifies Downstream Targets in Pod1 Knockout Mice. Journal of the American Society of Nephrology: JASN, 2005, 16, 3247-3255.	6.1	46
63	Activated Fps/Fes partially rescues the in vivo developmental potential of Flk1-deficient vascular progenitor cells. Blood, 2004, 103, 912-920.	1.4	15
64	HLF/HIF-2α is a key factor in retinopathy of prematurity in association with erythropoietin. EMBO Journal, 2003, 22, 1134-1146.	7.8	220
65	Cell Fate Decisions in Early Blood Vessel Formation. Trends in Cardiovascular Medicine, 2003, 13, 254-259.	4.9	69
66	Heterodimers of bHLH-PAS Protein Fragments Derived from AhR, AhRR, and Arnt Prepared by Co-Expression in Escherichia coli: Characterization of Their DNA Binding Activity and Preparation of a DNA Complex. Journal of Biochemistry, 2003, 134, 83-90.	1.7	35
67	Combinatorial effects of Flk1 and Tal1 on vascular and hematopoietic development in the mouse. Genes and Development, 2003, 17, 380-393.	5.9	232
68	Defective development of secretory neurones in the hypothalamus of Arnt2â€knockout mice. Genes To Cells, 2001, 6, 361-374.	1.2	99
69	Mild Impairment of Learning and Memory in Mice Overexpressing the mSim2 Gene Located on Chromosome 16: An Animal Model of Down's Syndrome. Human Molecular Genetics, 1999, 8, 1409-1415.	2.9	79
70	Transcriptionally Active Heterodimer Formation of an Arnt-like PAS Protein, Arnt3, with HIF-1a, HLF, and Clock. Biochemical and Biophysical Research Communications, 1998, 248, 789-794.	2.1	128
71	Inhibition of hypoxia-inducible factor 1 activity by nitric oxide donors in hypoxia. Proceedings of the National Academy of Sciences of the United States of America, 1998, 95, 7368-7373.	7.1	221
72	Regulation of Dioxin Receptor Function by Omeprazole. Journal of Biological Chemistry, 1997, 272, 12705-12713.	3.4	72

#	Article	IF	CITATIONS
73	cDNA Cloning of a Murine Homologue ofDrosophilaSingle-Minded, Its mRNA Expression in Mouse Development, and Chromosome Localization. Biochemical and Biophysical Research Communications, 1996, 218, 588-594.	2.1	50
74	cDNA cloning and structure of mouse putative Ah receptor. Biochemical and Biophysical Research Communications, 1992, 184, 246-253.	2.1	396