

# Carlos Martin-Cantera

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7495119/publications.pdf>

Version: 2024-02-01

50  
papers

1,120  
citations

430442

18  
h-index

454577

30  
g-index

61  
all docs

61  
docs citations

61  
times ranked

2138  
citing authors

#	ARTICLE	IF	CITATIONS
1	Effectiveness of a healthcare-based mobile intervention on sedentary patterns, physical activity, mental well-being and clinical and productivity outcomes in office employees with type 2 diabetes: study protocol for a randomized controlled trial. <i>BMC Public Health</i> , 2022, 22, .	1.2	6
2	Patterns of Physical Activity Progression in Patients With COPD. <i>Archivos De Bronconeumologia</i> , 2021, 57, 214-223.	0.4	9
3	Patterns of Physical Activity Progression in Patients With COPD. <i>Archivos De Bronconeumologia</i> , 2021, 57, 214-223.	0.4	1
4	6-Year Risk of Developing Lung Cancer in Spain: Analysis by Autonomous Communities. <i>Archivos De Bronconeumologia</i> , 2021, 57, 521-527.	0.4	2
5	Good practice regarding smoking cessation management in Spain: Challenges and opportunities for primary care physicians and nurses. <i>Tobacco Prevention and Cessation</i> , 2020, 6, 1-9.	0.2	7
6	The Relationship of the Atlantic Diet with Cardiovascular Risk Factors and Markers of Arterial Stiffness in Adults without Cardiovascular Disease. <i>Nutrients</i> , 2019, 11, 742.	1.7	20
7	Combined use of a healthy lifestyle smartphone application and usual primary care counseling to improve arterial stiffness, blood pressure and wave reflections: a Randomized Controlled Trial (EVIDENT II Study). <i>Hypertension Research</i> , 2019, 42, 852-862.	1.5	14
8	Effectiveness of the Epley manoeuvre in posterior canal benign paroxysmal positional vertigo: a randomised clinical trial in primary care. <i>British Journal of General Practice</i> , 2019, 69, e52-e60.	0.7	17
9	Attitudes towards anti-smoking legislation and prevalence of tobacco consumption in Spanish primary healthcare personnel. <i>Tobacco Prevention and Cessation</i> , 2019, 5, 9.	0.2	6
10	Complex multiple risk intervention to promote healthy behaviours in people between 45 to 75 years attended in primary health care (EIRA study): study protocol for a hybrid trial. <i>BMC Public Health</i> , 2018, 18, 874.	1.2	13
11	Long-term efficacy and effectiveness of a behavioural and community-based exercise intervention (Urban Training) to increase physical activity in patients with COPD: a randomised controlled trial. <i>European Respiratory Journal</i> , 2018, 52, 1800063.	3.1	79
12	Identifying high-risk individuals for lung cancer screening: Going beyond NLST criteria. <i>PLoS ONE</i> , 2018, 13, e0195441.	1.1	7
13	Patterns of sedentary behavior in overweight and moderately obese users of the Catalan primary-health care system. <i>PLoS ONE</i> , 2018, 13, e0190750.	1.1	15
14	The EVIDENT diet quality index is associated with cardiovascular risk and arterial stiffness in adults. <i>BMC Public Health</i> , 2017, 17, 305.	1.2	14
15	Physical Activity and Adiposity Among Older Adults of the EVIDENT Study. <i>Journal of Aging and Physical Activity</i> , 2017, 25, 254-260.	0.5	6
16	Smokefree legislation effects on respiratory and sensory disorders: A systematic review and meta-analysis. <i>PLoS ONE</i> , 2017, 12, e0181035.	1.1	20
17	Validation of Walking Trails for the Urban Training™ of Chronic Obstructive Pulmonary Disease Patients. <i>PLoS ONE</i> , 2016, 11, e0146705.	1.1	20
18	Dietary glycemic index and retinal microvasculature in adults: a cross-sectional study. <i>Nutrition Journal</i> , 2016, 15, 88.	1.5	2

#	ARTICLE	IF	CITATIONS
19	Short-Term Effectiveness of a Mobile Phone App for Increasing Physical Activity and Adherence to the Mediterranean Diet in Primary Care: A Randomized Controlled Trial (EVIDENT II Study). <i>Journal of Medical Internet Research</i> , 2016, 18, e331.	2.1	72
20	Clustering of lifestyle characteristics and their association with cardio-metabolic health: the Lifestyles and Endothelial Dysfunction (EVIDENT) study. <i>British Journal of Nutrition</i> , 2015, 114, 943-951.	1.2	17
21	Sedentary behaviour patterns and arterial stiffness in a Spanish adult population – The EVIDENT trial. <i>Atherosclerosis</i> , 2015, 243, 516-522.	0.4	15
22	Information and communication technologies for approaching smokers: a descriptive study in primary healthcare. <i>BMC Public Health</i> , 2015, 15, 2.	1.2	22
23	Continued smoking abstinence in diabetic patients in primary care: A cluster randomized controlled multicenter study. <i>Diabetes Research and Clinical Practice</i> , 2015, 107, 94-103.	1.1	15
24	Sedentary behaviour patterns and carotid intima-media thickness in Spanish healthy adult population. <i>Atherosclerosis</i> , 2015, 239, 571-576.	0.4	14
25	Effectiveness of multicomponent interventions in primary healthcare settings to promote continuous smoking cessation in adults: a systematic review. <i>BMJ Open</i> , 2015, 5, e008807.	0.8	35
26	Effectiveness of a brief primary care intervention to reduce passive smoking in babies: a cluster randomised clinical trial. <i>Journal of Epidemiology and Community Health</i> , 2015, 69, 249-260.	2.0	6
27	Glycemic index, glycemic load, and pulse wave reflection in adults. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2015, 25, 68-74.	1.1	12
28	Prolonged Sitting Time: Barriers, Facilitators and Views on Change among Primary Healthcare Patients Who Are Overweight or Moderately Obese. <i>PLoS ONE</i> , 2015, 10, e0125739.	1.1	22
29	Screening Physical Activity in Family Practice: Validity of the Spanish Version of a Brief Physical Activity Questionnaire. <i>PLoS ONE</i> , 2015, 10, e0136870.	1.1	35
30	Intervention for Smokers through New Communication Technologies: What Perceptions Do Patients and Healthcare Professionals Have? A Qualitative Study. <i>PLoS ONE</i> , 2015, 10, e0137415.	1.1	11
31	Effectiveness of a smartphone application for improving healthy lifestyles, a randomized clinical trial (EVIDENT II): study protocol. <i>BMC Public Health</i> , 2014, 14, 254.	1.2	53
32	Electrocardiographic Left Ventricular Hypertrophy Criteria and Ambulatory Blood Pressure Monitoring Parameters in Adults. <i>American Journal of Hypertension</i> , 2014, 27, 355-362.	1.0	6
33	Effectiveness of a primary care-based intervention to reduce sitting time in overweight and obese patients (SEDESTACTIV): a randomized controlled trial; rationale and study design. <i>BMC Public Health</i> , 2014, 14, 228.	1.2	13
34	Effectiveness of the Epley’s maneuver performed in primary care to treat posterior canal benign paroxysmal positional vertigo: study protocol for a randomized controlled trial. <i>Trials</i> , 2014, 15, 179.	0.7	11
35	Relationship between objectively measured physical activity and cardiovascular aging in the general population – The EVIDENT trial. <i>Atherosclerosis</i> , 2014, 233, 434-440.	0.4	36
36	Relationship between Physical Activity and Plasma Fibrinogen Concentrations in Adults without Chronic Diseases. <i>PLoS ONE</i> , 2014, 9, e87954.	1.1	19

#	ARTICLE	IF	CITATIONS
37	Effectiveness of an intensive E-mail based intervention in smoking cessation (TABATIC study): study protocol for a randomized controlled trial. BMC Public Health, 2013, 13, 364.	1.2	8
38	Association between smoking status and the parameters of vascular structure and function in adults: results from the EVIDENT study. BMC Cardiovascular Disorders, 2013, 13, 109.	0.7	8
39	The Effect of a Physical Activity Program on the Total Number of Primary Care Visits in Inactive Patients: A 15-Month Randomized Controlled Trial. PLoS ONE, 2013, 8, e66392.	1.1	5
40	Comparison of two measuring instruments, B-pro and SphygmoCor system as reference, to evaluate central systolic blood pressure and radial augmentation index. Hypertension Research, 2012, 35, 617-623.	1.5	42
41	Spinal Cord Injury and Traumatic Brain Injury: A Cost-of-Illness Study. Neuroepidemiology, 2012, 39, 103-108.	1.1	40
42	Incidence trends of traumatic spinal cord injury and traumatic brain injury in Spain, 2000-2009. Accident Analysis and Prevention, 2012, 46, 37-44.	3.0	68
43	Relationship between intima-media thickness of the common carotid artery and arterial stiffness in subjects with and without type 2 diabetes: a case-series report. Cardiovascular Diabetology, 2011, 10, 3.	2.7	39
44	Risk levels for suffering a traffic injury in primary health care. The LESIONAT* project. BMC Public Health, 2010, 10, 136.	1.2	2
45	Physical exercise, fitness and dietary pattern and their relationship with circadian blood pressure pattern, augmentation index and endothelial dysfunction biological markers: EVIDENT study protocol. BMC Public Health, 2010, 10, 233.	1.2	50
46	Passive smoking in babies: The BIBE study (Brief Intervention in babies. Effectiveness). BMC Public Health, 2010, 10, 772.	1.2	14
47	Therapeutic implications of selecting the SCORE (European) versus the D'AGOSTINO (American) risk charts for cardiovascular risk assessment in hypertensive patients. BMC Cardiovascular Disorders, 2009, 9, 17.	0.7	7
48	¿Aumentan los anticolinérgicos inhalados el riesgo de eventos cardiovasculares en pacientes con enfermedad pulmonar obstructiva crónica?. FMC Formacion Medica Continuada En Atencion Primaria, 2009, 16, 315.	0.0	0
49	El tratamiento ambulatorio de la neumonía de bajo riesgo es seguro y eficaz con levofloxacino. FMC Formacion Medica Continuada En Atencion Primaria, 2005, 12, 415.	0.0	0
50	Efficacy and Safety of Metamizol vs. Acetylsalicylic Acid in Patients With Moderate Episodic Tension-Type Headache: A Randomized, Double-Blind, Placebo- and Active-Controlled, Multicentre Study. Cephalalgia, 2001, 21, 604-610.	1.8	65