

# Oliver Cooper

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/750054/oliver-cooper-publications-by-year.pdf>

**Version:** 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

24  
papers

3,740  
citations

18  
h-index

37  
g-index

37  
ext. papers

4,071  
ext. citations

10.2  
avg, IF

4.48  
L-index

#	Paper	IF	Citations
24	Successful function of autologous iPSC-derived dopamine neurons following transplantation in a non-human primate model of Parkinson's disease. <i>Cell Stem Cell</i> , <b>2015</b> , 16, 269-74	18	214
23	LRRK2 mutations cause mitochondrial DNA damage in iPSC-derived neural cells from Parkinson's disease patients: reversal by gene correction. <i>Neurobiology of Disease</i> , <b>2014</b> , 62, 381-6	7.5	194
22	Long-term health of dopaminergic neuron transplants in Parkinson's disease patients. <i>Cell Reports</i> , <b>2014</b> , 7, 1755-61	10.6	112
21	Improved cell therapy protocols for Parkinson's disease based on differentiation efficiency and safety of hESC-, hiPSC-, and non-human primate iPSC-derived dopaminergic neurons. <i>Stem Cells</i> , <b>2013</b> , 31, 1548-62	5.8	168
20	Using stem cells and iPS cells to discover new treatments for Parkinson's disease. <i>Parkinsonism and Related Disorders</i> , <b>2012</b> , 18 Suppl 1, S14-6	3.6	13
19	Transcript expression levels of full-length alpha-synuclein and its three alternatively spliced variants in Parkinson's disease brain regions and in a transgenic mouse model of alpha-synuclein overexpression. <i>Molecular and Cellular Neurosciences</i> , <b>2012</b> , 49, 230-9	4.8	35
18	Characterization and criteria of embryonic stem and induced pluripotent stem cells for a dopamine replacement therapy. <i>Progress in Brain Research</i> , <b>2012</b> , 200, 265-76	2.9	12
17	Pharmacological rescue of mitochondrial deficits in iPSC-derived neural cells from patients with familial Parkinson's disease. <i>Science Translational Medicine</i> , <b>2012</b> , 4, 141ra90	17.5	381
16	Oct4-induced reprogramming is required for adult brain neural stem cell differentiation into midbrain dopaminergic neurons. <i>PLoS ONE</i> , <b>2011</b> , 6, e19926	3.7	33
15	Differentiated Parkinson patient-derived induced pluripotent stem cells grow in the adult rodent brain and reduce motor asymmetry in Parkinsonian rats. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2010</b> , 107, 15921-6	11.5	375
14	Differentiation of human ES and Parkinson's disease iPSCs into ventral midbrain dopaminergic neurons requires a high activity form of SHH, FGF8a and specific regionalization by retinoic acid. <i>Molecular and Cellular Neurosciences</i> , <b>2010</b> , 45, 258-66	4.8	175
13	No evidence for disease-like processes in fetal transplants. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2009</b> , 106, E104; author reply E105	11.5	2
12	Lack of functional relevance of isolated cell damage in transplants of Parkinson's disease patients. <i>Journal of Neurology</i> , <b>2009</b> , 256 Suppl 3, 310-6	5.5	42
11	Parkinson's disease patient-derived induced pluripotent stem cells free of viral reprogramming factors. <i>Cell</i> , <b>2009</b> , 136, 964-77	56.2	1262
10	Parkinson's Disease Patient-Derived Induced Pluripotent Stem Cells Free of Viral Reprogramming Factors. <i>Cell</i> , <b>2009</b> , 137, 1356	56.2	6
9	Klhl31 is associated with skeletal myogenesis and its expression is regulated by myogenic signals and Myf-5. <i>Mechanisms of Development</i> , <b>2009</b> , 126, 852-62	1.7	15
8	Recent advances in cell-based therapy for Parkinson disease. <i>Neurosurgical Focus</i> , <b>2008</b> , 24, E6	4.2	30

7	The migration of paraxial and lateral plate mesoderm cells emerging from the late primitive streak is controlled by different Wnt signals. <i>BMC Developmental Biology</i> , <b>2008</b> , 8, 63	3.1	51
6	Expression of avian prickle genes during early development and organogenesis. <i>Developmental Dynamics</i> , <b>2008</b> , 237, 1442-8	2.9	13
5	Fate mapping and lineage analyses demonstrate the production of a large number of striatal neuroblasts after transforming growth factor alpha and noggin striatal infusions into the dopamine-depleted striatum. <i>Stem Cells</i> , <b>2008</b> , 26, 2349-60	5.8	48
4	Neuroblast protuberances in the subventricular zone of the regenerative MRL/MpJ mouse. <i>Journal of Comparative Neurology</i> , <b>2006</b> , 498, 747-61	3.4	28
3	Cell type analysis of functional fetal dopamine cell suspension transplants in the striatum and substantia nigra of patients with Parkinson disease. <i>Brain</i> , <b>2005</b> , 128, 1498-510	11.2	352
2	Context-dependent neuronal differentiation and germ layer induction of Smad4 <sup>-/-</sup> and Cripto <sup>-/-</sup> embryonic stem cells. <i>Molecular and Cellular Neurosciences</i> , <b>2005</b> , 28, 417-29	4.8	32
1	Intrastriatal transforming growth factor alpha delivery to a model of Parkinson disease induces proliferation and migration of endogenous adult neural progenitor cells without differentiation into dopaminergic neurons. <i>Journal of Neuroscience</i> , <b>2004</b> , 24, 8924-31	6.6	147