## Erling A Hoivik

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7506722/publications.pdf Version: 2024-02-01



| #  | Article  | IF   | CITATIONS |
|----|--|------|-----------|
| 1  | Landscape of genomic alterations in cervical carcinomas. Nature, 2014, 506, 371-375.   | 27.8 | 708       |
| 2  | The genomic landscape and evolution of endometrial carcinoma progression and abdominopelvic metastasis. Nature Genetics, 2016, 48, 848-855.  | 21.4 | 174       |
| 3  | Lack of Estrogen Receptor-α Is Associated with Epithelial–Mesenchymal Transition and PI3K Alterations<br>in Endometrial Carcinoma. Clinical Cancer Research, 2013, 19, 1094-1105.                        | 7.0  | 120       |
| 4  | A Novel Wnt Regulatory Axis in Endometrioid Endometrial Cancer. Cancer Research, 2014, 74, 5103-5117.  | 0.9  | 114       |
| 5  | Loss of progesterone receptor links to high proliferation and increases from primary to metastatic endometrial cancer lesions. European Journal of Cancer, 2014, 50, 3003-3010.                          | 2.8  | 73        |
| 6  | Molecular profiling of endometrial carcinoma precursor, primary and metastatic lesions suggests different targets for treatment in obese compared to non-obese patients. Oncotarget, 2015, 6, 1327-1339. | 1.8  | 50        |
| 7  | Hypomethylation of the CTCFL/BORIS promoter and aberrant expression during endometrial cancer progression suggests a role as an Epi-driver gene. Oncotarget, 2014, 5, 1052-1061.                         | 1.8  | 35        |
| 8  | A Common Variant at the 14q32 Endometrial Cancer Risk Locus Activates AKT1 through YY1 Binding.<br>American Journal of Human Genetics, 2016, 98, 1159-1169.  | 6.2  | 32        |
| 9  | High degree of heterogeneity of PD-L1 and PD-1 from primary to metastatic endometrial cancer.<br>Gynecologic Oncology, 2020, 157, 260-267.   | 1.4  | 32        |
| 10 | Class I Phosphoinositide 3-Kinase PIK3CA/p110α and PIK3CB/p110β Isoforms in Endometrial Cancer.<br>International Journal of Molecular Sciences, 2018, 19, 3931.  | 4.1  | 26        |
| 11 | Patient-derived organoids reflect the genetic profile of endometrial tumors and predict patient prognosis. Communications Medicine, 2021, 1, .   | 4.2  | 20        |
| 12 | PIK3CA exon9 mutations associate with reduced survival, and are highly concordant between matching primary tumors and metastases in endometrial cancer. Scientific Reports, 2017, 7, 10240.              | 3.3  | 19        |
| 13 | <i>PIK3CA</i> Amplification Associates with Aggressive Phenotype but Not Markers of AKT-MTOR<br>Signaling in Endometrial Carcinoma. Clinical Cancer Research, 2019, 25, 334-345.                         | 7.0  | 17        |
| 14 | An MRI-Based Radiomic Prognostic Index Predicts Poor Outcome and Specific Genetic Alterations in<br>Endometrial Cancer. Journal of Clinical Medicine, 2021, 10, 538.                                     | 2.4  | 15        |
| 15 | Endometrial cancer cells exhibit high expression of p110l² and its selective inhibition induces variable responses on PI3K signaling, cell survival and proliferation. Oncotarget, 2017, 8, 3881-3894.   | 1.8  | 15        |
| 16 | A radiogenomics application for prognostic profiling of endometrial cancer. Communications Biology, 2021, 4, 1363.   | 4.4  | 14        |
| 17 | Identification of highly connected and differentially expressed gene subnetworks in metastasizing endometrial cancer. PLoS ONE, 2018, 13, e0206665.  | 2.5  | 11        |