

Grigorios Panagiotou

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7511161/grigorios-panagiotou-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

22
papers

1,355
citations

11
h-index

23
g-index

23
ext. papers

1,620
ext. citations

6
avg, IF

4.22
L-index

| # | Paper | IF | Citations |
|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----------|
| 22 | Endocrine manifestations and new developments in mitochondrial disease. <i>Endocrine Reviews</i> , 2021 | 27.2 | 1 |
| 21 | Obesity and COVID-19: A jigsaw puzzle with still missing pieces. <i>Clinical Obesity</i> , 2021 , 11, e12420 | 3.6 | 6 |
| 20 | Serum Levels of Irisin and Omentin-1 in Breast Neoplasms and Their Association with Tumor Histology. <i>International Journal of Endocrinology</i> , 2021 , 2021, 6656671 | 2.7 | 5 |
| 19 | Serum Follistatin Is Increased in Thyroid Cancer and Is Associated With Adverse Tumor Characteristics in Humans. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021 , 106, e2137-e2150 | 5.6 | 2 |
| 18 | Low serum 25-hydroxyvitamin D (25[OH]D) levels in patients hospitalized with COVID-19 are associated with greater disease severity. <i>Clinical Endocrinology</i> , 2020 , 93, 508-511 | 3.4 | 106 |
| 17 | Original publication: Low serum 25-hydroxyvitamin D (25[OH]D) levels in patients hospitalized with COVID-19 are associated with greater disease severity. <i>Clinical Endocrinology</i> , 2020 , 93, 629-630 | 3.4 | 8 |
| 16 | Adipocytes express tissue factor and FVII and are procoagulant in a TF/FVIIa-dependent manner. <i>Upsala Journal of Medical Sciences</i> , 2019 , 124, 158-167 | 2.8 | 2 |
| 15 | Serum Levels of Activins, Follistatins, and Growth Factors in Neoplasms of the Breast: A Case-Control Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 349-358 | 5.6 | 11 |
| 14 | Altered Glucose Uptake in Muscle, Visceral Adipose Tissue, and Brain Predict Whole-Body Insulin Resistance and may Contribute to the Development of Type 2 Diabetes: A Combined PET/MR Study. <i>Hormone and Metabolic Research</i> , 2018 , 50, 627-639 | 3.1 | 23 |
| 13 | Altered Glucose Uptake in Muscle, Visceral Adipose Tissue, and Brain Predict Whole-Body Insulin Resistance and may Contribute to the Development of Type 2 Diabetes: A Combined PET/MR Study. <i>Hormone and Metabolic Research</i> , 2018 , 50, e10 | 3.1 | 4 |
| 12 | Association between lifestyle and anthropometric parameters and thyroid nodule features. <i>Endocrine</i> , 2017 , 56, 560-567 | 4 | 17 |
| 11 | Physiological parameters regulating circulating levels of the IGFBP-4/Stanniocalcin-2/PAPP-A axis. <i>Metabolism: Clinical and Experimental</i> , 2017 , 75, 16-24 | 12.7 | 10 |
| 10 | Changes in Thyroid Hormone Levels Within the Normal and/or Subclinical Hyper- or Hypothyroid Range Do Not Affect Circulating Irisin Levels in Humans. <i>Thyroid</i> , 2016 , 26, 1039-45 | 6.2 | 13 |
| 9 | SERUM ADIPONECTIN AND INSULIN-LIKE GROWTH FACTOR 1 IN PREDOMINANTLY FEMALE PATIENTS WITH THYROID CANCER: ASSOCIATION WITH THE HISTOLOGIC CHARACTERISTICS OF THE TUMOR. <i>Endocrine Practice</i> , 2016 , 22, 68-75 | 3.2 | 15 |
| 8 | Effects of a 1-year exercise and lifestyle intervention on irisin, adipokines, and inflammatory markers in obese children. <i>Obesity</i> , 2014 , 22, 1701-8 | 8 | 81 |
| 7 | Exercise-induced irisin secretion is independent of age or fitness level and increased irisin may directly modulate muscle metabolism through AMPK activation. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, E2154-61 | 5.6 | 193 |
| 6 | Circulating irisin in healthy, young individuals: day-night rhythm, effects of food intake and exercise, and associations with gender, physical activity, diet, and body composition. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, 3247-55 | 5.6 | 102 |

| | | | |
|---|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----|
| 5 | Circulating irisin, omentin-1, and lipoprotein subparticles in adults at higher cardiovascular risk. <i>Metabolism: Clinical and Experimental</i> , 2014 , 63, 1265-71 | 12.7 | 69 |
| 4 | Irisin mRNA and circulating levels in relation to other myokines in healthy and morbidly obese humans. <i>European Journal of Endocrinology</i> , 2013 , 169, 829-34 | 6.5 | 46 |
| 3 | Responses of circulating irisin to different exercises in humans. <i>FASEB Journal</i> , 2013 , 27, 712.17 | 0.9 | |
| 2 | FNDC5 and irisin in humans: I. Predictors of circulating concentrations in serum and plasma and II. mRNA expression and circulating concentrations in response to weight loss and exercise. <i>Metabolism: Clinical and Experimental</i> , 2012 , 61, 1725-38 | 12.7 | 631 |
| 1 | Low serum 25-hydroxyvitamin D (25[OH]D) levels in patients hospitalised with COVID-19 are associated with greater disease severity: results of a local audit of practice | | 10 |