## Robert Lachmann

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7512855/publications.pdf

Version: 2024-02-01

1039880 1199470 1,059 15 9 12 citations h-index g-index papers 16 16 16 1500 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Human-specific gene <i>ARHGAP11B</i> promotes basal progenitor amplification and neocortex expansion. Science, 2015, 347, 1465-1470.	6.0	487
2	Differences and similarities between human and chimpanzee neural progenitors during cerebral cortex development. ELife, 2016, 5, .	2.8	200
3	Posterior brain in fetuses with open spina bifida at 11 to 13 weeks. Prenatal Diagnosis, 2011, 31, 103-106.	1.1	144
4	Threeâ€dimensional sonography of the posterior fossa in fetuses with open spina bifida at 11–13 weeks' gestation. Ultrasound in Obstetrics and Gynecology, 2011, 38, 625-629.	0.9	56
5	Frontomaxillary facial angle in fetuses with spina bifida at 11–13 weeks' gestation. Ultrasound in Obstetrics and Gynecology, 2010, 36, 268-271.	0.9	53
6	Posterior brain in fetuses with Dandy–Walker malformation with complete agenesis of the cerebellar vermis at 11–13 weeks: a pilot study. Prenatal Diagnosis, 2012, 32, 765-769.	1.1	45
7	Explorative investigation of vascular endothelial growth factor receptor expression in primary ovarian cancer and its clinical relevance. Gynecologic Oncology, 2014, 133, 467-472.	0.6	23
8	Isolated Cleft Lip and Palate: Maxillary Gap Sign and Palatino-Maxillary Diameter at 11–13 Weeks. Fetal Diagnosis and Therapy, 2018, 44, 241-246.	0.6	23
9	Midbrain and Falx in Fetuses with Absent Corpus Callosum at 11–13 Weeks. Fetal Diagnosis and Therapy, 2013, 33, 41-46.	0.6	17
10	Signs of Reduced Basal Progenitor Levels and Cortical Neurogenesis in Human Fetuses with Open Spina Bifida at 11–15 Weeks of Gestation. Journal of Neuroscience, 2020, 40, 1766-1777.	1.7	5
11	Correspondence regarding research letter published by Arigita <i>et al.</i> . Prenatal Diagnosis, 2012, 32, 201-201.	1.1	3
12	Comment on "A prospective study on fetal posterior cranial fossa assessment for early detection of open spina bifida at 11–13 weeks―by Kose et al Congenital Anomalies (discontinued), 2018, 58, 77-77.	0.3	2
13	Re: Absent †superimposed†ine†sign: novel marker in early diagnosis of cleft of fetal secondary palate. Ultrasound in Obstetrics and Gynecology, 2021, 58, 954-955.	0.9	1
14	OC20.03: Hypertensive disorders in pregnancy: screening by uterine artery Doppler and blood pressure at 11-13 weeks. Ultrasound in Obstetrics and Gynecology, 2009, 34, 37-37.	0.9	0
15	Clinical relevance of VEGF-receptor status in primary ovarian cancer: A pilot study for future biomarker analyses Journal of Clinical Oncology, 2013, 31, 5556-5556.	0.8	0