

Daniela Cabibi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/751976/daniela-cabibi-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

106
papers

2,756
citations

27
h-index

49
g-index

111
ext. papers

3,257
ext. citations

4.3
avg, IF

4.54
L-index

#	Paper	IF	Citations
106	TargetPlex FFPE-Direct DNA Library Preparation Kit for SiRe NGS panel: an international performance evaluation study. <i>Journal of Clinical Pathology</i> , 2021 ,	3.9	4
105	A Case of Follicular Tumor of Uncertain Malignant Potential (FT-UMP) with Glomeruloid Features Showing Capsular Mucinous Degeneration. <i>Case Reports in Pathology</i> , 2021 , 2021, 1686025	0.9	1
104	Hsp60 Quantification in Human Gastric Mucosa Shows Differences between Pathologies with Various Degrees of Proliferation and Malignancy Grade. <i>Applied Sciences (Switzerland)</i> , 2021 , 11, 3582	2.6	0
103	Mild exacerbation of obesity- and age-dependent liver disease progression by senolytic cocktail dasatinib + quercetin. <i>Cell Communication and Signaling</i> , 2021 , 19, 44	7.5	13
102	Effectiveness and safety of lomitapide in a patient with familial chylomicronemia syndrome. <i>Endocrine</i> , 2021 , 71, 344-350	4	4
101	Not all 557/558 codons mutations have the same prognostic influence on recurrence-free survival: breaking the exon 11 mutations in gastrointestinal stromal tumors (GISTs). <i>Therapeutic Advances in Medical Oncology</i> , 2021 , 13, 17588359211049779	5.4	0
100	Type and Gene Location of Mutations Predict Progression-Free Survival to First-Line Imatinib in Gastrointestinal Stromal Tumors: A Look into the Exon. <i>Cancers</i> , 2021 , 13,	6.6	5
99	Caucasian lean subjects with non-alcoholic fatty liver disease share long-term prognosis of non-lean: time for reappraisal of BMI-driven approach?. <i>Gut</i> , 2021 ,	19.2	27
98	NUPR1 protects liver from lipotoxic injury by improving the endoplasmic reticulum stress response. <i>FASEB Journal</i> , 2021 , 35, e21395	0.9	1
97	Long-term outcomes and predictive ability of non-invasive scoring systems in patients with non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2021 , 75, 786-794	13.4	21
96	The Prevalence of NAFLD and Fibrosis in Bariatric Surgery Patients and the Reliability of Noninvasive Diagnostic Methods. <i>BioMed Research International</i> , 2020 , 2020, 5023157	3	7
95	Current Role of Intraoperative Frozen Section Examination of Sentinel Lymph Node in Early Breast Cancer. <i>Anticancer Research</i> , 2020 , 40, 1711-1717	2.3	8
94	TTF-1/p63-Positive Poorly Differentiated NSCLC: A Histogenetic Hypothesis from the Basal Reserve Cell of the Terminal Respiratory Unit. <i>Diagnostics</i> , 2020 , 10,	3.8	5
93	GDF11 induces mild hepatic fibrosis independent of metabolic health. <i>Aging</i> , 2020 , 12, 20024-20046	5.6	6
92	Fever and dyspnoea in a tracheostomised patient. <i>Breathe</i> , 2020 , 16, 200115	1.8	
91	Hepatic and circulating levels of PCSK9 in morbidly obese patients: Relation with severity of liver steatosis. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2020 , 1865, 158792	5	8
90	Presence of Serum Antinuclear Antibodies Does Not Impact Long-Term Outcomes in Nonalcoholic Fatty Liver Disease. <i>American Journal of Gastroenterology</i> , 2020 , 115, 1289-1292	0.7	5

89	Immunomorphological Pattern of Molecular Chaperones in Normal and Pathological Thyroid Tissues and Circulating Exosomes: Potential Use in Clinics. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	23
88	Glypican-3 and Hep Par-1 are Useful Biomarkers in the Cytologic Assessment of Ascites. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2019 , 27, 65-69	1.9	
87	Evaluating the association of serum ferritin and hepatic iron with disease severity in non-alcoholic fatty liver disease. <i>Liver International</i> , 2019 , 39, 1325-1334	7.9	23
86	Quantitative Immunomorphological Analysis of Heat Shock Proteins in Thyroid Follicular Adenoma and Carcinoma Tissues Reveals Their Potential for Differential Diagnosis and Points to a Role in Carcinogenesis. <i>Applied Sciences (Switzerland)</i> , 2019 , 9, 4324	2.6	3
85	Are Long Noncoding RNAs New Potential Biomarkers in Gastrointestinal Stromal Tumors (GISTs)? The Role of H19 and MALAT1. <i>Journal of Oncology</i> , 2019 , 2019, 5458717	4.5	8
84	Reactive hyperemia index (RHI) and cognitive performance indexes are associated with histologic markers of liver disease in subjects with non-alcoholic fatty liver disease (NAFLD): a case control study. <i>Cardiovascular Diabetology</i> , 2018 , 17, 28	8.7	22
83	Fibronectin Type III Domain-Containing Protein 5 rs3480 A>G Polymorphism, Irisin, and Liver Fibrosis in Patients With Nonalcoholic Fatty Liver Disease. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017 , 102, 2660-2669	5.6	30
82	Gastric and Rectal Metastases from Malignant Melanoma Presenting with Hypochromic Anemia and Treated with Immunotherapy. <i>Case Reports in Oncological Medicine</i> , 2017 , 2017, 2079068	0.9	7
81	Pleural epithelioid angiosarcoma with lymphatic differentiation arisen after radiometabolic therapy for thyroid carcinoma: immunohistochemical findings and review of the literature. <i>Diagnostic Pathology</i> , 2017 , 12, 60	3	4
80	Interferon lambda 4 rs368234815 TT>C variant is associated with liver damage in patients with nonalcoholic fatty liver disease. <i>Hepatology</i> , 2017 , 66, 1885-1893	11.2	59
79	Analysis of tissue and circulating microRNA expression during metaplastic transformation of the esophagus. <i>Oncotarget</i> , 2016 , 7, 47821-47830	3.3	28
78	The Presence of White Matter Lesions Is Associated With the Fibrosis Severity of Nonalcoholic Fatty Liver Disease. <i>Medicine (United States)</i> , 2016 , 95, e3446	1.8	25
77	TM6SF2 rs58542926 is not associated with steatosis and fibrosis in large cohort of patients with genotype 1 chronic hepatitis C. <i>Liver International</i> , 2016 , 36, 198-204	7.9	12
76	CD1A-positive cells and HSP60 (HSPD1) levels in keratoacanthoma and squamous cell carcinoma. <i>Cell Stress and Chaperones</i> , 2016 , 21, 131-137	4	6
75	The hepatic expression of vitamin D receptor is inversely associated with the severity of liver damage in genotype 1 chronic hepatitis C patients. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 193-200	5.6	21
74	Epicardial fat, cardiac geometry and cardiac function in patients with non-alcoholic fatty liver disease: association with the severity of liver disease. <i>Journal of Hepatology</i> , 2015 , 62, 928-33	13.4	115
73	The combination of liver stiffness measurement and NAFLD fibrosis score improves the noninvasive diagnostic accuracy for severe liver fibrosis in patients with nonalcoholic fatty liver disease. <i>Liver International</i> , 2015 , 35, 1566-73	7.9	94
72	Amphiregulin activates human hepatic stellate cells and is upregulated in non alcoholic steatohepatitis. <i>Scientific Reports</i> , 2015 , 5, 8812	4.9	27

71	The severity of steatosis influences liver stiffness measurement in patients with nonalcoholic fatty liver disease. <i>Hepatology</i> , 2015 , 62, 1101-10	11.2	131
70	Comparison of Histochemical Stainings in Evaluation of Liver Fibrosis and Correlation with Transient Elastography in Chronic Hepatitis. <i>Analytical Cellular Pathology</i> , 2015 , 2015, 431750	3.4	13
69	Nodular histiocytic/mesothelial hyperplasia as consequence of chronic mesothelium irritation by subphrenic abscess. <i>Future Oncology</i> , 2015 , 11, 51-5	3.6	2
68	Comparison of Histochemical Staining Methods and Correlation with Transient Elastography in Acute Hepatitis. <i>Pathobiology</i> , 2015 , 82, 48-52	3.6	8
67	Elevated blood Hsp60, its structural similarities and cross-reactivity with thyroid molecules, and its presence on the plasma membrane of oncocytes point to the chaperonin as an immunopathogenic factor in Hashimoto's thyroiditis. <i>Cell Stress and Chaperones</i> , 2014 , 19, 343-53	4	43
66	Clinical features and outcomes of patients with drug-induced autoimmune hepatitis: a retrospective cohort study. <i>Digestive and Liver Disease</i> , 2014 , 46, 1116-20	3.3	28
65	Discordance rate of HER2 status in primary breast carcinomas versus synchronous axillary lymph node metastases: a multicenter retrospective investigation. <i>OncoTargets and Therapy</i> , 2014 , 7, 1267-72	4.4	16
64	Immunohistochemical/histochemical double staining method in the study of the columnar metaplasia of the oesophagus. <i>European Journal of Histochemistry</i> , 2014 , 58, 2326	2.1	6
63	Multimodal surgical and medical treatment for extensive rhinocerebral mucormycosis in an elderly diabetic patient: a case report and literature review. <i>Case Reports in Medicine</i> , 2014 , 2014, 527062	0.7	10
62	Body mass index and liver stiffness affect accuracy of ultrasonography in detecting steatosis in patients with chronic hepatitis C virus genotype 1 infection. <i>Clinical Gastroenterology and Hepatology</i> , 2014 , 12, 878-884.e1	6.9	10
61	Steatosis affects the performance of liver stiffness measurement for fibrosis assessment in patients with genotype 1 chronic hepatitis C. <i>Journal of Hepatology</i> , 2014 , 61, 523-9	13.4	57
60	Glucokinase regulatory protein gene polymorphism affects liver fibrosis in non-alcoholic fatty liver disease. <i>PLoS ONE</i> , 2014 , 9, e87523	3.7	83
59	Industrial, not fruit fructose intake is associated with the severity of liver fibrosis in genotype 1 chronic hepatitis C patients. <i>Journal of Hepatology</i> , 2013 , 59, 1169-76	13.4	28
58	Surgical pathology and the diagnosis of invasive visceral yeast infection: two case reports and literature review. <i>World Journal of Emergency Surgery</i> , 2013 , 8, 38	9.2	19
57	High sCD36 plasma level is associated with steatosis and its severity in patients with genotype 1 chronic hepatitis C. <i>Journal of Viral Hepatitis</i> , 2013 , 20, 174-82	3.4	10
56	PNPLA3 GG genotype and carotid atherosclerosis in patients with non-alcoholic fatty liver disease. <i>PLoS ONE</i> , 2013 , 8, e74089	3.7	47
55	Primary biliary cirrhosis and hereditary hemorrhagic telangiectasia: When two rare diseases coexist. <i>World Journal of Hepatology</i> , 2013 , 5, 288-91	3.4	2
54	Antimitochondrial antibody -M2 positive autoimmune hepatitis during standard of care for chronic hepatitis C. <i>Hepatology Research</i> , 2012 , 42, 428-32	5.1	3

53	Methylenetetrahydrofolate reductase homozygosis and low-density lipoproteins in patients with genotype 1 chronic hepatitis C. <i>Journal of Viral Hepatitis</i> , 2012 , 19, 465-72	3.4	8
52	Malignant tumor-like gastric lesion due to <i>Candida albicans</i> in a diabetic patient treated with cyclosporin: a case report and review of the literature. <i>Clinical and Experimental Medicine</i> , 2012 , 12, 201-5	4.9	5
51	CD10 and HHF35 actin in the differential diagnosis between Collagenous spherulosis and adenoid-cystic carcinoma of the breast. <i>Pathology Research and Practice</i> , 2012 , 208, 405-9	3.4	9
50	Hyperuricaemia: another metabolic feature affecting the severity of chronic hepatitis because of HCV infection. <i>Liver International</i> , 2012 , 32, 1443-50	7.9	15
49	IL28B and PNPLA3 polymorphisms affect histological liver damage in patients with non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2012 , 56, 1356-62	13.4	72
48	Serum γ -glutamyl transferase levels, insulin resistance and liver fibrosis in patients with chronic liver diseases. <i>PLoS ONE</i> , 2012 , 7, e51165	3.7	24
47	Morphological and biomolecular characteristics of subcentimetric invasive breast carcinomas in Sicily: A multicentre retrospective study in relation to trastuzumab treatment. <i>Oncology Letters</i> , 2012 , 3, 141-146	2.6	3
46	Carotid atherosclerosis and chronic hepatitis C: a prospective study of risk associations. <i>Hepatology</i> , 2012 , 55, 1317-23	11.2	105
45	Liver follicular helper T-cells predict the achievement of virological response following interferon-based treatment in HCV-infected patients. <i>Antiviral Therapy</i> , 2012 , 17, 111-8	1.6	8
44	Vitamin D levels and IL28B polymorphisms are related to rapid virological response to standard of care in genotype 1 chronic hepatitis C. <i>Antiviral Therapy</i> , 2012 , 17, 823-31	1.6	28
43	High liver RBP4 protein content is associated with histological features in patients with genotype 1 chronic hepatitis C and with nonalcoholic steatohepatitis. <i>Digestive and Liver Disease</i> , 2011 , 43, 404-10	3.3	24
42	Glut-1 expression and in situ CD1a/CD57 immunologic deficit in keratoacanthoma and squamous cell carcinoma of immunocompetent patients. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2011 , 19, 239-45	1.9	6
41	TyG index, HOMA score and viral load in patients with chronic hepatitis C due to genotype 1. <i>Journal of Viral Hepatitis</i> , 2011 , 18, e372-80	3.4	17
40	Hepatic steatosis and insulin resistance are associated with severe fibrosis in patients with chronic hepatitis caused by HBV or HCV infection. <i>Liver International</i> , 2011 , 31, 507-15	7.9	58
39	Metastatic seeding of colon adenocarcinoma manifesting as synchronous breast and chest wall localization: report of a case. <i>Surgery Today</i> , 2011 , 41, 242-6	3	7
38	Cystic mucinous adenocarcinoma of the lung: a case report. <i>Journal of Cardiothoracic Surgery</i> , 2011 , 6, 128	1.6	6
37	Accuracy of the water-siphon test associated to barium study in a high prevalence gastro-oesophageal reflux disease population: a novel statistical approach. <i>Journal of Evaluation in Clinical Practice</i> , 2010 , 16, 550-5	2.5	1
36	Noninvasive assessment of liver fibrosis in thalassaemia major patients by transient elastography (TE) - lack of interference by iron deposition. <i>British Journal of Haematology</i> , 2010 , 148, 476-9	4.5	45

35	Intrahepatic IgG/IgM plasma cells ratio helps in classifying autoimmune liver diseases. <i>Digestive and Liver Disease</i> , 2010 , 42, 585-92	3.3	24
34	BRAF(V600E) mutation influences hypoxia-inducible factor-1alpha expression levels in papillary thyroid cancer. <i>Modern Pathology</i> , 2010 , 23, 1052-60	9.8	38
33	Post-bronchoscopy fatal endobronchial hemorrhage in a woman with bronchopulmonary mucormycosis: a case report. <i>Journal of Medical Case Reports</i> , 2010 , 4, 398	1.2	4
32	The value of intraoperative frozen section examination of sentinel lymph nodes in surgical management of breast carcinoma. <i>Langenbeck's Archives of Surgery</i> , 2010 , 395, 685-91	3.4	13
31	Low vitamin D serum level is related to severe fibrosis and low responsiveness to interferon-based therapy in genotype 1 chronic hepatitis C. <i>Hepatology</i> , 2010 , 51, 1158-67	11.2	318
30	Visceral adiposity index is associated with histological findings and high viral load in patients with chronic hepatitis C due to genotype 1. <i>Hepatology</i> , 2010 , 52, 1543-52	11.2	76
29	Pyogenic granuloma-like Kaposi's sarcoma on the hands: immunohistochemistry and human herpesvirus-8 detection. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2009 , 23, 587-9	4.6	9
28	Keratin 7 expression as an early marker of reflux-related columnar mucosa without intestinal metaplasia in the esophagus. <i>Medical Science Monitor</i> , 2009 , 15, CR203-210	3.2	10
27	Diagnostic and prognostic value of CD10 in peripheral nerve sheath tumors. <i>Anticancer Research</i> , 2009 , 29, 3149-55	2.3	6
26	Differing expression of metalloprotease and of adhesion molecules in signet-ring cell and intestinal colorectal carcinoma. <i>Anticancer Research</i> , 2009 , 29, 4417-22	2.3	10
25	Time course of insulin resistance during antiviral therapy in non-diabetic, non-cirrhotic patients with genotype 1 HCV infection. <i>Antiviral Therapy</i> , 2009 , 14, 631-639	1.6	18
24	Liver eosinophilic infiltrate is a significant finding in patients with chronic hepatitis C. <i>Journal of Viral Hepatitis</i> , 2008 , 15, 523-30	3.4	15
23	Solitary necrotic nodule of the liver: different pathological findings express a different histogenesis. <i>Case Reports in Gastroenterology</i> , 2008 , 2, 149-54	1	9
22	Insulin resistance and diabetes increase fibrosis in the liver of patients with genotype 1 HCV infection. <i>American Journal of Gastroenterology</i> , 2008 , 103, 1136-44	0.7	149
21	The oral melanoma: atypical immunophenotype in the metastatic area. <i>Melanoma Research</i> , 2008 , 18, 70-2	3.3	
20	Liver disease in chelated transfusion-dependent thalasseemics: the role of iron overload and chronic hepatitis C. <i>Haematologica</i> , 2008 , 93, 1243-6	6.6	41
19	The accuracy of sentinel lymph node biopsy in the treatment of multicentric invasive breast cancer using a subareolar injection of tracer. <i>World Journal of Surgery</i> , 2008 , 32, 2483-7	3.3	16
18	Increased expression of transketolase-like-1 in papillary thyroid carcinomas smaller than 1.5 cm in diameter is associated with lymph-node metastases. <i>Cancer</i> , 2008 , 113, 936-44	6.4	32

17	Retinol-binding protein 4: a new marker of virus-induced steatosis in patients infected with hepatitis c virus genotype 1. <i>Hepatology</i> , 2008 , 48, 28-37	11.2	49
16	Autoimmune hepatitis following Epstein-Barr virus infection. <i>BMJ Case Reports</i> , 2008 , 2008, bcr0620080071	0.4	13
15	Cervico-Facial Actinomycosis: Epidemiological and Clinical Comments. <i>American Journal of Infectious Diseases</i> , 2008 , 4, 204-208	0.4	6
14	BRAF V600E mutation and p27 kip1 expression in papillary carcinomas of the thyroid . <i>Cancer</i> , 2007 , 110, 1218-26	6.4	73
13	Re: Low-grade fibromyxoid sarcoma arising in the big toe. <i>Pathology International</i> , 2007 , 57, 458; author reply 459-60	1.8	2
12	Associations between Notch-2, Akt-1 and HER2/neu expression in invasive human breast cancer: a tissue microarray immunophenotypic analysis on 98 patients. <i>Pathobiology</i> , 2007 , 74, 317-22	3.6	19
11	Immunohistochemistry differentiates papillary thyroid carcinoma arising in ectopic thyroid tissue from secondary lymph node metastases. <i>Thyroid</i> , 2007 , 17, 603-7	6.2	5
10	Fine-needle aspiration molecular analysis for the diagnosis of papillary thyroid carcinoma through BRAF V600E mutation and RET/PTC rearrangement. <i>Thyroid</i> , 2007 , 17, 1109-15	6.2	81
9	Carcinosarcoma of monoclonal origin arising in a dermoid cyst of ovary: a case report. <i>BMC Cancer</i> , 2006 , 6, 47	4.8	20
8	Unexpectedly high frequency of genital involvement in women with clinical and histological features of oral lichen planus. <i>Acta Dermato-Venereologica</i> , 2006 , 86, 433-8	2.2	20
7	Thyroid sarcoidosis as a unique localization. <i>Thyroid</i> , 2006 , 16, 1175-7	6.2	11
6	Secretory breast carcinoma with metastatic sentinel lymph node. <i>World Journal of Surgical Oncology</i> , 2006 , 4, 88	3.4	35
5	Difference in Ki67 and thymidylate synthase expression in primary tumour compared with metastatic nodes in breast cancer patients. <i>Nucleosides, Nucleotides and Nucleic Acids</i> , 2006 , 25, 1193-6	1.4	5
4	Solid variant of mammary "adenoid cystic carcinoma with basaloid features" merging with "small cell carcinoma". <i>Pathology Research and Practice</i> , 2005 , 201, 705-11	3.4	18
3	Overexpression of cyclin D1 and interaction between p27Kip1 and tumour thickness predict lymph node metastases occurrence in lower lip squamous cell carcinoma. <i>Oral Oncology</i> , 2005 , 41, 268-75	4.4	7
2	A case of bowel schistosomiasis not adhering to endoscopic findings. <i>World Journal of Gastroenterology</i> , 2005 , 11, 7044-7	5.6	8
1	Expression of cytokeratin 7 and 20 in pathological conditions of the bile tract. <i>Pathology Research and Practice</i> , 2003 , 199, 65-70	3.4	16