

Fu-xing Pei

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7531824/fu-xing-pei-publications-by-citations.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

66

papers

1,606

citations

23

h-index

38

g-index

75

ext. papers

2,056

ext. citations

3.9

avg, IF

5.08

L-index

#	Paper	IF	Citations
66	Multiple Boluses of Intravenous Tranexamic Acid to Reduce Hidden Blood Loss After Primary Total Knee Arthroplasty Without Tourniquet: A Randomized Clinical Trial. <i>Journal of Arthroplasty</i> , 2016 , 31, 2458-2464	4.4	112
65	Topical application of tranexamic acid in primary total hip arthroplasty: a randomized double-blind controlled trial. <i>Journal of Arthroplasty</i> , 2014 , 29, 2452-6	4.4	111
64	Combination of intravenous and topical application of tranexamic acid in primary total knee arthroplasty: a prospective randomized controlled trial. <i>Journal of Arthroplasty</i> , 2014 , 29, 2342-6	4.4	97
63	Comparison of Early Functional Recovery After Total Hip Arthroplasty Using a Direct Anterior or Posterolateral Approach: A Randomized Controlled Trial. <i>Journal of Arthroplasty</i> , 2017 , 32, 3421-3428	4.4	93
62	Long-Term Results of Cementless Total Hip Arthroplasty With Subtrochanteric Shortening Osteotomy in Crowe Type IV Developmental Dysplasia. <i>Journal of Arthroplasty</i> , 2017 , 32, 1211-1219	4.4	84
61	Intravenous and Topical Tranexamic Acid Alone Are Superior to Tourniquet Use for Primary Total Knee Arthroplasty: A Prospective, Randomized Controlled Trial. <i>Journal of Bone and Joint Surgery - Series A</i> , 2017 , 99, 2053-2061	5.6	84
60	The efficacy and safety of multiple-dose intravenous tranexamic acid on blood loss following total knee arthroplasty: a randomized controlled trial. <i>International Orthopaedics</i> , 2017 , 41, 2053-2059	3.8	55
59	Oral vs Intravenous vs Topical Tranexamic Acid in Primary Hip Arthroplasty: A Prospective, Randomized, Double-Blind, Controlled Study. <i>Journal of Arthroplasty</i> , 2018 , 33, 786-793	4.4	55
58	Plasma Fibrin Degradation Product and D-Dimer Are of Limited Value for Diagnosing Periprosthetic Joint Infection. <i>Journal of Arthroplasty</i> , 2019 , 34, 2454-2460	4.4	49
57	Adductor canal block versus femoral nerve block for total knee arthroplasty: a meta-analysis of randomized controlled trials. <i>Scientific Reports</i> , 2017 , 7, 40721	4.9	46
56	Comparison of intravenous versus topical tranexamic acid in primary total hip and knee arthroplasty: An updated meta-analysis. <i>Thrombosis Research</i> , 2017 , 153, 28-36	8.2	46
55	Does tranexamic acid alter the risk of thromboembolism following primary total knee arthroplasty with sequential earlier anticoagulation? A large, single center, prospective cohort study of consecutive cases. <i>Thrombosis Research</i> , 2015 , 136, 234-8	8.2	39
54	Risk factors for venous thromboembolism of total hip arthroplasty and total knee arthroplasty: a systematic review of evidences in ten years. <i>BMC Musculoskeletal Disorders</i> , 2015 , 16, 24	2.8	36
53	Two doses of low-dose perioperative dexamethasone improve the clinical outcome after total knee arthroplasty: a randomized controlled study. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2018 , 26, 1549-1556	5.5	33
52	The efficacy and safety of two low-dose peri-operative dexamethasone on pain and recovery following total hip arthroplasty: a randomized controlled trial. <i>International Orthopaedics</i> , 2018 , 42, 499-505	3.8	33
51	Single-dose tranexamic acid for reducing bleeding and transfusions in total hip arthroplasty: A double-blind, randomized controlled trial of different doses. <i>Thrombosis Research</i> , 2016 , 141, 119-23	8.2	32
50	Closed Suction Drainage Is Not Associated with Faster Recovery after Total Knee Arthroplasty: A Prospective Randomized Controlled Study of 80 Patients. <i>Orthopaedic Surgery</i> , 2016 , 8, 226-33	2.5	30

49	Multiple-Dose Intravenous Tranexamic Acid Further Reduces Hidden Blood Loss After Total Hip Arthroplasty: A Randomized Controlled Trial. <i>Journal of Arthroplasty</i> , 2018 , 33, 2940-2945	4.4	30
48	Multiple Doses of Perioperative Dexamethasone Further Improve Clinical Outcomes After Total Knee Arthroplasty: A Prospective, Randomized, Controlled Study. <i>Journal of Arthroplasty</i> , 2018 , 33, 3448-3454	4.4	29
47	Correlations between inflammatory cytokines, muscle damage markers and acute postoperative pain following primary total knee arthroplasty. <i>BMC Musculoskeletal Disorders</i> , 2017 , 18, 265	2.8	28
46	Comparison of Early and Delayed Open Reduction and Internal Fixation for Treating Closed Tibial Pilon Fractures. <i>Foot and Ankle International</i> , 2014 , 35, 657-64	3.3	27
45	Subtrochanteric shortening osteotomy during cementless total hip arthroplasty in young patients with severe developmental dysplasia of the hip. <i>BMC Musculoskeletal Disorders</i> , 2017 , 18, 491	2.8	25
44	Efficacy of oral tranexamic acid on blood loss in primary total hip arthroplasty using a direct anterior approach: a prospective randomized controlled trial. <i>International Orthopaedics</i> , 2018 , 42, 2535-2542	3.8	24
43	The effect of oral versus intravenous tranexamic acid in reducing blood loss after primary total hip arthroplasty: A randomized clinical trial. <i>Thrombosis Research</i> , 2018 , 164, 48-53	8.2	23
42	Multiple Low-Dose Dexamethasone Further Improves Clinical Outcomes Following Total Hip Arthroplasty. <i>Journal of Arthroplasty</i> , 2018 , 33, 1426-1431	4.4	22
41	Preoperative sleep quality affects postoperative pain and function after total joint arthroplasty: a prospective cohort study. <i>Journal of Orthopaedic Surgery and Research</i> , 2019 , 14, 378	2.8	22
40	Finding the Optimal Regimen for Oral Tranexamic Acid Administration in Primary Total Hip Arthroplasty: A Randomized Controlled Trial. <i>Journal of Bone and Joint Surgery - Series A</i> , 2019 , 101, 438-445	5.6	19
39	Effectiveness and safety of an optimized blood management program in total hip and knee arthroplasty: A large, single-center, retrospective study. <i>Medicine (United States)</i> , 2018 , 97, e9429	1.8	18
38	Roles of microRNA and signaling pathway in osteoarthritis pathogenesis. <i>Journal of Zhejiang University: Science B</i> , 2016 , 17, 200-8	4.5	18
37	The efficacy and safety of multiple-dose oral tranexamic acid on blood loss following total hip arthroplasty: a randomized controlled trial. <i>International Orthopaedics</i> , 2019 , 43, 299-305	3.8	18
36	Multicenter Study on Observation of Acute-phase Responses After Infusion of Zoledronic Acid 5 mg in Chinese Women with Postmenopausal Osteoporosis. <i>Orthopaedic Surgery</i> , 2017 , 9, 284-289	2.5	17
35	The antifibrinolytic and anti-inflammatory effects of a high initial-dose tranexamic acid in total knee arthroplasty: a randomized controlled trial. <i>International Orthopaedics</i> , 2020 , 44, 477-486	3.8	17
34	Effect of Multiple Doses of Oral Tranexamic Acid on Haemostasis and Inflammatory Reaction in Total Hip Arthroplasty: A Randomized Controlled Trial. <i>Thrombosis and Haemostasis</i> , 2019 , 119, 92-103	7	17
33	Closed suction drainage following routine primary total joint arthroplasty is associated with a higher transfusion rate and longer postoperative length of stay: a retrospective cohort study. <i>Journal of Orthopaedic Surgery and Research</i> , 2019 , 14, 163	2.8	16
32	Oral tranexamic acid is equivalent to topical tranexamic acid without drainage in primary total hip arthroplasty: A double-blind randomized clinical trial. <i>Thrombosis Research</i> , 2018 , 167, 1-5	8.2	14

31	Six-Dose Intravenous Tranexamic Acid Regimen Further Inhibits Postoperative Fibrinolysis and Reduces Hidden Blood Loss following Total Knee Arthroplasty. <i>Journal of Knee Surgery</i> , 2021 , 34, 224-232 ^{2,4}		14
30	Efficacy and safety of multiple boluses of oral versus intravenous tranexamic acid at reducing blood loss after primary total knee arthroplasty without a tourniquet: A prospective randomized clinical trial. <i>Thrombosis Research</i> , 2018 , 171, 68-73	8.2	13
29	Total hospital cost, length of stay, and complications between simultaneous and staged bilateral total hip arthroplasty: A nationwide retrospective cohort study in China. <i>Medicine (United States)</i> , 2019 , 98, e14687	1.8	12
28	The Use of Porous Tantalum Augments for the Reconstruction of Acetabular Defect in Primary Total Hip Arthroplasty. <i>Journal of Arthroplasty</i> , 2018 , 33, 453-459	4.4	12
27	Comparison of oral versus intra-articular tranexamic acid in enhanced-recovery primary total knee arthroplasty without tourniquet application: a randomized controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2018 , 19, 85	2.8	12
26	Intravenous and subsequent long-term oral tranexamic acid in enhanced-recovery primary total knee arthroplasty without the application of a tourniquet: a randomized placebo-controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2019 , 20, 478	2.8	12
25	Tranexamic acid in primary total knee arthroplasty without tourniquet: a randomized, controlled trial of oral versus intravenous versus topical administration. <i>Scientific Reports</i> , 2018 , 8, 13579	4.9	12
24	Intravenous Combined with Topical Administration of Tranexamic Acid in Primary Total Hip Arthroplasty: A Randomized Controlled Trial. <i>Orthopaedic Surgery</i> , 2017 , 9, 174-179	2.5	11
23	The efficacy and safety of tranexamic acid for reducing blood loss following simultaneous bilateral total knee arthroplasty: a multicenter retrospective study. <i>BMC Musculoskeletal Disorders</i> , 2019 , 20, 325 ^{2,8}	2.8	11
22	Long-Term Results of Cementless Total Hip Arthroplasty for Patients With High Hip Dislocation After Childhood Pyogenic Infection. <i>Journal of Arthroplasty</i> , 2019 , 34, 2420-2426	4.4	10
21	Additional benefits of multiple-dose tranexamic acid to anti-fibrinolysis and anti-inflammation in total knee arthroplasty: a randomized controlled trial. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2020 , 140, 1087-1095	3.6	9
20	Combination of erythropoietin and tranexamic acid in bilateral simultaneous total hip arthroplasty: a randomised, controlled trial. <i>HIP International</i> , 2016 , 26, 331-7	1.7	8
19	Accuracy of a Simple Digital Templating in Primary Uncemented Total Hip Arthroplasty. <i>Chinese Medical Sciences Journal</i> , 2015 , 30, 150-5	1.3	6
18	Effects of Total Hip Arthroplasty on Axial Alignment of the Lower Limb in Patients with Unilateral Developmental Hip Dysplasia (Crowe type IV). <i>Journal of Arthroplasty</i> , 2019 , 34, 2406-2414	4.4	5
17	Repeat Doses of Dexamethasone up to 48 Hours Further Reduce Pain and Inflammation After Total Hip Arthroplasty: A Randomized Controlled Trial. <i>Journal of Arthroplasty</i> , 2020 , 35, 3223-3229	4.4	5
16	Monobloc implants in cementless total hip arthroplasty in patients with Legg-Calve-Perthes disease: a long-term follow-up. <i>BMC Musculoskeletal Disorders</i> , 2017 , 18, 386	2.8	5
15	Benefits of early ambulation within 24 h after total knee arthroplasty: a multicenter retrospective cohort study in China. <i>Military Medical Research</i> , 2021 , 8, 17	19.3	4
14	Is a split-dose intravenous dexamethasone regimen superior to a single high dose in reducing pain and improving function after total hip arthroplasty? A randomized blinded placebo-controlled trial. <i>Bone and Joint Journal</i> , 2020 , 102-B, 1497-1504	5.6	3

13	Comparison of cortisol and inflammatory response between aged and middle-aged patients undergoing total hip arthroplasty: a prospective observational study. <i>BMC Musculoskeletal Disorders</i> , 2017 , 18, 541	2.8	3
12	Finding the optimal control level of intraoperative blood pressure in no tourniquet primary total knee arthroplasty combine with tranexamic acid: a retrospective cohort study which supports the enhanced recovery strategy. <i>Journal of Orthopaedic Surgery and Research</i> , 2020 , 15, 350	2.8	3
11	Role of D-dimer and Fibrinogen in the Diagnosis of Periprosthetic Joint Infection: A Systematic Review and Meta-Analysis. <i>Orthopaedic Surgery</i> , 2021 , 13, 692-700	2.5	3
10	Intraoperative Femoral Condylar Fracture during Primary Total Knee Arthroplasty: Report of Two Cases. <i>Orthopaedic Surgery</i> , 2015 , 7, 180-4	2.5	2
9	Long-term efficacy of repeated sodium hyaluronate injections in adult patients with Kashin-Beck disease of the knee. <i>International Journal of Rheumatic Diseases</i> , 2019 , 22, 392-398	2.3	2
8	Risk factors of opioid use associated with an enhanced-recovery programme after total knee arthroplasty.. <i>BMC Musculoskeletal Disorders</i> , 2021 , 22, 1046	2.8	2
7	One-Stage Arthroplasty or Revision for Seronegative Infections in Hip and Knee. <i>Orthopaedic Surgery</i> , 2020 , 12, 38-49	2.5	1
6	Effectiveness of preemptive antifibrinolysis with tranexamic acid in rheumatoid arthritis patients undergoing total knee arthroplasty: a study protocol for a randomized controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2020 , 21, 465	2.8	1
5	Is there a maximal effect of tranexamic acid in patients undergoing total knee arthroplasty? A randomized controlled trial. <i>MedComm</i> , 2020 , 1, 219-227	2.2	1
4	Dose optimization of intravenous dexamethasone for total knee arthroplasty: when two is not better than one. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2021 , 1	3.6	1
3	Is there a role of tranexamic acid in rheumatoid arthritis with total knee arthroplasty? Findings from a multicenter prospective cohort study in China. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2021 , 141, 489-496	3.6	1
2	Negative pressure pulmonary edema after percutaneous endoscopic interlaminar lumbar discectomy-a case report. <i>BMC Musculoskeletal Disorders</i> , 2018 , 19, 401	2.8	1
1	Oral mosapride can provide additional anti-emetic efficacy following total joint arthroplasty under general anesthesia: a randomized, double-blinded clinical trial. <i>BMC Anesthesiology</i> , 2020 , 20, 297	2.4	