Steven J Rothstein

List of Publications by Citations

Source: https://exaly.com/author-pdf/7533647/steven-j-rothstein-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

149 papers

11,302 citations

61 h-index

103 g-index

151 ext. papers

12,929 ext. citations

7.6 avg, IF

6.13 L-index

#	Paper	IF	Citations
149	A restriction enzyme cleavage map of Tn5 and location of a region encoding neomycin resistance. <i>Molecular Genetics and Genomics</i> , 1979 , 177, 65-72		530
148	Tung tree DGAT1 and DGAT2 have nonredundant functions in triacylglycerol biosynthesis and are localized to different subdomains of the endoplasmic reticulum. <i>Plant Cell</i> , 2006 , 18, 2294-313	11.6	393
147	Molecular cloning of complementary DNA encoding the lignin-forming peroxidase from tobacco: Molecular analysis and tissue-specific expression. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987 , 84, 7542-6	11.5	371
146	C6-volatiles derived from the lipoxygenase pathway induce a subset of defense-related genes. <i>Plant Journal</i> , 1998 , 16, 561-9	6.9	348
145	Understanding plant response to nitrogen limitation for the improvement of crop nitrogen use efficiency. <i>Journal of Experimental Botany</i> , 2011 , 62, 1499-509	7	332
144	Control of inflorescence architecture in Antirrhinum. <i>Nature</i> , 1996 , 379, 791-7	50.4	331
143	Heat induces the splicing by IRE1 of a mRNA encoding a transcription factor involved in the unfolded protein response in Arabidopsis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 7247-52	11.5	300
142	Genetic regulation by NLA and microRNA827 for maintaining nitrate-dependent phosphate homeostasis in arabidopsis. <i>PLoS Genetics</i> , 2011 , 7, e1002021	6	256
141	Arabidopsis homolog of the yeast TREX-2 mRNA export complex: components and anchoring nucleoporin. <i>Plant Journal</i> , 2010 , 61, 259-70	6.9	241
140	Tissue specificity of tobacco peroxidase isozymes and their induction by wounding and tobacco mosaic virus infection. <i>Plant Physiology</i> , 1987 , 84, 438-42	6.6	224
139	A mutation in NLA, which encodes a RING-type ubiquitin ligase, disrupts the adaptability of Arabidopsis to nitrogen limitation. <i>Plant Journal</i> , 2007 , 50, 320-37	6.9	218
138	The relationship between anion exchange and net anion flow across the human red blood cell membrane. <i>Journal of General Physiology</i> , 1977 , 69, 363-86	3.4	216
137	The S-locus receptor kinase gene in a self-incompatible Brassica napus line encodes a functional serine/threonine kinase. <i>Plant Cell</i> , 1992 , 4, 1273-81	11.6	208
136	The inverted repeats of Tn5 are functionally different. <i>Cell</i> , 1980 , 19, 795-805	56.2	198
135	Expression of allene oxide synthase determines defense gene activation in tomato. <i>Plant Physiology</i> , 2000 , 122, 1335-42	6.6	159
134	SAUR39, a small auxin-up RNA gene, acts as a negative regulator of auxin synthesis and transport in rice. <i>Plant Physiology</i> , 2009 , 151, 691-701	6.6	157
133	ABCG transporters are required for suberin and pollen wall extracellular barriers in Arabidopsis. <i>Plant Cell</i> , 2014 , 26, 3569-88	11.6	156

(2012-2007)

132	Genome-wide analysis of Arabidopsis responsive transcriptome to nitrogen limitation and its regulation by the ubiquitin ligase gene NLA. <i>Plant Molecular Biology</i> , 2007 , 65, 775-97	4.6	152
131	Adaptation of Arabidopsis to nitrogen limitation involves induction of anthocyanin synthesis which is controlled by the NLA gene. <i>Journal of Experimental Botany</i> , 2008 , 59, 2933-44	7	151
130	Global transcription profiling reveals differential responses to chronic nitrogen stress and putative nitrogen regulatory components in Arabidopsis. <i>BMC Genomics</i> , 2007 , 8, 281	4.5	151
129	Biological functions of asparagine synthetase in plants. <i>Plant Science</i> , 2010 , 179, 141-153	5.3	148
128	The Genetics of Nitrogen Use Efficiency in Crop Plants. <i>Annual Review of Genetics</i> , 2015 , 49, 269-89	14.5	146
127	Genetic analysis of Arabidopsis GATA transcription factor gene family reveals a nitrate-inducible member important for chlorophyll synthesis and glucose sensitivity. <i>Plant Journal</i> , 2005 , 44, 680-92	6.9	144
126	The role of epigenetic processes in controlling flowering time in plants exposed to stress. <i>Journal of Experimental Botany</i> , 2011 , 62, 3727-35	7	139
125	The functional differences in the inverted repeats of Tn5 are caused by a single base pair nonhomology. <i>Cell</i> , 1981 , 23, 191-9	56.2	133
124	Genome-wide identification of microRNAs in response to low nitrate availability in maize leaves and roots. <i>PLoS ONE</i> , 2011 , 6, e28009	3.7	127
123	Isolation of cDNA clones coding for spinach nitrite reductase: complete sequence and nitrate induction. <i>Molecular Genetics and Genomics</i> , 1988 , 212, 20-6		123
122	The Response of Leaf Photosynthesis and Dry Matter Accumulation to Nitrogen Supply in an Older and a Newer Maize Hybrid. <i>Crop Science</i> , 2008 , 48, 656-665	2.4	122
121	Secretion of a wheat Emylase expressed in yeast. <i>Nature</i> , 1984 , 308, 662-665	50.4	122
120	AMT1;1 transgenic rice plants with enhanced NH4(+) permeability show superior growth and higher yield under optimal and suboptimal NH4(+) conditions. <i>Journal of Experimental Botany</i> , 2014 , 65, 965-79	9 7	121
119	Increased nitrogen-use efficiency in transgenic rice plants over-expressing a nitrogen-responsive early nodulin gene identified from rice expression profiling. <i>Plant, Cell and Environment</i> , 2009 , 32, 1749-	-80 ⁴	112
118	Characterization of Antisense Transformed Plants Deficient in the Tobacco Anionic Peroxidase. <i>Plant Physiology</i> , 1997 , 114, 1187-1196	6.6	108
117	Molecular characterization of an Arabidopsis gene encoding hydroperoxide lyase, a cytochrome P-450 that is wound inducible. <i>Plant Physiology</i> , 1998 , 117, 1393-400	6.6	107
116	Peroxidase-Induced Wilting in Transgenic Tobacco Plants. <i>Plant Cell</i> , 1990 , 2, 7-18	11.6	107
115	The rice R2R3-MYB transcription factor OsMYB55 is involved in the tolerance to high temperature and modulates amino acid metabolism. <i>PLoS ONE</i> , 2012 , 7, e52030	3.7	104

114	The APETALA-2-like transcription factor OsAP2-39 controls key interactions between abscisic acid and gibberellin in rice. <i>PLoS Genetics</i> , 2010 , 6, e1001098	6	101
113	Arabidopsis thaliana GPAT8 and GPAT9 are localized to the ER and possess distinct ER retrieval signals: functional divergence of the dilysine ER retrieval motif in plant cells. <i>Plant Physiology and Biochemistry</i> , 2009 , 47, 867-79	5.4	98
112	NIN-like protein 8 is a master regulator of nitrate-promoted seed germination in Arabidopsis. <i>Nature Communications</i> , 2016 , 7, 13179	17.4	95
111	Molecular cloning of complementary DNA encoding maize nitrite reductase: molecular analysis and nitrate induction. <i>Plant Physiology</i> , 1988 , 88, 741-6	6.6	93
110	Stable and heritable inhibition of the expression of nopaline synthase in tobacco expressing antisense RNA. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987 , 84, 8439-43	11.5	91
109	Genome-wide expression profiling of maize in response to individual and combined water and nitrogen stresses. <i>BMC Genomics</i> , 2013 , 14, 3	4.5	89
108	GNC and CGA1 modulate chlorophyll biosynthesis and glutamate synthase (GLU1/Fd-GOGAT) expression in Arabidopsis. <i>PLoS ONE</i> , 2011 , 6, e26765	3.7	87
107	Arabidopsis PEROXIN11c-e, FISSION1b, and DYNAMIN-RELATED PROTEIN3A cooperate in cell cycle-associated replication of peroxisomes. <i>Plant Cell</i> , 2008 , 20, 1567-85	11.6	87
106	The cloning of two tomato lipoxygenase genes and their differential expression during fruit ripening. <i>Plant Physiology</i> , 1994 , 106, 109-18	6.6	85
105	Promoter cassettes, antibiotic-resistance genes, and vectors for plant transformation. <i>Gene</i> , 1987 , 53, 153-61	3.8	85
104	Expression of OsMYB55 in maize activates stress-responsive genes and enhances heat and drought tolerance. <i>BMC Genomics</i> , 2016 , 17, 312	4.5	83
103	Structural and transcriptional comparative analysis of the S locus regions in two self-incompatible Brassica napus lines. <i>Plant Cell</i> , 1999 , 11, 2217-31	11.6	82
102	Cloning of tomato (Lycopersicon esculentum Mill.) arginine decarboxylase gene and its expression during fruit ripening. <i>Plant Physiology</i> , 1993 , 103, 829-34	6.6	80
101	Reappraisal of nitrogen use efficiency in rice overexpressing glutamine synthetase1. <i>Physiologia Plantarum</i> , 2011 , 141, 361-72	4.6	77
100	Molecular characterization of the S locus in two self-incompatible Brassica napus lines. <i>Plant Cell</i> , 1996 , 8, 2369-80	11.6	75
99	Improving yield potential in crops under elevated CO(2): Integrating the photosynthetic and nitrogen utilization efficiencies. <i>Frontiers in Plant Science</i> , 2012 , 3, 162	6.2	74
98	Transformation of a partial nopaline synthase gene into tobacco suppresses the expression of a resident wild-type gene. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991 , 88, 1770-4	11.5	71
97	Over-expression of STP13, a hexose transporter, improves plant growth and nitrogen use in Arabidopsis thaliana seedlings. <i>Plant, Cell and Environment</i> , 2009 , 32, 271-85	8.4	70

(1977-2009)

96	Assimilation of excess ammonium into amino acids and nitrogen translocation in Arabidopsis thalianaroles of glutamate synthases and carbamoylphosphate synthetase in leaves. <i>FEBS Journal</i> , 2009 , 276, 4061-76	5.7	70	
95	Expression of Nuclear and Chloroplast Photosynthesis-Specific Genes During Leaf Senescence. Journal of Experimental Botany, 1991 , 42, 801-811	7	7º	
94	ROS Induces Anthocyanin Production Via Late Biosynthetic Genes and Anthocyanin Deficiency Confers the Hypersensitivity to ROS-Generating Stresses in Arabidopsis. <i>Plant and Cell Physiology</i> , 2017 , 58, 1364-1377	4.9	67	
93	Rice cytokinin GATA transcription Factor1 regulates chloroplast development and plant architecture. <i>Plant Physiology</i> , 2013 , 162, 132-44	6.6	65	
92	OsPIN5b modulates rice (Oryza sativa) plant architecture and yield by changing auxin homeostasis, transport and distribution. <i>Plant Journal</i> , 2015 , 83, 913-25	6.9	63	
91	A developmental transcriptional network for maize defines coexpression modules. <i>Plant Physiology</i> , 2013 , 161, 1830-43	6.6	63	
90	The Arabidopsis halophytic relative Thellungiella halophila tolerates nitrogen-limiting conditions by maintaining growth, nitrogen uptake, and assimilation. <i>Plant Physiology</i> , 2008 , 147, 1168-80	6.6	63	
89	Regulation of the Accumulation and Reduction of Nitrate by Nitrogen and Carbon Metabolites in Maize Seedlings. <i>Plant Physiology</i> , 1997 , 114, 583-589	6.6	62	
88	Asparagine Metabolic Pathways in Arabidopsis. Plant and Cell Physiology, 2016, 57, 675-89	4.9	61	
87	Effect of light/dark cycles on expression of nitrate assimilatory genes in maize shoots and roots. <i>Plant Physiology</i> , 1991 , 95, 281-5	6.6	59	
86	MicroRNA-mediated repression of the seed maturation program during vegetative development in Arabidopsis. <i>PLoS Genetics</i> , 2012 , 8, e1003091	6	55	
85	Synergistic repression of the embryonic programme by SET DOMAIN GROUP 8 and EMBRYONIC FLOWER 2 in Arabidopsis seedlings. <i>Journal of Experimental Botany</i> , 2012 , 63, 1391-404	7	55	
84	Two Members of the Thioredoxin-h Family Interacts with the Kinase Domain of a Brassica S Locus Receptor Kinase. <i>Plant Cell</i> , 1996 , 8, 1641	11.6	55	
83	The self-incompatibility phenotype in brassica is altered by the transformation of a mutant S locus receptor kinase. <i>Plant Cell</i> , 1998 , 10, 209-18	11.6	55	
82	Structure and regulation of ferredoxin-dependent glutamase synthase from Arabidopsis thaliana. Cloning of cDNA expression in different tissues of wild-type and gltS mutant strains, and light induction. <i>FEBS Journal</i> , 1997 , 243, 708-18		54	
81	Footprinting of the spinach nitrite reductase gene promoter reveals the preservation of nitrate regulatory elements between fungi and higher plants. <i>Plant Molecular Biology</i> , 1997 , 34, 465-76	4.6	53	
80	Isolation of the spinach nitrite reductase gene promoter which confers nitrate inducibility on GUS gene expression in transgenic tobacco. <i>Plant Molecular Biology</i> , 1991 , 17, 9-18	4.6	52	
79	RNA polymerase binding sites in lambdaplac5 DNA. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1977 , 74, 4914-8	11.5	52	

78	The Transcription Factor ANAC032 Represses Anthocyanin Biosynthesis in Response to High Sucrose and Oxidative and Abiotic Stresses. <i>Frontiers in Plant Science</i> , 2016 , 7, 1548	6.2	50
77	Returning to our roots: making plant biology research relevant to future challenges in agriculture. <i>Plant Cell</i> , 2007 , 19, 2695-9	11.6	48
76	AtMBD9: a protein with a methyl-CpG-binding domain regulates flowering time and shoot branching in Arabidopsis. <i>Plant Journal</i> , 2006 , 46, 282-96	6.9	48
75	Purification and partial characterization of a membrane-associated lipoxygenase in tomato fruit. <i>Plant Physiology</i> , 1992 , 100, 1802-7	6.6	48
74	Role of microRNAs involved in plant response to nitrogen and phosphorous limiting conditions. <i>Frontiers in Plant Science</i> , 2015 , 6, 629	6.2	47
73	A 330 bp region of the spinach nitrite reductase gene promoter directs nitrate-inducible tissue-specific expression in transgenic tobacco. <i>Plant Journal</i> , 1993 , 4, 317-326	6.9	47
72	Transient Accumulation of Nitrite Reductase mRNA in Maize following the Addition of Nitrate. <i>Plant Physiology</i> , 1989 , 90, 1214-20	6.6	47
71	Binding of pea cytochrome f to the inner membrane of Escherichia coli requires the bacterial secA gene product. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1985 , 82, 7955-9	11.5	47
70	AtMBD9 modulates Arabidopsis development through the dual epigenetic pathways of DNA methylation and histone acetylation. <i>Plant Journal</i> , 2009 , 59, 123-35	6.9	44
69	Physiological and genetic analysis of Arabidopsis thaliana anthocyanin biosynthesis mutants under chronic adverse environmental conditions. <i>Journal of Experimental Botany</i> , 2013 , 64, 229-40	7	43
68	Hydrophobic-domain-dependent protein-protein interactions mediate the localization of GPAT enzymes to ER subdomains. <i>Traffic</i> , 2011 , 12, 452-72	5.7	43
67	Biosynthesis of cannflavins A and B from Cannabis sativa L. <i>Phytochemistry</i> , 2019 , 164, 162-171	4	42
66	ANAC032 Positively Regulates Age-Dependent and Stress-Induced Senescence in Arabidopsis thaliana. <i>Plant and Cell Physiology</i> , 2016 , 57, 2029-2046	4.9	42
65	Co-expression of both the maize large and wheat small subunit genes of ribulose-bisphosphate carboxylase in Escherichia coli. <i>FEBS Journal</i> , 1987 , 168, 227-31		41
64	Synthesis and secretion of wheat alpha-amylase in Saccharomyces cerevisiae. <i>Gene</i> , 1987 , 55, 353-6	3.8	38
63	The S locus glycoprotein and the S receptor kinase are sufficient for self-pollen rejection in Brassica. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000 , 97, 371.	3- 7 1.5	38
62	Use of the polymerase chain reaction to isolate an S-locus glycoprotein cDNA introgressed from Brassica campestris into B. napus ssp. oleifera. <i>Molecular Genetics and Genomics</i> , 1992 , 234, 185-92		37
61	Translational coupling of the maize chloroplast atpB and atpE genes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989 , 86, 4066-70	11.5	36

(2015-2016)

60	Ammonium-induced architectural and anatomical changes with altered suberin and lignin levels significantly change water and solute permeabilities of rice (Oryza sativa L.) roots. <i>Planta</i> , 2016 , 243, 231-49	4.7	35
59	Nitrate effects on nitrate reductase activity and nitrite reductase mRNA levels in maize suspension cultures. <i>Plant Physiology</i> , 1989 , 90, 962-7	6.6	33
58	Alteration of the bZIP60/IRE1 pathway affects plant response to ER stress in Arabidopsis thaliana. <i>PLoS ONE</i> , 2012 , 7, e39023	3.7	33
57	ROS-Induced anthocyanin production provides feedback protection by scavenging ROS and maintaining photosynthetic capacity in Arabidopsis. <i>Plant Signaling and Behavior</i> , 2018 , 13, e1451708	2.5	32
56	Regulation by light and metabolites of ferredoxin-dependent glutamate synthase in maize. <i>Physiologia Plantarum</i> , 2001 , 112, 524-530	4.6	30
55	Metabolic and co-expression network-based analyses associated with nitrate response in rice. <i>BMC Genomics</i> , 2014 , 15, 1056	4.5	29
54	Global DNA methylation analysis using methyl-sensitive amplification polymorphism (MSAP). <i>Methods in Molecular Biology</i> , 2014 , 1062, 285-98	1.4	29
53	Molecular mechanisms of self-recognition in Brassica self-incompatibility. <i>Trends in Plant Science</i> , 2000 , 5, 432-8	13.1	29
52	Overexpression of OsGATA12 regulates chlorophyll content, delays plant senescence and improves rice yield under high density planting. <i>Plant Molecular Biology</i> , 2017 , 94, 215-227	4.6	28
51	Transcript and metabolite signature of maize source leaves suggests a link between transitory starch to sucrose balance and the autonomous floral transition. <i>Journal of Experimental Botany</i> , 2012 , 63, 5079-92	7	28
50	Identification of an S-locus glycoprotein allele introgressed from B. napus ssp. rapifera to B. napus ssp. oleifera. <i>Plant Journal</i> , 1992 , 2, 983-989	6.9	28
49	Regulation of maize root nitrate reductase mRNA levels. <i>Physiologia Plantarum</i> , 1992 , 85, 561-566	4.6	28
48	Overexpression of miR169o, an Overlapping MicroRNA in Response to Both Nitrogen Limitation and Bacterial Infection, Promotes Nitrogen Use Efficiency and Susceptibility to Bacterial Blight in Rice. <i>Plant and Cell Physiology</i> , 2018 , 59, 1234-1247	4.9	27
47	High throughput RNA sequencing of a hybrid maize and its parents shows different mechanisms responsive to nitrogen limitation. <i>BMC Genomics</i> , 2014 , 15, 77	4.5	27
46	Characterization and immunolocalization of a cytosolic calcium-binding protein from Brassica napus and Arabidopsis pollen. <i>Plant Physiology</i> , 1999 , 120, 787-98	6.6	27
45	Regulation of seed dormancy and germination by nitrate. <i>Seed Science Research</i> , 2018 , 28, 150-157	1.3	26
44	Auxin-responsive SAUR39 gene modulates auxin level in rice. <i>Plant Signaling and Behavior</i> , 2009 , 4, 1174	l-25 5	26
43	Overexpression of the CC-type glutaredoxin, OsGRX6 affects hormone and nitrogen status in rice plants. <i>Frontiers in Plant Science</i> , 2015 , 6, 934	6.2	25

42	Quantitative and qualitative differences in C6-volatile production from the lipoxygenase pathway in an alcohol dehydrogenase mutant of Arabidopsis thaliana. <i>Physiologia Plantarum</i> , 1998 , 104, 97-104	4.6	25
41	A complete cDNA for adenine phosphoribosyltransferase from Arabidopsis thaliana. <i>Plant Molecular Biology</i> , 1992 , 18, 653-62	4.6	25
40	Functional characterization of the rice UDP-glucose 4-epimerase 1, OsUGE1: a potential role in cell wall carbohydrate partitioning during limiting nitrogen conditions. <i>PLoS ONE</i> , 2014 , 9, e96158	3.7	24
39	Peroxidase-Induced Wilting in Transgenic Tobacco Plants. <i>Plant Cell</i> , 1990 , 2, 7	11.6	23
38	Active Translation of the D-1 Protein of Photosystem II in Senescing Leaves. <i>Plant Physiology</i> , 1992 , 99, 589-94	6.6	21
37	Agrobacterium-derived cytokinin influences plastid morphology and starch accumulation in Nicotiana benthamiana during transient assays. <i>BMC Plant Biology</i> , 2014 , 14, 127	5.3	20
36	Developmental regulation of two tomato lipoxygenase promoters in transgenic tobacco and tomato. <i>Plant Molecular Biology</i> , 1997 , 33, 835-46	4.6	19
35	The challenges of commercializing second-generation transgenic crop traits necessitate the development of international public sector research infrastructure. <i>Journal of Experimental Botany</i> , 2014 , 65, 5673-82	7	17
34	Loss of callose in the stigma papillae does not affect the Brassica self-incompatibility phenotype. <i>Planta</i> , 1997 , 203, 327-331	4.7	17
33	Transformation of Arabidopsis with a Brassica SLG/SRK region and ARC1 gene is not sufficient to transfer the self-incompatibility phenotype. <i>Molecular Genetics and Genomics</i> , 2000 , 263, 648-54		17
32	Synthesis of a wheat storage protein subunit in Escherichia coli using novel expression vectors. <i>Gene</i> , 1985 , 35, 159-67	3.8	17
31	An S Receptor Kinase Gene in Self-Compatible Brassica napus Has a 1-bp Deletion. <i>Plant Cell</i> , 1993 , 5, 531	11.6	16
30	Nitrogen effects on the induction of ferredoxin-dependent glutamate synthase and its mRNA in maize leaves under the light. <i>Plant Science</i> , 1996 , 114, 83-91	5.3	15
29	Features of the extracellular domain of the S-locus receptor kinase from Brassica. <i>Molecular Genetics and Genomics</i> , 1994 , 244, 630-7		15
28	Overexpression of ANAC046 Promotes Suberin Biosynthesis in Roots of. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	15
27	The SNAC-A Transcription Factor ANAC032 Reprograms Metabolism in Arabidopsis. <i>Plant and Cell Physiology</i> , 2019 , 60, 999-1010	4.9	14
26	Analysis of cis-acting DNA elements mediating induction and repression of the spinach nitrite reductase gene. <i>Planta</i> , 1998 , 206, 66-71	4.7	14
25	Nitrogen transporter and assimilation genes exhibit developmental stage-selective expression in maize (Zea mays L.) associated with distinct cis-acting promoter motifs. <i>Plant Signaling and Behavior</i> , 2013, 8,	2.5	13

24	Expression of a wheat alpha-amylase gene in Escherichia coli: recognition of the translational initiation site and the signal peptide. <i>Gene</i> , 1986 , 45, 11-8	3.8	12
23	Nitrogen limitation and high density responses in rice suggest a role for ethylene under high density stress. <i>BMC Genomics</i> , 2014 , 15, 681	4.5	10
22	Exploring the molecular and metabolic factors contributing to the adaptation of maize seedlings to nitrate limitation. <i>Frontiers in Plant Science</i> , 2011 , 2, 49	6.2	9
21	A novel strategy for regulated expression of a cytotoxic gene. <i>Gene</i> , 2001 , 279, 175-9	3.8	9
20	Engineering resistance to trypsin inactivation into L-asparaginase through the production of a chimeric protein between the enzyme and a protective single-chain antibody. <i>Enzyme and Microbial Technology</i> , 1995 , 17, 757-64	3.8	9
19	Synthesis of maize chloroplast ATP-synthase beta-subunit fusion proteins in Escherichia coli and binding to the inner membrane. <i>Gene</i> , 1986 , 41, 241-7	3.8	9
18	A multivariate Poisson-log normal mixture model for clustering transcriptome sequencing data. <i>BMC Bioinformatics</i> , 2019 , 20, 394	3.6	8
17	Genome-wide binding analysis of AtGNC and AtCGA1 demonstrates their cross-regulation and common and specific functions. <i>Plant Direct</i> , 2017 , 1, e00016	3.3	7
16	Identification and characterization of a putative light-harvesting chlorophyll a/b-binding protein gene encoded at a fertility restorer locus for the Ogura CMS in Brassica napus L <i>Theoretical and Applied Genetics</i> , 2001 , 102, 759-766	6	7
15	A binary vector-based large insert library for Brassica napus and identification of clones linked to a fertility restorer locus for Ogura cytoplasmic male sterility (CMS). <i>Genome</i> , 2000 , 43, 102-109	2.4	7
14	Altered Expression of OsNLA1 Modulates Pi Accumulation in Rice (L.) Plants. <i>Frontiers in Plant Science</i> , 2017 , 8, 928	6.2	6
13	Using bacteria to analyze sequences involved in chloroplast gene expression. <i>Photosynthesis Research</i> , 1988 , 19, 7-22	3.7	6
12	S-Locus Receptor Kinase Genes and Self-incompatibility in Brassica napus. <i>Plant Gene Research</i> , 1996 , 217-230		4
11	Identification of regulatory genes to improve nitrogen use efficiency. <i>Canadian Journal of Plant Science</i> , 2014 , 94, 1009-1012	1	3
10	Evidence that the Arabidopsis Ubiquitin C-terminal Hydrolases 1 and 2 associate with the 26S proteasome and the TREX-2 complex. <i>Plant Signaling and Behavior</i> , 2012 , 7, 1415-9	2.5	3
9	The AtPP gene of the Brassica napus S locus region is specifically expressed in the stigma and encodes a protein similar to a methyltransferase involved in plant defense. <i>Sexual Plant Reproduction</i> , 2001 , 13, 309-314		3
8	Regulation of maize root nitrate reductase mRNA levels. <i>Physiologia Plantarum</i> , 1992 , 85, 561-566	4.6	2
7	Antigenic similarity between the Bubunits of the ATPases of a bacterium, a yeast and a higher plant. <i>Phytochemistry</i> , 1985 , 24, 259-260	4	1

6	Distinct domains within the NITROGEN LIMITATION ADAPTATION protein mediate its subcellular localization and function in the nitrate-dependent phosphate homeostasis pathway. <i>Botany</i> , 2018 , 96, 79-96	1.3 1
5	Molecular Genetics of Self-incompatibility in Brassica napus. <i>Annals of Botany</i> , 2000 , 85, 133-139	4.1
4	The Self-Incompatibility Phenotype in Brassica Is Altered by the Transformation of a Mutant S Locus Receptor Kinase. <i>Plant Cell</i> , 1998 , 10, 209	11.6
3	Structural and Transcriptional Comparative Analysis of the S Locus Regions in Two Self-Incompatible Brassica napus Lines. <i>Plant Cell</i> , 1999 , 11, 2217	11.6
3		11.6