

Andreia M Ionescu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7537419/publications.pdf>

Version: 2024-02-01

9
papers

255
citations

1478505

6
h-index

1588992

8
g-index

9
all docs

9
docs citations

9
times ranked

426
citing authors

#	ARTICLE	IF	CITATIONS
1	FoxA Family Members Are Crucial Regulators of the Hypertrophic Chondrocyte Differentiation Program. <i>Developmental Cell</i> , 2012, 22, 927-939.	7.0	70
2	GATA6 Is a Crucial Regulator of Shh in the Limb Bud. <i>PLoS Genetics</i> , 2014, 10, e1004072.	3.5	48
3	At the Crossroads of the Adipocyte and Osteoclast Differentiation Programs: Future Therapeutic Perspectives. <i>International Journal of Molecular Sciences</i> , 2020, 21, 2277.	4.1	48
4	The Role of <i>Bmp2</i> in the Maturation and Maintenance of the Murine Knee Joint. <i>Journal of Bone and Mineral Research</i> , 2018, 33, 1708-1717.	2.8	34
5	A FoxA2+ long-term stem cell population is necessary for growth plate cartilage regeneration after injury. <i>Nature Communications</i> , 2022, 13, 2515.	12.8	22
6	BMP-mediated induction of GATA4/5/6 blocks somitic responsiveness to SHH. <i>Development (Cambridge)</i> , 2014, 141, 3978-3987.	2.5	21
7	The secreted tyrosine kinase VLK is essential for normal platelet activation and thrombus formation. <i>Blood</i> , 2022, 139, 104-117.	1.4	6
8	Overexpression of transcription factor FoxA2 in the developing skeleton causes an enlargement of the cartilage hypertrophic zone, but it does not trigger ectopic differentiation in immature chondrocytes. <i>Bone</i> , 2022, 160, 116418.	2.9	6
9	The Secreted Tyrosine Kinase Vlk Is Essential for Normal Platelet Activation and Thrombus Formation. <i>Blood</i> , 2020, 136, 10-11.	1.4	0