Beata Guzow-Krzemińska

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7566708/publications.pdf

Version: 2024-02-01

687220 642610 32 571 13 23 g-index citations h-index papers 34 34 34 700 docs citations times ranked citing authors all docs

#	Article	IF	Citations
1	The Isoxazole Derivative of Usnic Acid Induces an ER Stress Response in Breast Cancer Cells That Leads to Paraptosis-like Cell Death. International Journal of Molecular Sciences, 2022, 23, 1802.	1.8	14
2	Phylogeny and Ecology of Trebouxia Photobionts From Bolivian Lichens. Frontiers in Microbiology, 2022, 13, 779784.	1.5	5
3	New lineages of photobionts in Bolivian lichens expand our knowledge on habitat preferences and distribution of Asterochloris algae. Scientific Reports, 2021, 11, 8701.	1.6	15
4	A molecular re-evaluation of <i>Parmelia encryptata</i> with notes on its distribution. Lichenologist, 2021, 53, 341-345.	0.5	1
5	Infraspecific variation of some brown Parmeliae (in Poland) – a comparison of ITS rDNA and non-molecular characters. MycoKeys, 2021, 85, 127-160.	0.8	4
6	Trentepohlialean Algae (Trentepohliales, Ulvophyceae) Show Preference to Selected Mycobiont Lineages in Lichen Symbioses. Journal of Phycology, 2020, 56, 979-993.	1.0	16
7	Two new Micarea species (Pilocarpaceae) from Western Europe. Plant and Fungal Systematics, 2020, 65, 189-199.	0.7	5
8	One Name – One Fungus: The Influence of Photosynthetic Partners on the Taxonomy and Systematics of Lichenized Fungi. Acta Societatis Botanicorum Poloniae, 2020, 89, .	0.8	3
9	Synthesis of Usnic Acid Derivatives and Evaluation of Their Antiproliferative Activity against Cancer Cells. Journal of Natural Products, 2019, 82, 1768-1778.	1.5	27
10	New species and records of lichens from Bolivia. Phytotaxa, 2019, 397, 257.	0.1	14
11	Phylogenetic placement of Lepraria cryptovouauxii sp. nov. (Lecanorales, Lecanoromycetes,) Tj ETQq1 1 0.78431	.4 rgBT /O	veglock 10 Tf
12	Understanding the evolution of phenotypical characters in the Micarea prasina group (Pilocarpaceae) and descriptions of six new species within the group. MycoKeys, 2019, 57, 1-30.	0.8	14
13	Morphology and secondary chemistry in species recognition of Parmelia omphalodes group – evidence from molecular data with notes on the ecological niche modelling and genetic variability of photobionts. MycoKeys, 2019, 61, 39-74.	0.8	6
14	<i>Bacidina mendax</i> sp. nov., a new widespread species in Central Europe, together with a new combination within the genus <i>Bacidina</i> . Lichenologist, 2018, 50, 43-57.	0.5	15
15	Phylogenetic approaches reveal a new sterile lichen in the genus Loxospora (Sarrameanales,) Tj ETQq $1\ 1\ 0.7843$	14 rgBT /C	verlock 10 <mark>Tf</mark>
16	Photobiont switching causes changes in the reproduction strategy and phenotypic dimorphism in the Arthoniomycetes. Scientific Reports, 2018, 8, 4952.	1.6	41
17	Evaluation of diagnostic chemical and morphological characters in five Parmelia species (Parmeliaceae, lichenized Ascomycota) with special emphasis on the thallus pruinosity. Phytotaxa, 2018, 383, 165.	0.1	11
18	Antibacterial and anticancer activities of acetone extracts from in vitro cultured lichen-forming fungi. BMC Complementary and Alternative Medicine, 2017, 17, 300.	3.7	22

#	Article	IF	CITATIONS
19	Lecanora stanislai, a new, sterile, usnic acid containing lichen species from Eurasia and North America. Phytotaxa, 2017, 329, 201.	0.1	18
20	<i>Micarea soralifera</i> sp. nov., a new sorediate species in the <i>M. prasina</i> group. Lichenologist, 2016, 48, 161-169.	0.5	30
21	Lichens and lichenicolous fungi of Magurski National Park (Poland, Western Carpathians). Polish Botanical Journal, 2016, 61, 127-160.	0.5	5
22	Phylogenetic relationship of the stringent response-related genes of marine bacteria. Acta Biochimica Polonica, 2015, 62, 773-783.	0.3	2
23	A rapidly progressing, deadly disease of Actias selene (Indian moon moth) larvae associated with a mixed bacterial and baculoviral infection. Journal of Biosciences, 2015, 40, 487-495.	0.5	3
24	The Lichen Order Peltigerales in Bolivia â€" The First Assessment of the Biodiversity. Herzogia, 2014, 27, 321-345.	0.1	3
25	Antibacterial activity of lichen secondary metabolite usnic acid is primarily caused by inhibition of RNA and DNA synthesis. FEMS Microbiology Letters, 2014, 353, 57-62.	0.7	71
26	<i>In vitro</i> culturing and resynthesis of the mycobiont <i>Protoparmeliopsis muralis</i> with algal bionts. Lichenologist, 2013, 45, 65-76.	0.5	17
27	Development of microsatellite markers in Protoparmeliopsis muralis (lichenized Ascomycete) – a common lichen species. Lichenologist, 2013, 45, 791-798.	0.5	7
28	ITS rDNA data confirm a delimitation of <i>Bacidina arnoldiana</i> and <i>B. sulphurella</i> and support a description of a new species within the genus <i>Bacidina</i> . Lichenologist, 2012, 44, 743-755.	0.5	14
29	A new Agonimia from Europe with a flabelliform thallus. Lichenologist, 2012, 44, 55-66.	0.5	23
30	A phylogenetic study of the <i>Micarea prasina </i> proup shows that <i>Micarea micrococca </i> includes three distinct lineages. Lichenologist, 2010, 42, 7-21.	0.5	43
31	Photobiont flexibility in the lichen Protoparmeliopsis muralis as revealed by ITS rDNA analyses. Lichenologist, 2006, 38, 469-476.	0.5	75
32	A preliminary study on the phylogeny of the genus Melanelia using nuclear large subunit ribosomal DNA sequences. Lichenologist, 2003, 35, 83-86.	0.5	9