

Zhen-Yu Du

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/756749/publications.pdf>

Version: 2024-02-01

136
papers

5,134
citations

87723

38
h-index

114278

63
g-index

138
all docs

138
docs citations

138
times ranked

4553
citing authors

#	ARTICLE	IF	CITATIONS
1	<i>Bacillus amyloliquefaciens</i> ameliorates high-carbohydrate diet-induced metabolic phenotypes by restoration of intestinal acetate-producing bacteria in Nile Tilapia. <i>British Journal of Nutrition</i> , 2022, 127, 653-665.	1.2	30
2	Are we actually measuring growth? An appeal to use a more comprehensive growth index system for advancing aquaculture research. <i>Reviews in Aquaculture</i> , 2022, 14, 525-527.	4.6	16
3	The lipids. , 2022, , 303-467.		18
4	More simple more worse: Simple carbohydrate diets cause alterations in glucose and lipid metabolism in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Aquaculture</i> , 2022, 550, 737857.	1.7	13
5	Dietary sodium lactate promotes protein and lipid deposition through increasing energy supply from glycolysis in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Aquaculture</i> , 2022, 550, 737858.	1.7	9
6	Mildronate triggers growth suppression and lipid accumulation in largemouth bass (<i>Micropterus</i>) Tj ETQq0 0 0 rgBT/Overlock 10 Tf 50 5	0.9	4
7	Dietary L-carnitine supplementation recovers the increased pH and hardness in fillets caused by high-fat diet in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Food Chemistry</i> , 2022, 382, 132367.	4.2	18
8	Peroxisome proliferator-activated receptor gamma is essential for stress adaptation by maintaining lipid homeostasis in female fish. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2022, 1867, 159162.	1.2	1
9	Dietary L-Carnitine Alleviates the Adverse Effects Caused by Reducing Protein and Increasing Fat Contents in Diet Juvenile Largemouth Bass (<i>Micropterus salmoides</i>). <i>Aquaculture Nutrition</i> , 2022, 2022, 1-14.	1.1	6
10	Intestinal Microbiota Mediates Gossypol-Induced Intestinal Inflammation, Oxidative Stress, and Apoptosis in Fish. <i>Journal of Agricultural and Food Chemistry</i> , 2022, 70, 6688-6697.	2.4	19
11	Microbiota derived butyrate affected the muscle texture of Nile tilapia (<i>Oreochromis niloticus</i>) fed with different protein sources. <i>Food Chemistry</i> , 2022, 393, 133392.	4.2	9
12	Inhibition of pyruvate dehydrogenase kinase improves carbohydrate utilization in Nile tilapia by regulating PDK2/4-PDHE1± axis and insulin sensitivity. <i>Animal Nutrition</i> , 2022, 11, 25-37.	2.1	6
13	<i>Lactobacillus plantarum</i> Ameliorates High-Carbohydrate Diet-Induced Hepatic Lipid Accumulation and Oxidative Stress by Upregulating Uridine Synthesis. <i>Antioxidants</i> , 2022, 11, 1238.	2.2	8
14	<i>Bacillus amyloliquefaciens</i> protects Nile tilapia against <i>Aeromonas hydrophila</i> infection and alleviates liver inflammation induced by high-carbohydrate diet. <i>Fish and Shellfish Immunology</i> , 2022, 127, 836-842.	1.6	5
15	Trehalose alleviated hepatic cholesterol accumulation via inhibiting transformation from glucose-derived acyl-CoA to cholesterol synthesis in Nile tilapia. <i>Aquaculture</i> , 2022, 560, 738600.	1.7	1
16	A global analysis on the systemic effects of antibiotics in cultured fish and their potential human health risk: a review. <i>Reviews in Aquaculture</i> , 2021, 13, 1015-1059.	4.6	105
17	Inulin alleviates adverse metabolic syndrome and regulates intestinal microbiota composition in Nile tilapia (<i>Oreochromis niloticus</i>) fed with high-carbohydrate diet. <i>British Journal of Nutrition</i> , 2021, 126, 161-171.	1.2	26
18	The reduction of lipid-sourced energy production caused by ATGL inhibition cannot be compensated by activation of HSL, autophagy, and utilization of other nutrients in fish. <i>Fish Physiology and Biochemistry</i> , 2021, 47, 173-188.	0.9	8

#	ARTICLE	IF	CITATIONS
19	Dietary tea polyphenols change flesh quality with dose-related manner in the GIFT tilapia fed with a high-fat diet. <i>Aquaculture Nutrition</i> , 2021, 27, 519-532.	1.1	17
20	Dietary aflatoxin impairs flesh quality through reducing nutritional value and changing myofiber characteristics in yellow catfish (<i>Pelteobagrus fulvidraco</i>). <i>Animal Feed Science and Technology</i> , 2021, 274, 114764.	1.1	11
21	The individual and combined effects of hypoxia and high-fat diet feeding on nutrient composition and flesh quality in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Food Chemistry</i> , 2021, 343, 128479.	4.2	35
22	High protein intake promotes the adaptation to chronic hypoxia in zebrafish (<i>Danio rerio</i>). <i>Aquaculture</i> , 2021, 535, 736356.	1.7	7
23	Reduced fatty acid β -oxidation improves glucose catabolism and liver health in Nile tilapia (<i>Oreochromis niloticus</i>) juveniles fed a high-starch diet. <i>Aquaculture</i> , 2021, 535, 736392.	1.7	19
24	Different effects of two dietary levels of tea polyphenols on the lipid deposition, immunity and antioxidant capacity of juvenile GIFT tilapia (<i>Oreochromis niloticus</i>) fed a high-fat diet. <i>Aquaculture</i> , 2021, 542, 736896.	1.7	28
25	Lipolysis and lipophagy play individual and interactive roles in regulating triacylglycerol and cholesterol homeostasis and mitochondrial form in zebrafish. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2021, 1866, 158988.	1.2	12
26	Alteration and the Function of Intestinal Microbiota in High-Fat-Diet- or Genetics-Induced Lipid Accumulation. <i>Frontiers in Microbiology</i> , 2021, 12, 741616.	1.5	4
27	Oligosaccharides improve the flesh quality and nutrition value of Nile tilapia fed with high carbohydrate diet. <i>Food Chemistry Molecular Sciences</i> , 2021, 3, 100040.	0.9	9
28	Effects of replacing soybean meal protein with cottonseed protein concentrate on the growth condition and intestinal health of Nile tilapia (<i>Oreochromis niloticus</i>). <i>Aquaculture Nutrition</i> , 2021, 27, 2436-2447.	1.1	13
29	Activation of peroxisome proliferator-activated receptor α stimulated lipid catabolism in liver of largemouth bass, <i>Micropterus salmoides</i> . <i>Aquaculture Nutrition</i> , 2021, 27, 2749-2758.	1.1	3
30	<i>Pediococcus pentosaceus</i> Enhances Host Resistance Against Pathogen by Increasing IL-1 β Production: Understanding Probiotic Effectiveness and Administration Duration. <i>Frontiers in Immunology</i> , 2021, 12, 766401.	2.2	9
31	Inhibition of intestinal lipases alleviates the adverse effects caused by high-fat diet in Nile tilapia. <i>Fish Physiology and Biochemistry</i> , 2020, 46, 111-123.	0.9	15
32	Metabolism of linoleic and linolenic acids in hepatocytes of two freshwater fish with different n-3 or n-6 fatty acid requirements. <i>Aquaculture</i> , 2020, 515, 734595.	1.7	20
33	Sodium acetate alleviated high-carbohydrate induced intestinal inflammation by suppressing MAPK and NF- κ B signaling pathways in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Fish and Shellfish Immunology</i> , 2020, 98, 758-765.	1.6	56
34	High carbohydrate diet partially protects Nile tilapia (<i>Oreochromis niloticus</i>) from oxytetracycline-induced side effects. <i>Environmental Pollution</i> , 2020, 256, 113508.	3.7	37
35	Inhibited carnitine synthesis impairs adaptation to high-fat diet in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Aquaculture Reports</i> , 2020, 16, 100249.	0.7	8
36	High-carbohydrate diet promotes the adaptation to acute hypoxia in zebrafish. <i>Fish Physiology and Biochemistry</i> , 2020, 46, 665-679.	0.9	17

#	ARTICLE	IF	CITATIONS
37	Gnotobiotic models: Powerful tools for deeply understanding intestinal microbiota-host interactions in aquaculture. <i>Aquaculture</i> , 2020, 517, 734800.	1.7	29
38	Peroxisomal proliferator-activated receptor β deficiency induces the reprogramming of nutrient metabolism in zebrafish. <i>Journal of Physiology</i> , 2020, 598, 4537-4553.	1.3	20
39	Gemfibrozil improves lipid metabolism in Nile tilapia <i>Oreochromis niloticus</i> fed a high-carbohydrate diet through peroxisome proliferator activated receptor- β activation. <i>General and Comparative Endocrinology</i> , 2020, 296, 113537.	0.8	24
40	<i>Citrobacter</i> Species Increase Energy Harvest by Modulating Intestinal Microbiota in Fish: Nondominant Species Play Important Functions. <i>MSystems</i> , 2020, 5, .	1.7	27
41	Impaired peroxisomal fat oxidation induces hepatic lipid accumulation and oxidative damage in Nile tilapia. <i>Fish Physiology and Biochemistry</i> , 2020, 46, 1229-1242.	0.9	15
42	Mitochondrial Fatty Acid β -Oxidation Inhibition Promotes Glucose Utilization and Protein Deposition through Energy Homeostasis Remodeling in Fish. <i>Journal of Nutrition</i> , 2020, 150, 2322-2335.	1.3	44
43	Dietary L-carnitine improves glycogen and protein accumulation in Nile tilapia via increasing lipid-sourced energy supply: An isotope-based metabolic tracking. <i>Aquaculture Reports</i> , 2020, 17, 100302.	0.7	12
44	The regulation of rapamycin on nutrient metabolism in Nile tilapia fed with high-energy diet. <i>Aquaculture</i> , 2020, 520, 734975.	1.7	22
45	Environmental estrogen exposure converts lipid metabolism in male fish to a female pattern mediated by AMPK and mTOR signaling pathways. <i>Journal of Hazardous Materials</i> , 2020, 394, 122537.	6.5	41
46	Functional differences between L- and D-carnitine in metabolic regulation evaluated using a low-carnitine Nile tilapia model. <i>British Journal of Nutrition</i> , 2019, 122, 625-638.	1.2	20
47	Inhibited Lipophagy Suppresses Lipid Metabolism in Zebrafish Liver Cells. <i>Frontiers in Physiology</i> , 2019, 10, 1077.	1.3	22
48	Concentration-dependent effects of 17 β -estradiol and bisphenol A on lipid deposition, inflammation and antioxidant response in male zebrafish (<i>Danio rerio</i>). <i>Chemosphere</i> , 2019, 237, 124422.	4.2	40
49	Inhibited autophagy impairs systemic nutrient metabolism in Nile tilapia. <i>Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology</i> , 2019, 236, 110521.	0.8	14
50	The Responses of Germ-Free Zebrafish (<i>Danio rerio</i>) to Varying Bacterial Concentrations, Colonization Time Points, and Exposure Duration. <i>Frontiers in Microbiology</i> , 2019, 10, 2156.	1.5	24
51	PPAR β activation enhances the ability of Nile tilapia (<i>Oreochromis niloticus</i>) to resist <i>Aeromonas hydrophila</i> infection. <i>Fish and Shellfish Immunology</i> , 2019, 94, 675-684.	1.6	16
52	Sex-specific alterations of lipid metabolism in zebrafish exposed to polychlorinated biphenyls. <i>Chemosphere</i> , 2019, 221, 768-777.	4.2	44
53	Reduced oxidative stress increases acute cold stress tolerance in zebrafish. <i>Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology</i> , 2019, 235, 166-173.	0.8	26
54	Diacylglycerol oil reduces fat accumulation and increases protein content by inducing lipid catabolism and protein metabolism in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Aquaculture</i> , 2019, 510, 90-99.	1.7	11

#	ARTICLE	IF	CITATIONS
55	High fat diet worsens the adverse effects of antibiotic on intestinal health in juvenile Nile tilapia (<i>Oreochromis niloticus</i>). <i>Science of the Total Environment</i> , 2019, 680, 169-180.	3.9	61
56	CIDEA and CIDEC are regulated by CREB and are not induced during fasting in grass carp <i>Ctenopharyngodon idella</i> adipocytes. <i>Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology</i> , 2019, 234, 50-57.	0.7	1
57	Comparison of effects of dietary-specific fatty acids on growth and lipid metabolism in Nile tilapia. <i>Aquaculture Nutrition</i> , 2019, 25, 862-872.	1.1	11
58	The comparisons in protective mechanisms and efficiencies among dietary α -lipoic acid, β -glucan and l-carnitine on Nile tilapia infected by <i>Aeromonas hydrophila</i> . <i>Fish and Shellfish Immunology</i> , 2019, 86, 785-793.	1.6	46
59	Forskolin reduces fat accumulation in Nile tilapia (<i>Oreochromis niloticus</i>) through stimulating lipolysis and beta-oxidation. <i>Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology</i> , 2019, 230, 7-15.	0.8	22
60	Fasting enhances cold resistance in fish through stimulating lipid catabolism and autophagy. <i>Journal of Physiology</i> , 2019, 597, 1585-1603.	1.3	96
61	Dietary oils modify lipid molecules and nutritional value of fillet in Nile tilapia: A deep lipidomics analysis. <i>Food Chemistry</i> , 2019, 277, 515-523.	4.2	50
62	The metabolic regulation of dietary L-carnitine in aquaculture nutrition: present status and future research strategies. <i>Reviews in Aquaculture</i> , 2019, 11, 1228-1257.	4.6	47
63	IGF-1 induces SOCS-2 but not SOCS-1 and SOCS-3 transcription in juvenile Nile tilapia (<i>Oreochromis niloticus</i>). <i>Journal of Endocrinology</i> , 2019, 202, 1-10.	1.0	14
64	Influence of Long-Term Feeding Antibiotics on the Gut Health of Zebrafish. <i>Zebrafish</i> , 2018, 15, 340-348.	0.5	65
65	Environmental concentrations of antibiotics impair zebrafish gut health. <i>Environmental Pollution</i> , 2018, 235, 245-254.	3.7	198
66	Chronic exposure to low environmental concentrations and legal aquaculture doses of antibiotics cause systemic adverse effects in Nile tilapia and provoke differential human health risk. <i>Environment International</i> , 2018, 115, 205-219.	4.8	241
67	G0S2a1 (G0/G1 switch gene 2a1) is downregulated by TNF- α in grass carp (<i>Ctenopharyngodon idellus</i>) hepatocytes through PPAR α inhibition. <i>Gene</i> , 2018, 641, 1-7.	1.0	11
68	Soybean and cottonseed meals are good candidates for fishmeal replacement in the diet of juvenile <i>Macrobrachium nipponense</i> . <i>Aquaculture International</i> , 2018, 26, 309-324.	1.1	24
69	Leptin Selectively Regulates Nutrients Metabolism in Nile Tilapia Fed on High Carbohydrate or High Fat Diet. <i>Frontiers in Endocrinology</i> , 2018, 9, 574.	1.5	36
70	The protein-sparing effect of α -lipoic acid in juvenile grass carp, <i>Ctenopharyngodon idellus</i> : effects on lipolysis, fatty acid β -oxidation and protein synthesis. <i>British Journal of Nutrition</i> , 2018, 120, 977-987.	1.2	40
71	Tracking pollutants in dietary fish oil: From ocean to table. <i>Environmental Pollution</i> , 2018, 240, 733-744.	3.7	21
72	Untargeted GC-MS metabolomics reveals metabolic differences in the Chinese mitten-crab (<i>Eriocheir sinensis</i>) fed with dietary palm oil or olive oil. <i>Aquaculture Nutrition</i> , 2018, 24, 1623-1637.	1.1	12

#	ARTICLE	IF	CITATIONS
73	Different physiological roles of insulin receptors in mediating nutrient metabolism in zebrafish. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2018, 315, E38-E51.	1.8	36
74	Inhibited Carnitine Synthesis Causes Systemic Alteration of Nutrient Metabolism in Zebrafish. <i>Frontiers in Physiology</i> , 2018, 9, 509.	1.3	17
75	The Presence or Absence of Intestinal Microbiota Affects Lipid Deposition and Related Genes Expression in Zebrafish (<i>Danio rerio</i>). <i>Frontiers in Microbiology</i> , 2018, 9, 1124.	1.5	63
76	Dietary microencapsulated oil improves immune function and intestinal health in Nile tilapia fed with high-fat diet. <i>Aquaculture</i> , 2018, 496, 19-29.	1.7	61
77	Lipophagy is essential for lipid metabolism in fish. <i>Science Bulletin</i> , 2018, 63, 879-882.	4.3	29
78	Pigment epithelium-derived factor improves TNF α -induced hepatic steatosis in grass carp (<i>Ctenopharyngodon idella</i>). <i>Developmental and Comparative Immunology</i> , 2017, 71, 8-17.	1.0	11
79	Nutritional background changes the hypolipidemic effects of fenofibrate in Nile tilapia (<i>Oreochromis</i>) Tj ETQq1 1 0.784314 r _{BT} /Ove	1.6	35
80	Systemic regulation of L-carnitine in nutritional metabolism in zebrafish, <i>Danio rerio</i> . <i>Scientific Reports</i> , 2017, 7, 40815.	1.6	53
81	Forkhead box O1 in grass carp <i>Ctenopharyngodon idella</i> : Molecular characterization, gene structure, tissue distribution and mRNA expression in insulin-inhibited adipocyte lipolysis. <i>Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology</i> , 2017, 204, 76-84.	0.8	14
82	α -lipoic acid ameliorates n-3 highly-unsaturated fatty acids induced lipid peroxidation via regulating antioxidant defenses in grass carp (<i>Ctenopharyngodon idellus</i>). <i>Fish and Shellfish Immunology</i> , 2017, 67, 359-367.	1.6	37
83	Physiological and metabolic differences between visceral and subcutaneous adipose tissues in Nile tilapia (<i>Oreochromis niloticus</i>). <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2017, 313, R608-R619.	0.9	18
84	Inhibited fatty acid β -oxidation impairs stress resistance ability in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Fish and Shellfish Immunology</i> , 2017, 68, 500-508.	1.6	37
85	The metabolomics responses of Chinese mitten-hand crab (<i>Eriocheir sinensis</i>) to different dietary oils. <i>Aquaculture</i> , 2017, 479, 188-199.	1.7	68
86	Molecular characterization and nutritional regulation of carnitine palmitoyltransferase (CPT) family in grass carp (<i>Ctenopharyngodon idellus</i>). <i>Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology</i> , 2017, 203, 11-19.	0.7	24
87	Influence of Endogenous and Exogenous Estrogenic Endocrine on Intestinal Microbiota in Zebrafish. <i>PLoS ONE</i> , 2016, 11, e0163895.	1.1	49
88	Lipolytic enzymes involving lipolysis in Teleost: Synteny, structure, tissue distribution, and expression in grass carp (<i>Ctenopharyngodon idella</i>). <i>Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology</i> , 2016, 198, 110-118.	0.7	33
89	An LC-MS-based lipidomics approach for studying the impact of dietary eicosapentaenoic acid on modulating methylmercury toxicity in mice. <i>Metabolomics</i> , 2016, 12, 1.	1.4	3
90	Dietary Arachidonic Acid Has a Time-Dependent Differential Impact on Adipogenesis Modulated via COX and LOX Pathways in Grass Carp (<i>Ctenopharyngodon idellus</i>). <i>Lipids</i> , 2016, 51, 1325-1338.	0.7	15

#	ARTICLE	IF	CITATIONS
91	Mechanisms and metabolic regulation of PPAR α activation in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2016, 1861, 1036-1048.	1.2	80
92	Growth, body composition, ammonia tolerance and hepatopancreas histology of white shrimp <i>Litopenaeus vannamei</i> fed diets containing different carbohydrate sources at low salinity. <i>Aquaculture Research</i> , 2016, 47, 1932-1943.	0.9	29
93	Methylmercury Increases and Eicosapentaenoic Acid Decreases the Relative Amounts of Arachidonic Acid-Containing Phospholipids in Mouse Brain. <i>Lipids</i> , 2016, 51, 61-73.	0.7	3
94	Molecular characterization and immune response to lipopolysaccharide (LPS) of the suppressor of cytokine signaling (SOCS)-1, 2 and 3 genes in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Fish and Shellfish Immunology</i> , 2016, 50, 160-167.	1.6	29
95	Systemic adaptation of lipid metabolism in response to low- and high-fat diet in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Physiological Reports</i> , 2015, 3, e12485.	0.7	113
96	Evaluation of the distribution of adipose tissues in fish using magnetic resonance imaging (MRI). <i>Aquaculture</i> , 2015, 448, 112-122.	1.7	38
97	Morphology, mitochondrial development and adipogenic-related genes expression during adipocytes differentiation in grass carp (<i>Ctenopharyngodon idellus</i>). <i>Science Bulletin</i> , 2015, 60, 1241-1251.	4.3	14
98	Nutrients and contaminants in tissues of five fish species obtained from Shanghai markets: Risk-benefit evaluation from human health perspectives. <i>Science of the Total Environment</i> , 2015, 536, 933-945.	3.9	32
99	Comparative analysis of the hepatopancreas transcriptome of grass carp (<i>Ctenopharyngodon idellus</i>) fed with lard oil and fish oil diets. <i>Gene</i> , 2015, 565, 192-200.	1.0	52
100	Molecular characterization, transcriptional activity and nutritional regulation of peroxisome proliferator activated receptor gamma in Nile tilapia (<i>Oreochromis niloticus</i>). <i>General and Comparative Endocrinology</i> , 2015, 223, 139-147.	0.8	25
101	Identification, characterization and nutritional regulation of two isoforms of acyl-coenzyme A oxidase 1 gene in Nile tilapia (<i>Oreochromis niloticus</i>). <i>Gene</i> , 2014, 545, 30-35.	1.0	33
102	Quantitative dietary isoleucine requirement of juvenile Pacific white shrimp, <i>Litopenaeus vannamei</i> (Boone) reared in low-salinity water. <i>Aquaculture International</i> , 2014, 22, 1481-1497.	1.1	23
103	Effects of temperature and salinity on metabolic rate of the Asiatic clam <i>Corbicula fluminea</i> (Müller). <i>TJ ETQq1 1 0.784314 rrgBT /Ov</i>	1.2	36
104	Urinary Loss of Tricarboxylic Acid Cycle Intermediates As Revealed by Metabolomics Studies: An Underlying Mechanism to Reduce Lipid Accretion by Whey Protein Ingestion?. <i>Journal of Proteome Research</i> , 2014, 13, 2560-2570.	1.8	42
105	Comparative Analysis of Fatty Acid Profiles in Brains and Eyes of Five Economic Fish Species in Winter and Summer. <i>Journal of Food and Nutrition Research (Newark, Del)</i> , 2014, 2, 722-730.	0.1	4
106	Different intakes of n-3 fatty acids among pregnant women in 3 regions of China with contrasting dietary patterns are reflected in maternal but not in umbilical erythrocyte phosphatidylcholine fatty acid composition. <i>Nutrition Research</i> , 2013, 33, 613-621.	1.3	13
107	Dietary eicosapentaenoic acid supplementation accentuates hepatic triglyceride accumulation in mice with impaired fatty acid oxidation capacity. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2013, 1831, 291-299.	1.2	39
108	Hydrolyzed Casein Reduces Diet-Induced Obesity in Male C57BL/6J Mice. <i>Journal of Nutrition</i> , 2013, 143, 1367-1375.	1.3	50

#	ARTICLE	IF	CITATIONS
109	Dietary inclusion of salmon, herring and pompano as oily fish reduces CVD risk markers in dyslipidaemic middle-aged and elderly Chinese women. <i>British Journal of Nutrition</i> , 2012, 108, 1455-1465.	1.2	53
110	Dietary glutathione as an antioxidant improves resistance to ammonia exposure in <i>Litopenaeus vannamei</i> . <i>Aquaculture Research</i> , 2012, 43, 311-316.	0.9	28
111	Risk-benefit evaluation of fish from Chinese markets: Nutrients and contaminants in 24 fish species from five big cities and related assessment for human health. <i>Science of the Total Environment</i> , 2012, 416, 187-199.	3.9	58
112	Effect of long-term administration of dietary β -1,3-glucan on growth, physiological, and immune responses in <i>Litopenaeus vannamei</i> (Boone, 1931). <i>Aquaculture International</i> , 2012, 20, 145-158.	1.1	18
113	Vaccenic and Elaidic Acid Equally Esterify into Triacylglycerols, but Differently into Phospholipids of Fed Rat Liver Cells. <i>Lipids</i> , 2011, 46, 647-657.	0.7	9
114	Nutritional Regulation of Bile Acid Metabolism Is Associated with Improved Pathological Characteristics of the Metabolic Syndrome. <i>Journal of Biological Chemistry</i> , 2011, 286, 28382-28395.	1.6	55
115	Depot-Dependent Effects of Adipose Tissue Explants on Co-Cultured Hepatocytes. <i>PLoS ONE</i> , 2011, 6, e20917.	1.1	27
116	Methionine limitation results in increased hepatic FAS activity, higher liver 18:1 to 18:0 fatty acid ratio and hepatic TAG accumulation in Atlantic salmon, <i>Salmo salar</i> . <i>Amino Acids</i> , 2010, 39, 449-460.	1.2	82
117	Dissimilar Properties of Vaccenic Versus Elaidic Acid in β -Oxidation Activities and Gene Regulation in Rat Liver Cells. <i>Lipids</i> , 2010, 45, 581-591.	0.7	12
118	Discrimination of ω -3 Rich Oils by Gas Chromatography. <i>Lipids</i> , 2010, 45, 1147-1158.	0.7	22
119	Unbound DHA causes a high blank value in β -oxidation assay: a concern for <i>in vitro</i> studies. <i>European Journal of Lipid Science and Technology</i> , 2010, 112, 333-342.	1.0	9
120	Persistent Organic Pollutant Exposure Leads to Insulin Resistance Syndrome. <i>Environmental Health Perspectives</i> , 2010, 118, 465-471.	2.8	326
121	β -oxidation modulates metabolic competition between eicosapentaenoic acid and arachidonic acid regulating prostaglandin E2 synthesis in rat hepatocytes-Kupffer cells. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2010, 1801, 526-536.	1.2	25
122	Inclusion of Atlantic salmon in the Chinese diet reduces cardiovascular disease risk markers in dyslipidemic adult men. <i>Nutrition Research</i> , 2010, 30, 447-454.	1.3	30
123	Elucidation of triacylglycerols in cod liver oil by liquid chromatography electrospray tandem ion-trap mass spectrometry. <i>Talanta</i> , 2010, 82, 1261-1270.	2.9	29
124	Toxic effects and residue of aflatoxin B1 in tilapia (<i>Oreochromis niloticus</i> - <i>O. aureus</i>) during long-term dietary exposure. <i>Aquaculture</i> , 2010, 307, 233-240.	1.7	145
125	The Intestinal Evacuation and Maximum Daily Consumption of Purified Formulated Diets by Juvenile Grass Carp (<i>Ctenopharyngodon idella</i>). <i>The Open Fish Science Journal</i> , 2009, 2, 1-5.	0.2	9
126	Effect of Dietary Energy to Protein Ratios on Growth Performance and Feed Efficiency of Juvenile Grass Carp (<i>Ctenopharyngodon idella</i>). <i>The Open Fish Science Journal</i> , 2009, 2, 25-31.	0.2	14

#	ARTICLE	IF	CITATIONS
127	Hypolipidaemic effects of fenofibrate and fasting in the herbivorous grass carp (<i>Ctenopharyngodon idella</i>) fed a high-fat diet. <i>British Journal of Nutrition</i> , 2008, 100, 1200-1212.	1.2	36
128	Regulation of Lipid Flux between Liver and Adipose Tissue during Transient Hepatic Steatosis in Carnitine-depleted Rats. <i>Journal of Biological Chemistry</i> , 2007, 282, 20816-20826.	1.6	34
129	Effect of dietary calcium and phosphorus on growth, feed efficiency, mineral content and body composition of juvenile grouper, <i>Epinephelus coioides</i> . <i>Aquaculture</i> , 2006, 255, 263-271.	1.7	97
130	The influence of feeding rate on growth, feed efficiency and body composition of juvenile grass carp (<i>Ctenopharyngodon idella</i>). <i>Aquaculture International</i> , 2006, 14, 247-257.	1.1	82
131	Biochemical hepatic alterations and body lipid composition in the herbivorous grass carp (<i>Ctenopharyngodon idella</i>) fed high-fat diets. <i>British Journal of Nutrition</i> , 2006, 95, 905-915.	1.2	154
132	Upregulation of liver VLDL receptor and FAT/CD36 expression in LDLR ^{-/-} /apoB100/100 mice fed trans-10,cis-12 conjugated linoleic acid. <i>Journal of Lipid Research</i> , 2006, 47, 2647-2655.	2.0	59
133	Effects of dietary carbohydrate level on growth and body composition of juvenile tilapia, <i>Oreochromis niloticus</i> × <i>O. aureus</i> . <i>Aquaculture Research</i> , 2005, 36, 1408-1413.	0.9	78
134	Effect of dietary lipid level on growth performance, lipid deposition, hepatic lipogenesis in juvenile cobia (<i>Rachycentron canadum</i>). <i>Aquaculture</i> , 2005, 249, 439-447.	1.7	289
135	Alteration of 20:5n-3 and 22:6n-3 fat contents and liver peroxisomal activities in fenofibrate-treated rainbow trout. <i>Lipids</i> , 2004, 39, 849-855.	0.7	27
136	Dietary Copper Requirement of Juvenile Oriental River Prawn <i>Macrobrachium nipponense</i> , and its Effects on Growth, Antioxidant Activities, and Resistance to <i>Aeromonas hydrophila</i> . <i>Israeli Journal of Aquaculture - Bamidgeh</i> , 0, 66, .	0.0	4