

Shufei Zang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7569095/publications.pdf>

Version: 2024-02-01

20
papers

578
citations

840776

11
h-index

752698

20
g-index

22
all docs

22
docs citations

22
times ranked

908
citing authors

#	ARTICLE	IF	CITATIONS
1	Targeting ferroptosis alleviates methionineâ€choline deficient (MCD)â€diet induced NASH by suppressing liver lipotoxicity. <i>Liver International</i> , 2020, 40, 1378-1394.	3.9	135
2	Lipocalin-2 mediates non-alcoholic steatohepatitis by promoting neutrophil-macrophage crosstalk via the induction of CXCR2. <i>Journal of Hepatology</i> , 2016, 65, 988-997.	3.7	134
3	Neutrophils Play a Crucial Role in the Early Stage of Nonalcoholic Steatohepatitis via Neutrophil Elastase in Mice. <i>Cell Biochemistry and Biophysics</i> , 2015, 73, 479-487.	1.8	52
4	Relationship of serum uric acid level with nonâ€alcoholic fatty liver disease and its inflammation progression in nonâ€obese adults. <i>Hepatology Research</i> , 2017, 47, E104-E112.	3.4	35
5	Increased ratio of neutrophil elastase to <i>â€antitrypsin</i> is closely associated with liver inflammation in patients with nonalcoholic steatohepatitis. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2016, 43, 13-21.	1.9	34
6	Knockout of neutrophil elastase protects against western diet induced nonalcoholic steatohepatitis in mice by regulating hepatic ceramides metabolism. <i>Biochemical and Biophysical Research Communications</i> , 2019, 518, 691-697.	2.1	30
7	The association between nonalcoholic fatty liver disease and risk of colorectal adenoma and cancer incident and recurrence: a meta-analysis of observational studies. <i>Expert Review of Gastroenterology and Hepatology</i> , 2019, 13, 385-395.	3.0	29
8	Elevated First-Trimester Neutrophil Count Is Closely Associated With the Development of Maternal Gestational Diabetes Mellitus and Adverse Pregnancy Outcomes. <i>Diabetes</i> , 2020, 69, 1401-1410.	0.6	27
9	Ratio of Conjugated Chenodeoxycholic to Muricholic Acids is Associated with Severity of Nonalcoholic Steatohepatitis. <i>Obesity</i> , 2019, 27, 2055-2066.	3.0	22
10	Berberine prevents non-alcoholic steatohepatitis-derived hepatocellular carcinoma by inhibiting inflammation and angiogenesis in mice. <i>American Journal of Translational Research (discontinued)</i> , 2019, 11, 2668-2682.	0.0	17
11	Haptoglobin Genotype and Vitaminâ€E Versus Placebo for the Treatment of Nondiabetic Patients with Nonalcoholic Steatohepatitis in China: A Multicenter, Randomized, Placebo-Controlled Trial Design. <i>Advances in Therapy</i> , 2018, 35, 218-231.	2.9	13
12	Decreased Monocyte Count Is Associated With Gestational Diabetes Mellitus Development, Macrosomia, and Inflammation. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, 107, 192-204.	3.6	10
13	Increased neutrophil count Is associated with the development of chronic kidney disease in patients with diabetes. <i>Journal of Diabetes</i> , 2022, 14, 442-454.	1.8	8
14	Haptoglobin 2-2 Genotype is Associated with More Advanced Disease in Subjects with Non-Alcoholic Steatohepatitis: A Retrospective Study. <i>Advances in Therapy</i> , 2019, 36, 880-895.	2.9	7
15	Serum ferritin as a risk factor for type 2 diabetes mellitus, regulated by liver transferrin receptor 2. <i>Endocrine Connections</i> , 2021, 10, 1513-1521.	1.9	6
16	Prealbumin to fibrinogen ratio is closely associated with diabetic peripheral neuropathy. <i>Endocrine Connections</i> , 2020, 9, 858-863.	1.9	5
17	Fibrosis Risk in Nonalcoholic Fatty Liver Disease Is Related to Chronic Kidney Disease in Older Type 2 Diabetes Patients. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, 107, e3661-e3669.	3.6	5
18	Alterations in CD8+ Tregs, CD56+ Natural Killer Cells and IL-10 Are Associated With Invasiveness of Nonfunctioning Pituitary Adenomas (NFPAs). <i>Pathology and Oncology Research</i> , 2021, 27, 598887.	1.9	3

#	ARTICLE	IF	CITATIONS
19	The effects of insulin therapy on maternal blood pressure and weight in women with gestational diabetes mellitus. <i>BMC Pregnancy and Childbirth</i> , 2021, 21, 657.	2.4	3
20	Increased Neutrophil elastase and proteinase 3 are closely associated with occurrence and severity of stroke and acute myocardial infarction in patients with type 2 diabetes mellitus. <i>Diabetes Research and Clinical Practice</i> , 2022, 186, 109853.	2.8	3