Masato Nagai

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7572701/publications.pdf

Version: 2024-02-01

172207 168136 2,998 62 29 53 citations h-index g-index papers 66 66 66 4890 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Does variety of social interactions associate with frequency of laughter among older people? The JAGES cross-sectional study. BMJ Open, 2021, 11, e039363.	0.8	2
2	The association between body mass index and recovery from post-traumatic stress disorder after the nuclear accident in Fukushima. Scientific Reports, 2021, 11, 5330.	1.6	2
3	dbTMM: an integrated database of large-scale cohort, genome and clinical data for the Tohoku Medical Megabank Project. Human Genome Variation, 2021, 8, 44.	0.4	7
4	Cohort Profile: Tohoku Medical Megabank Project Birth and Three-Generation Cohort Study (TMM) Tj ETQq0 0 0 2020, 49, 18-19m.	rgBT /Over 0.9	lock 10 Tf 50 107
5	Effectiveness of seasonal inactivated influenza vaccination in Japanese schoolchildren: an epidemiologic study at the community level. Human Vaccines and Immunotherapeutics, 2020, 16, 295-300.	1.4	2
6	Low birth weight and abnormal pre-pregnancy body mass index were at higher risk for hypertensive disorders of pregnancy. Pregnancy Hypertension, 2020, 22, 119-125.	0.6	5
7	Influence of post-disaster evacuation on incidence of hyperuricemia in residents of Fukushima Prefecture: the Fukushima Health Management Survey. Clinical and Experimental Nephrology, 2020, 24, 1025-1032.	0.7	7
8	Clustering by phenotype and genome-wide association study in autism. Translational Psychiatry, 2020, 10, 290.	2.4	29
9	Relationships among personality traits, metabolic syndrome, and metabolic syndrome scores: The Kakegawa cohort study. Journal of Psychosomatic Research, 2018, 107, 20-25.	1.2	12
10	Impact of evacuation onstrends in the prevalence, treatment, and control of hypertension before and after a disaster. Journal of Hypertension, 2018, 36, 924-932.	0.3	15
11	Associations of disaster-related and psychosocial factors with changes in smoking status after a disaster: a cross-sectional survey after the Great East Japan Earthquake. BMJ Open, 2018, 8, e018943.	0.8	14
12	Potential identification of vitamin B6 responsiveness in autism spectrum disorder utilizing phenotype variables and machine learning methods. Scientific Reports, 2018, 8, 14840.	1.6	18
13	The Relationship between Sleep Time and Mental Health Problems According to the Strengths and Difficulties Questionnaire in Children after an Earthquake Disaster: The Fukushima Health Management Survey. International Journal of Environmental Research and Public Health, 2018, 15, 633.	1.2	12
14	Impact of social relationships on income–laughter relationships among older people: the JAGES cross-sectional study. BMJ Open, 2018, 8, e019104.	0.8	7
15	Effect of evacuation on liver function after the Fukushima Daiichi Nuclear Power Plant accident: The Fukushima Health Management Survey. Journal of Epidemiology, 2017, 27, 180-185.	1.1	25
16	Fasting but not casual blood glucose is associated with pancreatic cancer mortality in Japanese: EPOCH-JAPAN. Cancer Causes and Control, 2017, 28, 625-633.	0.8	15
17	Changes of Posttraumatic Stress Responses in Evacuated Residents and Their Related Factors. Asia-Pacific Journal of Public Health, 2017, 29, 182S-192S.	0.4	21
18	Exercise Habits Are Important for the Mental Health of Children in Fukushima After the Fukushima Daiichi Disaster. Asia-Pacific Journal of Public Health, 2017, 29, 171S-181S.	0.4	17

#	Article	IF	CITATIONS
19	Changes in Cardiovascular Risk Factors After the Great East Japan Earthquake. Asia-Pacific Journal of Public Health, 2017, 29, 47S-55S.	0.4	64
20	The impact of evacuation on the incidence of chronic kidney disease after the Great East Japan Earthquake: The Fukushima Health Management Survey. Clinical and Experimental Nephrology, 2017, 21, 995-1002.	0.7	26
21	Effects of socioeconomic factors on cardiovascular-related symptoms among residents in Fukushima after the Great East Japan Earthquake: a cross-sectional study using data from the Fukushima Health Management Survey. BMJ Open, 2017, 7, e014077.	0.8	13
22	Lifestyle-related factors that explain disaster-induced changes in socioeconomic status and poor subjective health: a cross-sectional study from the Fukushima health management survey. BMC Public Health, 2017, 17, 340.	1.2	19
23	Evacuation after the Great East Japan Earthquake was associated with poor dietary intake: The Fukushima Health Management Survey. Journal of Epidemiology, 2017, 27, 14-23.	1.1	47
24	Influence of Post-disaster Evacuation on Incidence of Metabolic Syndrome. Journal of Atherosclerosis and Thrombosis, 2017, 24, 327-337.	0.9	48
25	Perception of Radiation Risk as a Predictor of Mid-Term Mental Health after a Nuclear Disaster: The Fukushima Health Management Survey. International Journal of Environmental Research and Public Health, 2017, 14, 1067.	1.2	32
26	Association of Total Energy Intake with 29-Year Mortality in the Japanese: NIPPON DATA80. Journal of Atherosclerosis and Thrombosis, 2016, 23, 339-354.	0.9	16
27	Threeâ€year trend survey of psychological distress, postâ€traumatic stress, and problem drinking among residents in the evacuation zone after the Fukushima Daiichi Nuclear Power Plant accident [The Fukushima Health Management Survey]. Psychiatry and Clinical Neurosciences, 2016, 70, 245-252.	1.0	74
28	Diagnostic interview study of the prevalence of depression among public employees engaged in longâ€term relief work in Fukushima. Psychiatry and Clinical Neurosciences, 2016, 70, 413-420.	1.0	24
29	Prevalence of Renal Dysfunction among Evacuees and Non-evacuees after the Great East Earthquake: Results from the Fukushima Health Management Survey. Internal Medicine, 2016, 55, 2563-2569.	0.3	12
30	Hypo-high-density Lipoprotein Cholesterolemia Caused by Evacuation after the Fukushima Daiichi Nuclear Power Plant Accident: Results from the Fukushima Health Management Survey. Internal Medicine, 2016, 55, 1967-1976.	0.3	34
31	Association between psychological distress and dietary intake among evacuees after the Great East Japan Earthquake in a cross-sectional study: the Fukushima Health Management Survey. BMJ Open, 2016, 6, e011534.	0.8	37
32	Serum Uric Acid and Mortality Form Cardiovascular Disease: EPOCH-JAPAN Study. Journal of Atherosclerosis and Thrombosis, 2016, 23, 692-703.	0.9	89
33	Predictors of severe psychological distress trajectory after nuclear disaster: evidence from the Fukushima Health Management Survey. BMJ Open, 2016, 6, e013400.	0.8	42
34	Secular trends of the impact of overweight and obesity on hypertension in Japan, 1980–2010. Hypertension Research, 2015, 38, 790-795.	1.5	39
35	Low-carbohydrate diets and cardiovascular and total mortality in Japanese: a 29-year follow-up of NIPPON DATA80. British Journal of Nutrition, 2014, 112, 916-924.	1.2	59
36	Consumption of fruits, vegetables, and seaweeds (sea vegetables) and pancreatic cancer risk: The Ohsaki Cohort Study. Cancer Epidemiology, 2014, 38, 129-136.	0.8	16

#	Article	IF	Citations
37	Long sleep duration and causeâ€specific mortality according to physical function and selfâ€rated health: the <scp>O</scp> hsaki <scp>C</scp> ohort <scp>S</scp> tudy. Journal of Sleep Research, 2013, 22, 209-216.	1.7	70
38	Association between sleep duration, weight gain, and obesity for long period. Sleep Medicine, 2013, 14, 206-210.	0.8	53
39	Lactation pattern and the risk for hormone-related female cancer in Japan. European Journal of Cancer Prevention, 2013, 22, 187-192.	0.6	13
40	Epidemiology of Hypertension in Japan. Circulation Journal, 2013, 77, 2226-2231.	0.7	155
41	Impact of Weight Change Since Age 20 and Cardiovascular Disease Mortality Risk. Circulation Journal, 2013, 77, 679-686.	0.7	14
42	Impact of obesity, overweight and underweight on life expectancy and lifetime medical expenditures: the Ohsaki Cohort Study. BMJ Open, 2012, 2, e000940.	0.8	40
43	Trends in Life Expectancy With Care Needs Based on Long-term Care Insurance Data in Japan. Journal of Epidemiology, 2012, 22, 238-243.	1.1	14
44	Gamma-Glutamyltransferase and Cancer Incidence: The Ohsaki Cohort Study. Journal of Epidemiology, 2012, 22, 144-150.	1.1	17
45	Gains in Disability-Free Life Expectancy From Elimination of Diseases and Injuries in Japan. Journal of Epidemiology, 2012, 22, 199-204.	1.1	15
46	History of cholelithiasis and the risk of prostate cancer: The Ohsaki Cohort Study. International Journal of Cancer, 2011, 128, 185-191.	2.3	9
47	Impact of walking on life expectancy and lifetime medical expenditure: the Ohsaki Cohort Study. BMJ Open, 2011, 1, e000240.	0.8	25
48	Association between Body-Mass Index and Risk of Death in More Than 1 Million Asians. New England Journal of Medicine, 2011, 364, 719-729.	13.9	730
49	Body Mass Index and Diabetes in Asia: A Cross-Sectional Pooled Analysis of 900,000 Individuals in the Asia Cohort Consortium. PLoS ONE, 2011, 6, e19930.	1.1	154
50	The Ohsaki Cohort 2006 Study: Design of Study and Profile of Participants at Baseline. Journal of Epidemiology, 2010, 20, 253-258.	1.1	70
51	Trends in Disability-Free Life Expectancy in Japan, 1995–2004. Journal of Epidemiology, 2010, 20, 308-312.	1.1	47
52	Effect of Age on the Association between Body Mass Index and All-Cause Mortality: The Ohsaki Cohort Study. Journal of Epidemiology, 2010, 20, 398-407.	1.1	46
53	History of diabetes mellitus and the risk of prostate cancer: the Ohsaki Cohort Study. Cancer Causes and Control, 2010, 21, 1025-1032.	0.8	64
54	Citrus consumption and cancer incidence: the Ohsaki cohort study. International Journal of Cancer, 2010, 127, 1913-1922.	2.3	52

#	Article	IF	CITATION
55	Estimating Visceral Fat Area by Multifrequency Bioelectrical Impedance. Diabetes Care, 2010, 33, 1077-1079.	4.3	54
56	Coffee Consumption and Mortality Due to All Causes, Cardiovascular Disease, and Cancer in Japanese Women. Journal of Nutrition, 2010, 140, 1007-1013.	1.3	88
57	Difference in lifetime medical expenditures between male smokers and non-smokers. Health Policy, 2010, 94, 84-89.	1.4	10
58	Participation in health check-ups and mortality using propensity score matched cohort analyses. Preventive Medicine, 2010, 51, 397-402.	1.6	43
59	Green tea consumption is associated with lower psychological distress in a general population: the Ohsaki Cohort 2006 Study. American Journal of Clinical Nutrition, 2009, 90, 1390-1396.	2.2	54
60	Factors Associated With Psychological Distress in a Community-Dwelling Japanese Population: The Ohsaki Cohort 2006 Study. Journal of Epidemiology, 2009, 19, 294-302.	1.1	64
61	Development of a New Method for Estimating Visceral Fat Area with Multi-Frequency Bioelectrical Impedance. Tohoku Journal of Experimental Medicine, 2008, 214, 105-112.	0.5	57
62	Developments in Estimating Visceral Fat Area from Medical Examination Data. Journal of Atherosclerosis and Thrombosis, 2008, 15, 193-198.	0.9	10