Paolo Scarpelli

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/757635/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	The Bile Acid Receptor GPBAR1 Regulates the M1/M2 Phenotype of Intestinal Macrophages and Activation of GPBAR1 Rescues Mice from Murine Colitis. Journal of Immunology, 2017, 199, 718-733.	0.8	198
2	Peroxynitrite Activates the NLRP3 Inflammasome Cascade in SOD1(G93A) Mouse Model of Amyotrophic Lateral Sclerosis. Molecular Neurobiology, 2018, 55, 2350-2361.	4.0	53
3	Metabolic Variability of a Multispecies Probiotic Preparation Impacts on the Anti-inflammatory Activity. Frontiers in Pharmacology, 2017, 8, 505.	3.5	49
4	Ursodeoxycholic acid is a GPBAR1 agonist and resets liver/intestinal FXR signaling in a model of diet-induced dysbiosis and NASH. Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids, 2019, 1864, 1422-1437.	2.4	37
5	GPBAR1 Functions as Gatekeeper for Liver NKT Cells and provides Counterregulatory Signals in Mouse Models of Immune-Mediated Hepatitis. Cellular and Molecular Gastroenterology and Hepatology, 2019, 8, 447-473.	4.5	37
6	Gpbar1 agonism promotes a Pgc-1α-dependent browning of white adipose tissue and energy expenditure and reverses diet-induced steatohepatitis in mice. Scientific Reports, 2017, 7, 13689.	3.3	36
7	<i>Saccharomyces cerevisiae</i> –based probiotic as novel anti-microbial agent for therapy of bacterial vaginosis. Virulence, 2018, 9, 954-966.	4.4	28
8	ROS-independent Nrf2 activation in prostate cancer. Oncotarget, 2017, 8, 67506-67518.	1.8	27
9	Disruption of TFGÎ ² -SMAD3 pathway by the nuclear receptor SHP mediates the antifibrotic activities of BAR704, a novel highly selective FXR ligand. Pharmacological Research, 2018, 131, 17-31.	7.1	25
10	Clostridium difficile toxin B induces senescence in enteric glial cells: A potential new mechanism of Clostridium difficile pathogenesis. Biochimica Et Biophysica Acta - Molecular Cell Research, 2018, 1865, 1945-1958.	4.1	24
11	Cyclo(His-Pro) inhibits NLRP3 inflammasome cascade in ALS microglial cells. Molecular and Cellular Neurosciences, 2019, 94, 23-31.	2.2	23
12	Extracellular Vesicles from Human Advanced-Stage Prostate Cancer Cells Modify the Inflammatory Response of Microenvironment-Residing Cells. Cancers, 2019, 11, 1276.	3.7	18
13	The bile acid activated receptors GPBAR1 and FXR exert antagonistic effects on autophagy. FASEB Journal, 2021, 35, e21271.	0.5	15
14	Nanotraps with biomimetic surface as decoys for chemokines. Nanomedicine: Nanotechnology, Biology, and Medicine, 2017, 13, 2575-2585.	3.3	14
15	Selected cholesterol biosynthesis inhibitors produce accumulation of the intermediate FF-MAS that targets nucleus and activates LXRα in HepG2 cells. Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids, 2017, 1862, 842-852.	2.4	12
16	Palmitate lipotoxicity in enteric glial cells: Lipid remodeling and mitochondrial ROS are responsible for cyt c release outside mitochondria. Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids, 2018, 1863, 895-908.	2.4	12
17	Amniotic fluid stem cellâ€derived extracellular vesicles are independent metabolic units capable of modulating inflammasome activation in THPâ€1 cells. FASEB Journal, 2022, 36, e22218.	0.5	11
18	Extracellular Vesicles-Mediated Transfer of miRNA Let-7b from PC3 Cells to Macrophages. Genes, 2020, 11, 1495.	2.4	9

#	Article	IF	CITATIONS
19	Tm7sf2 gene promotes adipocyte differentiation of mouse embryonic fibroblasts and improves insulin sensitivity. Biochimica Et Biophysica Acta - Molecular Cell Research, 2021, 1868, 118897.	4.1	8