

Luis A Garca-Rodrguez

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7581840/luis-a-garcia-rodriguez-publications-by-year.pdf>
Version: 2024-04-11

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| | | | |
|--------------------|--------------------------|----------------|-----------------|
| 199 papers | 10,893 citations | 51 h-index | 100 g-index |
| 209 ext. papers | 12,325 ext. citations | 5.3 avg, IF | 6.42 L-index |

| # | Paper | IF | Citations |
|-----|---|------|-----------|
| 199 | Safety and effectiveness of appropriately and inappropriately dosed rivaroxaban or apixaban versus warfarin in patients with atrial fibrillation: a cohort study with nested case-control analyses from UK primary care. <i>BMJ Open</i> , 2022 , 12, e059311 | 3 | |
| 198 | Trends in the use of oral anticoagulants, antiplatelets and statins in four European countries: a population-based study. <i>European Journal of Clinical Pharmacology</i> , 2021 , 1 | 2.8 | 3 |
| 197 | Trends in Incidence of Intracerebral Hemorrhage and Association With Antithrombotic Drug Use in Denmark, 2005-2018. <i>JAMA Network Open</i> , 2021 , 4, e218380 | 10.4 | 2 |
| 196 | Validation of venous thromboembolism diagnoses in patients receiving rivaroxaban or warfarin in The Health Improvement Network. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 229-236 | 2.6 | 2 |
| 195 | Use of non-vitamin K antagonist oral anticoagulants in Colombia: A descriptive study using a national administrative healthcare database. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 426-434 | 2.6 | 2 |
| 194 | Public health impact of low-dose aspirin on colorectal cancer, cardiovascular disease and safety in the UK - Results from micro-simulation model. <i>IJC Heart and Vasculature</i> , 2021 , 36, 100851 | 2.4 | 1 |
| 193 | The Validity of Intracerebral Hemorrhage Diagnoses in the Danish Patient Registry and the Danish Stroke Registry. <i>Clinical Epidemiology</i> , 2020 , 12, 1313-1325 | 5.9 | 4 |
| 192 | Reduction in use of cyproterone/ethinylestradiol (Diane-35 and generics) after risk minimization measures in the Netherlands, UK and Italy. <i>Contraception</i> , 2020 , 102, 243-245 | 2.5 | |
| 191 | Risk of cardiovascular events and death associated with initiation of SGLT2 inhibitors compared with DPP-4 inhibitors: an analysis from the CVD-REAL 2 multinational cohort study. <i>Lancet Diabetes and Endocrinology</i> , 2020 , 8, 606-615 | 18.1 | 37 |
| 190 | Bleeding associated with low-dose aspirin: Comparison of data from the COMPASS randomized controlled trial and routine clinical practice. <i>International Journal of Cardiology</i> , 2020 , 318, 21-24 | 3.2 | 3 |
| 189 | No association between use of phosphodiesterase 5 inhibitors and colorectal cancer in men with erectile dysfunction. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 605-608 | 2.6 | 4 |
| 188 | Low-dose aspirin and risk of gastric and oesophageal cancer: A population-based study in the United Kingdom using The Health Improvement Network. <i>International Journal of Cancer</i> , 2020 , 147, 2394-2404 | 7.5 | 6 |
| 187 | Antidepressant drug use and subdural hematoma risk. <i>Journal of Thrombosis and Haemostasis</i> , 2020 , 18, 318-327 | 15.4 | 4 |
| 186 | Kidney outcomes associated with use of SGLT2 inhibitors in real-world clinical practice (CVD-REAL 3): a multinational observational cohort study. <i>Lancet Diabetes and Endocrinology</i> , 2020 , 8, 27-35 | 18.1 | 109 |
| 185 | Impact of chronic kidney disease definition on assessment of its incidence and risk factors in patients with newly diagnosed type 1 and type 2 diabetes in the UK: A cohort study using primary care data from the United Kingdom. <i>Primary Care Diabetes</i> , 2020 , 14, 381-387 | 2.4 | 7 |
| 184 | Feasibility study to identify women of childbearing age at risk of pregnancy not using any contraception in The Health Improvement Network (THIN) database. <i>BMC Medical Informatics and Decision Making</i> , 2020 , 20, 164 | 3.6 | 0 |
| 183 | Rationale and design of a European epidemiological post-authorization safety study (PASS) program: rivaroxaban use in routine clinical practice. <i>Expert Opinion on Drug Safety</i> , 2020 , 19, 1513-1520 | 4.1 | 2 |

| | | | |
|-----|---|-----|----|
| 182 | Risk of bleeding events among patients with systemic sclerosis and the general population in the UK: a large population-based cohort study. <i>Clinical Rheumatology</i> , 2020 , 39, 19-26 | 3.9 | 0 |
| 181 | Effect of Proton Pump Inhibitors on Risks of Upper and Lower Gastrointestinal Bleeding among Users of Low-Dose Aspirin: A Population-Based Observational Study. <i>Journal of Clinical Medicine</i> , 2020 , 9, | 5.1 | 3 |
| 180 | Discontinuation of oral anticoagulation in atrial fibrillation and risk of ischaemic stroke. <i>Heart</i> , 2020 , , | 5.1 | 5 |
| 179 | Development of hypotension in patients newly diagnosed with heart failure in UK general practice: retrospective cohort and nested case-control analyses. <i>BMJ Open</i> , 2019 , 9, e028750 | 3 | 2 |
| 178 | The effect of low-dose aspirin on colorectal cancer prevention and gastrointestinal bleeding according to bodyweight and body mass index: Analysis of UK primary care data. <i>International Journal of Cardiology</i> , 2019 , 297, 135-139 | 3.2 | 3 |
| 177 | Clopidogrel and Low-Dose Aspirin, Alone or Together, Reduce Risk of Colorectal Cancer. <i>Clinical Gastroenterology and Hepatology</i> , 2019 , 17, 2024-2033.e2 | 6.9 | 26 |
| 176 | Low-dose aspirin and risk of upper/lower gastrointestinal bleeding by bleed severity: a cohort study with nested case-control analysis using primary care electronic health records from the United Kingdom. <i>Annals of Medicine</i> , 2019 , 51, 182-192 | 1.5 | 6 |
| 175 | Ascertainment and validation of major bleeding events in a primary care database. <i>Pharmacoepidemiology and Drug Safety</i> , 2019 , 28, 148-155 | 2.6 | 10 |
| 174 | Population-based case-control study: chemoprotection of colorectal cancer with non-aspirin nonsteroidal anti-inflammatory drugs and other drugs for pain control. <i>Alimentary Pharmacology and Therapeutics</i> , 2019 , 50, 295-305 | 6.1 | 6 |
| 173 | Feasibility and validity of The Health Improvement Network database of primary care electronic health records to identify and characterise patients with small cell lung cancer in the United Kingdom. <i>BMC Cancer</i> , 2019 , 19, 91 | 4.8 | 1 |
| 172 | Appropriateness of initial dose of non-vitamin K antagonist oral anticoagulants in patients with non-valvular atrial fibrillation in the UK. <i>BMJ Open</i> , 2019 , 9, e031341 | 3 | 21 |
| 171 | Discontinuation of non-Vitamin K antagonist oral anticoagulants in patients with non-valvular atrial fibrillation: a population-based cohort study using primary care data from The Health Improvement Network in the UK. <i>BMJ Open</i> , 2019 , 9, e031342 | 3 | 9 |
| 170 | Risk of Acute Myocardial Infarction Among New Users of Allopurinol According to Serum Urate Level: A Nested Case-Control Study. <i>Journal of Clinical Medicine</i> , 2019 , 8, | 5.1 | 5 |
| 169 | The importance of validating intracranial bleeding diagnoses in The Health Improvement Network, United Kingdom: Misclassification of onset and its impact on the risk associated with low-dose aspirin therapy. <i>Pharmacoepidemiology and Drug Safety</i> , 2019 , 28, 134-139 | 2.6 | 5 |
| 168 | Incidence of Upper and Lower Gastrointestinal Bleeding in New Users of Low-Dose Aspirin. <i>Clinical Gastroenterology and Hepatology</i> , 2019 , 17, 887-895.e6 | 6.9 | 16 |
| 167 | Predictors of Over-Anticoagulation in Warfarin Users in the UK General Population: A Nested Case-Control Study in a Primary Health Care Database. <i>Thrombosis and Haemostasis</i> , 2019 , 119, 66-76 | 7 | 10 |
| 166 | Validation study of colorectal cancer diagnosis in the Spanish primary care database, BIFAP. <i>Pharmacoepidemiology and Drug Safety</i> , 2019 , 28, 209-216 | 2.6 | 9 |
| 165 | Contemporary epidemiology of systemic sclerosis: A population-based cohort study in the United Kingdom. <i>Seminars in Arthritis and Rheumatism</i> , 2019 , 49, 105-111 | 5.3 | 6 |

| | | | |
|-----|---|------|-----|
| 164 | Safety of non-insulin glucose-lowering drugs in pregnant women with pre-gestational diabetes: A cohort study. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 1642-1651 | 6.7 | 9 |
| 163 | Lack of Association Between Proton Pump Inhibitor Use and Cognitive Decline. <i>Clinical Gastroenterology and Hepatology</i> , 2018 , 16, 681-689 | 6.9 | 21 |
| 162 | Population Impact of Drug Interactions with Warfarin: A Real-World Data Approach. <i>Thrombosis and Haemostasis</i> , 2018 , 118, 461-470 | 7 | 14 |
| 161 | Time trends in peripheral artery disease incidence, prevalence and secondary preventive therapy: a cohort study in The Health Improvement Network in the UK. <i>BMJ Open</i> , 2018 , 8, e018184 | 3 | 30 |
| 160 | Electronic medical records can be used to emulate target trials of sustained treatment strategies. <i>Journal of Clinical Epidemiology</i> , 2018 , 96, 12-22 | 5.7 | 47 |
| 159 | Predictors of Bleeding in Patients with Symptomatic Peripheral Artery Disease: A Cohort Study Using The Health Improvement Network in the United Kingdom. <i>Thrombosis and Haemostasis</i> , 2018 , 118, 1101-1112 | 7 | 4 |
| 158 | Cardiovascular Events Associated With SGLT-2 Inhibitors Versus Other Glucose-Lowering Drugs: The CVD-REAL 2 Study. <i>Journal of the American College of Cardiology</i> , 2018 , 71, 2628-2639 | 15.1 | 263 |
| 157 | Trends in the contemporary incidence of colorectal cancer and patient characteristics in the United Kingdom: a population-based cohort study using The Health Improvement Network. <i>BMC Cancer</i> , 2018 , 18, 402 | 4.8 | 11 |
| 156 | Real world management of pregestational diabetes not achieving glycemic control for many patients in the UK. <i>Pharmacoepidemiology and Drug Safety</i> , 2018 , 27, 940-948 | 2.6 | 3 |
| 155 | Intracerebral hemorrhage: positive predictive value of diagnosis codes in two nationwide Danish registries. <i>Clinical Epidemiology</i> , 2018 , 10, 941-948 | 5.9 | 20 |
| 154 | Hormonal contraception is not associated with increased risk for seizures in the general population: results from a cohort study using The Health Improvement Network. <i>European Journal of Clinical Pharmacology</i> , 2018 , 74, 1175-1180 | 2.8 | 3 |
| 153 | Association of Antithrombotic Drug Use With Subdural Hematoma Risk. <i>JAMA - Journal of the American Medical Association</i> , 2017 , 317, 836-846 | 27.4 | 77 |
| 152 | Characteristics of children and adolescents first prescribed proton pump inhibitors or histamine-2-receptor antagonists: an observational cohort study. <i>Current Medical Research and Opinion</i> , 2017 , 33, 2251-2259 | 2.5 | 2 |
| 151 | Incidence of intracranial bleeds in new users of low-dose aspirin: a cohort study using The Health Improvement Network. <i>Journal of Thrombosis and Haemostasis</i> , 2017 , 15, 1055-1064 | 15.4 | 10 |
| 150 | Statins and polyneuropathy revisited: case-control study in Denmark, 1999-2013. <i>British Journal of Clinical Pharmacology</i> , 2017 , 83, 2087-2095 | 3.8 | 14 |
| 149 | Mortality in patients who discontinue low-dose acetylsalicylic acid therapy after upper gastrointestinal bleeding. <i>Pharmacoepidemiology and Drug Safety</i> , 2017 , 26, 215-222 | 2.6 | 3 |
| 148 | Prescribing of cyproterone acetate/ethinylestradiol in UK general practice: a retrospective descriptive study using The Health Improvement Network. <i>Contraception</i> , 2017 , 95, 299-305 | 2.5 | 6 |
| 147 | Statins and the Risk of Intracerebral Hemorrhage in Patients With Previous Ischemic Stroke or Transient Ischemic Attack. <i>Stroke</i> , 2017 , 48, 3245-3251 | 6.7 | 12 |

| | | | |
|-----|---|-----|----|
| 146 | Incidence, treatment and recurrence of endometriosis in a UK-based population analysis using data from The Health Improvement Network and the Hospital Episode Statistics database. <i>European Journal of Contraception and Reproductive Health Care</i> , 2017 , 22, 334-343 | 1.8 | 14 |
| 145 | Incidence of colorectal cancer in new users and non-users of low-dose aspirin without existing cardiovascular disease: A cohort study using The Health Improvement Network. <i>International Journal of Cardiology</i> , 2017 , 248, 376-381 | 3.2 | 4 |
| 144 | Low-dose aspirin and risk of intracranial bleeds: An observational study in UK general practice. <i>Neurology</i> , 2017 , 89, 2280-2287 | 6.5 | 25 |
| 143 | Cardiovascular outcomes for patients with symptomatic peripheral artery disease: A cohort study in The Health Improvement Network (THIN) in the UK. <i>European Journal of Preventive Cardiology</i> , 2017 , 24, 1927-1937 | 3.9 | 17 |
| 142 | New use of low-dose aspirin and risk of colorectal cancer by stage at diagnosis: a nested case-control study in UK general practice. <i>BMC Cancer</i> , 2017 , 17, 637 | 4.8 | 25 |
| 141 | Utilization and safety of proton-pump inhibitors and histamine-2 receptor antagonists in children and adolescents: an observational cohort study. <i>Current Medical Research and Opinion</i> , 2017 , 33, 2201-2209 | 2.5 | 5 |
| 140 | Low-Dose Aspirin after an Episode of Haemorrhagic Stroke Is Associated with Improved Survival. <i>Thrombosis and Haemostasis</i> , 2017 , 117, 2396-2405 | 7 | 7 |
| 139 | Risk of bleeding after hospitalization for a serious coronary event: a retrospective cohort study with nested case-control analyses. <i>BMC Cardiovascular Disorders</i> , 2016 , 16, 164 | 2.3 | 8 |
| 138 | Subdural hematoma cases identified through a Danish patient register: diagnosis validity, clinical characteristics, and preadmission antithrombotic drug use. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25, 1253-1262 | 2.6 | 18 |
| 137 | Type 2 diabetes, high blood pressure and acute pancreatitis. <i>Hepatobiliary and Pancreatic Diseases International</i> , 2016 , 15, 443-5 | 2.1 | 2 |
| 136 | Impact of hyperkalaemia definition on incidence assessment: implications for epidemiological research based on a large cohort study in newly diagnosed heart failure patients in primary care. <i>BMC Family Practice</i> , 2016 , 17, 51 | 2.6 | 6 |
| 135 | Incidence and risk factors for severe renal impairment after first diagnosis of heart failure: A cohort and nested case-control study in UK general practice. <i>International Journal of Cardiology</i> , 2016 , 207, 252-7 | 3.2 | 8 |
| 134 | Association between low-dose acetylsalicylic acid reinitiation and the risk of myocardial infarction or coronary heart disease death. <i>European Journal of Preventive Cardiology</i> , 2016 , 23, 1029-36 | 3.9 | 0 |
| 133 | The Protective Effect of Low-Dose Aspirin against Colorectal Cancer Is Unlikely Explained by Selection Bias: Results from Three Different Study Designs in Clinical Practice. <i>PLoS ONE</i> , 2016 , 11, e0159179 | 3.7 | 24 |
| 132 | Bleeding Risk with Long-Term Low-Dose Aspirin: A Systematic Review of Observational Studies. <i>PLoS ONE</i> , 2016 , 11, e0160046 | 3.7 | 93 |
| 131 | Heart failure hospitalization: An important prognostic factor for heart failure re-admission and mortality. <i>International Journal of Cardiology</i> , 2016 , 220, 855-61 | 3.2 | 19 |
| 130 | Incidence and risk factors for atrial fibrillation in patients with newly diagnosed heart failure. <i>Journal of Cardiovascular Medicine</i> , 2016 , 17, 608-15 | 1.9 | 5 |
| 129 | Prescribing patterns of combined hormonal products containing cyproterone acetate, levonorgestrel and drospirenone in the UK. <i>Journal of Family Planning and Reproductive Health Care</i> , 2016 , 42, 247-254 | | 3 |

| | | | |
|-----|---|------|----|
| 128 | Reply to Comment on New-onset type 2 diabetes, elevated HbA1c, anti-diabetic medications, and risk of pancreatic cancer. <i>British Journal of Cancer</i> , 2016 , 114, e12 | 8.7 | |
| 127 | Use of tricyclic antidepressants and risk of glioma: a nationwide case-control study. <i>British Journal of Cancer</i> , 2016 , 114, 1265-8 | 8.7 | 12 |
| 126 | Validation of low-dose aspirin prescription data in The Health Improvement Network: how much misclassification due to over-the-counter use?. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25, 392-8 | 2.6 | 26 |
| 125 | Risk of seizure associated with use of acid-suppressive drugs: An observational cohort study. <i>Epilepsy and Behavior</i> , 2016 , 62, 72-80 | 3.2 | 6 |
| 124 | Do case-only designs yield consistent results across design and different databases? A case study of hip fractures and benzodiazepines. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 79-87 | 2.6 | 11 |
| 123 | Prevalence of antibiotic use: a comparison across various European health care data sources. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 11-20 | 2.6 | 40 |
| 122 | Risk of acute liver injury associated with use of antibiotics. Comparative cohort and nested case-control studies using two primary care databases in Europe. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 29-38 | 2.6 | 14 |
| 121 | Validity and completeness of colorectal cancer diagnoses in a primary care database in the United Kingdom. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25, 385-91 | 2.6 | 23 |
| 120 | Allopurinol use and risk of non-fatal acute myocardial infarction. <i>Heart</i> , 2015 , 101, 679-85 | 5.1 | 37 |
| 119 | Incidence and Predictors of Hemorrhagic Stroke in Users of Low-Dose Acetylsalicylic Acid. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2015 , 24, 2321-8 | 2.8 | 4 |
| 118 | Risk factors for hyperkalaemia in a cohort of patients with newly diagnosed heart failure: a nested case-control study in UK general practice. <i>European Journal of Heart Failure</i> , 2015 , 17, 205-13 | 12.3 | 46 |
| 117 | Cardiovascular events and all-cause mortality in a cohort of 57,946 patients with type 2 diabetes: associations with renal function and cardiovascular risk factors. <i>Cardiovascular Diabetology</i> , 2015 , 14, 38 | 8.7 | 47 |
| 116 | Sa1971 Low-Dose Acetylsalicylic Acid and Risk of Colorectal Cancer: Evidence From a Primary Care Study in the UK. <i>Gastroenterology</i> , 2015 , 148, S-371 | 13.3 | 3 |
| 115 | The incidence of hysterectomy, uterus-preserving procedures and recurrent treatment in the management of uterine fibroids. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2015 , 194, 147-52 | 2.4 | 9 |
| 114 | The continuation rates of long-acting reversible contraceptives in UK general practice using data from The Health Improvement Network. <i>Pharmacoepidemiology and Drug Safety</i> , 2015 , 24, 52-8 | 2.6 | 16 |
| 113 | Antithrombotic drugs and subarachnoid haemorrhage risk. A nationwide case-control study in Denmark. <i>Thrombosis and Haemostasis</i> , 2015 , 114, 1064-75 | 7 | 26 |
| 112 | Patterns of Antiplatelet Therapy in Patients Who Have Experienced an Acute Coronary Event: A Descriptive Study in UK Primary Care. <i>Journal of Cardiovascular Pharmacology and Therapeutics</i> , 2015 , 20, 378-86 | 2.6 | 2 |
| 111 | New-onset type 2 diabetes, elevated HbA1c, anti-diabetic medications, and risk of pancreatic cancer. <i>British Journal of Cancer</i> , 2015 , 113, 1607-14 | 8.7 | 44 |

| | | | |
|-----|---|-----|-----|
| 110 | Reply: To PMID 25460554. <i>Clinical Gastroenterology and Hepatology</i> , 2015 , 13, 2024-5 | 6.9 | |
| 109 | Risk of upper and lower gastrointestinal bleeding in patients taking nonsteroidal anti-inflammatory drugs, antiplatelet agents, or anticoagulants. <i>Clinical Gastroenterology and Hepatology</i> , 2015 , 13, 906-12.e2 | 6.9 | 151 |
| 108 | Diabetes, cardiovascular morbidity, and risk of age-related macular degeneration in a primary care population. <i>Investigative Ophthalmology and Visual Science</i> , 2015 , 56, 1585-92 | | 23 |
| 107 | Risk of nonfatal acute myocardial infarction associated with non-steroidal antiinflammatory drugs, non-narcotic analgesics and other drugs used in osteoarthritis: a nested case-control study. <i>Pharmacoepidemiology and Drug Safety</i> , 2014 , 23, 1128-38 | 2.6 | 40 |
| 106 | Low-dose aspirin and upper gastrointestinal bleeding in primary versus secondary cardiovascular prevention: a population-based, nested case-control study. <i>Circulation: Cardiovascular Quality and Outcomes</i> , 2014 , 7, 70-7 | 5.8 | 12 |
| 105 | Patterns in the use of low-dose acetylsalicylic acid and other therapies following upper gastrointestinal bleeding. <i>American Journal of Cardiovascular Drugs</i> , 2014 , 14, 443-50 | 4 | 5 |
| 104 | Incidence rates and trends of hip/femur fractures in five European countries: comparison using e-healthcare records databases. <i>Calcified Tissue International</i> , 2014 , 94, 580-9 | 3.9 | 36 |
| 103 | Represcription of low-dose acetylsalicylic acid after discontinuation in patients receiving treatment for secondary cardiovascular disease prevention in the UK. <i>American Journal of Cardiovascular Drugs</i> , 2014 , 14, 319-26 | 4 | 4 |
| 102 | Migraine and risk of hemorrhagic stroke: a study based on data from general practice. <i>Journal of Headache and Pain</i> , 2014 , 15, 74 | 8.8 | 19 |
| 101 | Risk of uncomplicated peptic ulcer disease in a cohort of new users of low-dose acetylsalicylic acid for secondary prevention of cardiovascular events. <i>BMC Gastroenterology</i> , 2014 , 14, 205 | 3 | 8 |
| 100 | Use of proton pump inhibitors and the risk of coronary events in new users of low-dose acetylsalicylic acid in UK primary care. <i>Thrombosis and Haemostasis</i> , 2014 , 111, 131-9 | 7 | 8 |
| 99 | Study of long-acting reversible contraceptive use in a UK primary care database: validation of methodology. <i>European Journal of Contraception and Reproductive Health Care</i> , 2014 , 19, 22-8 | 1.8 | 7 |
| 98 | Prescription of antihypertensive medications during pregnancy in the UK. <i>Pharmacoepidemiology and Drug Safety</i> , 2014 , 23, 1051-8 | 2.6 | 13 |
| 97 | Study of the association between hip fracture and acid-suppressive drug use in a UK primary care setting. <i>Pharmacotherapy</i> , 2014 , 34, 570-81 | 5.8 | 30 |
| 96 | Use of long-acting reversible contraceptives in the UK from 2004 to 2010: analysis using The Health Improvement Network Database. <i>European Journal of Contraception and Reproductive Health Care</i> , 2014 , 19, 439-47 | 1.8 | 12 |
| 95 | Risk factors associated with uncomplicated peptic ulcer and changes in medication use after diagnosis. <i>PLoS ONE</i> , 2014 , 9, e101768 | 3.7 | 13 |
| 94 | Upper gastrointestinal bleeding associated with NSAIDs, other drugs and interactions: a nested case-control study in a new general practice database. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 691-701 | 2.8 | 57 |
| 93 | Risk of complications after a peptic ulcer diagnosis: effectiveness of proton pump inhibitors. <i>Digestive Diseases and Sciences</i> , 2013 , 58, 1653-62 | 4 | 8 |

| | | | |
|----|--|------|-----|
| 92 | Statins and risk of diabetes: an analysis of electronic medical records to evaluate possible bias due to differential survival. <i>Diabetes Care</i> , 2013 , 36, 1236-40 | 14.6 | 51 |
| 91 | Effect of baseline gastrointestinal risk and use of proton pump inhibitors on frequency of discontinuation of aspirin for secondary cardiovascular prevention in United kingdom primary care. <i>American Journal of Cardiology</i> , 2013 , 112, 1075-82 | 3 | 14 |
| 90 | Warfarin and absolute risk of hemorrhagic stroke. <i>Cmaj</i> , 2013 , 185, 687 | 3.5 | 2 |
| 89 | Antithrombotic drugs and risk of hemorrhagic stroke in the general population. <i>Neurology</i> , 2013 , 81, 566-74 | 6.5 | 105 |
| 88 | Risk factors for falls with use of acid-suppressive drugs. <i>Epidemiology</i> , 2013 , 24, 600-7 | 3.1 | 11 |
| 87 | Observational data for comparative effectiveness research: an emulation of randomised trials of statins and primary prevention of coronary heart disease. <i>Statistical Methods in Medical Research</i> , 2013 , 22, 70-96 | 2.3 | 124 |
| 86 | Mortality after hemorrhagic stroke: data from general practice (The Health Improvement Network). <i>Neurology</i> , 2013 , 81, 559-65 | 6.5 | 81 |
| 85 | Challenges of using primary care electronic medical records in the UK to study medications in pregnancy. <i>Pharmacoepidemiology and Drug Safety</i> , 2013 , 22, 977-85 | 2.6 | 19 |
| 84 | Coxibs: pharmacology, toxicity and efficacy in cancer clinical trials. <i>Recent Results in Cancer Research</i> , 2013 , 191, 67-93 | 1.5 | 33 |
| 83 | Incidence of hemorrhagic stroke in the general population: validation of data from The Health Improvement Network. <i>Pharmacoepidemiology and Drug Safety</i> , 2013 , 22, 176-82 | 2.6 | 36 |
| 82 | Cardiovascular and upper gastrointestinal bleeding consequences of low-dose acetylsalicylic acid discontinuation. <i>Thrombosis and Haemostasis</i> , 2013 , 110, 1298-304 | 7 | 25 |
| 81 | Use of clopidogrel and proton pump inhibitors after a serious acute coronary event: risk of coronary events and peptic ulcer bleeding. <i>Thrombosis and Haemostasis</i> , 2013 , 110, 1014-24 | 7 | 13 |
| 80 | Discontinuation of low-dose acetylsalicylic acid therapy in UK primary care: incidence and predictors in patients with cardiovascular disease. <i>Journal of Pragmatic and Observational Research</i> , 2012 , 3, 1-9 | 7.4 | 5 |
| 79 | Predictors and time trends in clopidogrel and proton pump inhibitor coprescription with low-dose acetylsalicylic acid. <i>Pharmacoepidemiology and Drug Safety</i> , 2012 , 21, 463-9 | 2.6 | 1 |
| 78 | Validation of diabetic retinopathy and maculopathy diagnoses recorded in a U.K. primary care database. <i>Diabetes Care</i> , 2012 , 35, 762-7 | 14.6 | 27 |
| 77 | Androgen deprivation therapy and the risk of coronary heart disease and heart failure in patients with prostate cancer: a nested case-control study in UK primary care. <i>Drug Safety</i> , 2011 , 34, 1061-77 | 5.1 | 50 |
| 76 | Acid suppressants reduce risk of gastrointestinal bleeding in patients on antithrombotic or anti-inflammatory therapy. <i>Gastroenterology</i> , 2011 , 141, 71-9 | 13.3 | 74 |
| 75 | Confounding adjustment via a semi-automated high-dimensional propensity score algorithm: an application to electronic medical records. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 849-57 | 2.6 | 66 |

| | | | |
|----|--|------|-----|
| 74 | Systematic review of peptic ulcer disease incidence rates: do studies without validation provide reliable estimates?. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 718-28 | 2.6 | 49 |
| 73 | Low-dose naproxen interferes with the antiplatelet effects of aspirin in healthy subjects: recommendations to minimize the functional consequences. <i>Arthritis and Rheumatism</i> , 2011 , 63, 850-9 | | 47 |
| 72 | Risk of upper gastrointestinal bleeding with low-dose acetylsalicylic acid alone and in combination with clopidogrel and other medications. <i>Circulation</i> , 2011 , 123, 1108-15 | 16.7 | 135 |
| 71 | Discontinuation of low dose aspirin and risk of myocardial infarction: case-control study in UK primary care. <i>BMJ, The</i> , 2011 , 343, d4094 | 5.9 | 97 |
| 70 | Increased risk of stroke after discontinuation of acetylsalicylic acid: a UK primary care study. <i>Neurology</i> , 2011 , 76, 740-6 | 6.5 | 88 |
| 69 | NSAID use selectively increases the risk of non-fatal myocardial infarction: a systematic review of randomised trials and observational studies. <i>PLoS ONE</i> , 2011 , 6, e16780 | 3.7 | 35 |
| 68 | Risk of Upper Gastrointestinal Bleeding in a Cohort of New Users of Low-Dose ASA for Secondary Prevention of Cardiovascular Outcomes. <i>Frontiers in Pharmacology</i> , 2010 , 1, 126 | 5.6 | 27 |
| 67 | Gastroesophageal reflux disease in children and adolescents in primary care. <i>Scandinavian Journal of Gastroenterology</i> , 2010 , 45, 139-46 | 2.4 | 60 |
| 66 | Acute pancreatitis in association with type 2 diabetes and antidiabetic drugs: a population-based cohort study. <i>Diabetes Care</i> , 2010 , 33, 2580-5 | 14.6 | 105 |
| 65 | Long-term use of anti-inflammatory drugs and risk of atrial fibrillation. <i>Archives of Internal Medicine</i> , 2010 , 170, 1450-5 | | 44 |
| 64 | Prospective drug safety monitoring using the UK primary-care General Practice Research Database: theoretical framework, feasibility analysis and extrapolation to future scenarios. <i>Drug Safety</i> , 2010 , 33, 223-32 | 5.1 | 17 |
| 63 | Heart failure, myocardial infarction, lung cancer and death in COPD patients: a UK primary care study. <i>Respiratory Medicine</i> , 2010 , 104, 1691-9 | 4.6 | 44 |
| 62 | Case validation in research using large databases. <i>British Journal of General Practice</i> , 2010 , 60, 160-1 | 1.6 | 15 |
| 61 | Use of non-steroidal antiinflammatory drugs and type-specific risk of acute coronary syndrome. <i>American Journal of Cardiology</i> , 2010 , 105, 1102-6 | 3 | 35 |
| 60 | Variability among nonsteroidal antiinflammatory drugs in risk of upper gastrointestinal bleeding. <i>Arthritis and Rheumatism</i> , 2010 , 62, 1592-601 | | 173 |
| 59 | Validation of ischemic cerebrovascular diagnoses in the health improvement network (THIN). <i>Pharmacoepidemiology and Drug Safety</i> , 2010 , 19, 579-85 | 2.6 | 65 |
| 58 | Use of multiple international healthcare databases for the detection of rare drug-associated outcomes: a pharmacoepidemiological programme comparing rosuvastatin with other marketed statins. <i>Pharmacoepidemiology and Drug Safety</i> , 2010 , 19, 1218-24 | 2.6 | 13 |
| 57 | Rheumatoid arthritis in UK primary care: incidence and prior morbidity. <i>Scandinavian Journal of Rheumatology</i> , 2009 , 38, 173-7 | 1.9 | 85 |

| | | | |
|----|---|------|-----|
| 56 | Chronic obstructive pulmonary disease in UK primary care: incidence and risk factors. <i>COPD: Journal of Chronic Obstructive Pulmonary Disease</i> , 2009 , 6, 369-79 | 2 | 30 |
| 55 | Cardiovascular risks of nonsteroidal antiinflammatory drugs in patients after hospitalization for serious coronary heart disease. <i>Circulation: Cardiovascular Quality and Outcomes</i> , 2009 , 2, 155-63 | 5.8 | 100 |
| 54 | Positive predictive value of computerized medical records for uncomplicated and complicated upper gastrointestinal ulcer. <i>Pharmacoepidemiology and Drug Safety</i> , 2009 , 18, 900-9 | 2.6 | 33 |
| 53 | The use of selective cyclooxygenase-2 inhibitors and the risk of acute myocardial infarction in Saskatchewan, Canada. <i>Pharmacoepidemiology and Drug Safety</i> , 2009 , 18, 1016-25 | 2.6 | 26 |
| 52 | Use of angiotensin II receptor blockers and the risk of acute pancreatitis: a nested case-control study. <i>Pancreatology</i> , 2009 , 9, 786-92 | 3.8 | 16 |
| 51 | Acid-suppressive drugs and community-acquired pneumonia. <i>Epidemiology</i> , 2009 , 20, 800-6 | 3.1 | 51 |
| 50 | Role of dose potency in the prediction of risk of myocardial infarction associated with nonsteroidal anti-inflammatory drugs in the general population. <i>Journal of the American College of Cardiology</i> , 2008 , 52, 1628-36 | 15.1 | 238 |
| 49 | Relationship between gastroesophageal reflux disease and COPD in UK primary care. <i>Chest</i> , 2008 , 134, 1223-1230 | 5.3 | 49 |
| 48 | Risk of myocardial infarction and overall mortality in survivors of venous thromboembolism. <i>Thrombosis Journal</i> , 2008 , 6, 10 | 5.6 | 8 |
| 47 | The safety of rosuvastatin in comparison with other statins in over 25,000 statin users in the Saskatchewan Health Databases. <i>Pharmacoepidemiology and Drug Safety</i> , 2008 , 17, 953-61 | 2.6 | 27 |
| 46 | The safety of rosuvastatin in comparison with other statins in over 100,000 statin users in UK primary care. <i>Pharmacoepidemiology and Drug Safety</i> , 2008 , 17, 943-52 | 2.6 | 53 |
| 45 | Effect of antisecretory drugs and nitrates on the risk of ulcer bleeding associated with nonsteroidal anti-inflammatory drugs, antiplatelet agents, and anticoagulants. <i>American Journal of Gastroenterology</i> , 2007 , 102, 507-15 | 0.7 | 230 |
| 44 | Use of acid-suppressing drugs and the risk of bacterial gastroenteritis. <i>Clinical Gastroenterology and Hepatology</i> , 2007 , 5, 1418-23 | 6.9 | 101 |
| 43 | Risk of upper gastrointestinal complications among users of traditional NSAIDs and COXIBs in the general population. <i>Gastroenterology</i> , 2007 , 132, 498-506 | 13.3 | 185 |
| 42 | Traditional nonsteroidal anti-inflammatory drugs and postmenopausal hormone therapy: a drug-drug interaction?. <i>PLoS Medicine</i> , 2007 , 4, e157 | 11.6 | 11 |
| 41 | NSAIDs and the risk of acute myocardial infarction. <i>Nature Clinical Practice Rheumatology</i> , 2007 , 3, 202-3 | | 0 |
| 40 | Use of oral corticosteroids and the risk of acute myocardial infarction. <i>Atherosclerosis</i> , 2007 , 192, 376-83 | 3.1 | 85 |
| 39 | Cardioprotective aspirin users and their excess risk of upper gastrointestinal complications. <i>BMC Medicine</i> , 2006 , 4, 22 | 11.4 | 130 |

| | | | |
|----|--|------|-----|
| 38 | The ever growing story of cyclo-oxygenase inhibition. <i>Lancet, The</i> , 2006 , 368, 1745-7 | 40 | 26 |
| 37 | Acute gastroenteritis is followed by an increased risk of inflammatory bowel disease. <i>Gastroenterology</i> , 2006 , 130, 1588-94 | 13.3 | 286 |
| 36 | Rosuvastatin safety: a comprehensive, international pharmacoepidemiology programme. <i>Pharmacoepidemiology and Drug Safety</i> , 2006 , 15, 454-61 | 2.6 | 5 |
| 35 | Upper gastrointestinal complications among users of paracetamol. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2006 , 98, 297-303 | 3.1 | 24 |
| 34 | Non-steroidal antiinflammatory drugs and the risk of acute myocardial infarction. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2006 , 98, 266-74 | 3.1 | 171 |
| 33 | Low-dose aspirin for the prevention of atherothrombosis. <i>New England Journal of Medicine</i> , 2005 , 353, 2373-83 | 59.2 | 870 |
| 32 | Predictors and prognosis of paroxysmal atrial fibrillation in general practice in the UK. <i>BMC Cardiovascular Disorders</i> , 2005 , 5, 20 | 2.3 | 72 |
| 31 | Long-term use of non-steroidal anti-inflammatory drugs and the risk of myocardial infarction in the general population. <i>BMC Medicine</i> , 2005 , 3, 17 | 11.4 | 49 |
| 30 | Risk of uncomplicated peptic ulcer among users of aspirin and nonaspirin nonsteroidal antiinflammatory drugs. <i>American Journal of Epidemiology</i> , 2004 , 159, 23-31 | 3.8 | 96 |
| 29 | Nonsteroidal antiinflammatory drugs and the risk of myocardial infarction in the general population. <i>Circulation</i> , 2004 , 109, 3000-6 | 16.7 | 167 |
| 28 | Association between captopril, other antihypertensive drugs and risk of prostate cancer. <i>Prostate</i> , 2004 , 58, 50-6 | 4.2 | 84 |
| 27 | Nonsteroidal antiinflammatory drugs as a trigger of clinical heart failure. <i>Epidemiology</i> , 2003 , 14, 240-6 | 3.1 | 112 |
| 26 | Validity of the general practice research database. <i>Pharmacotherapy</i> , 2003 , 23, 686-9 | 5.8 | 437 |
| 25 | Users of oral steroids are at a reduced risk of developing irritable bowel syndrome. <i>Pharmacoepidemiology and Drug Safety</i> , 2003 , 12, 583-8 | 2.6 | 9 |
| 24 | Risk of stroke associated with nonsteroidal anti-inflammatory drugs: a nested case-control study. <i>Stroke</i> , 2003 , 34, 379-86 | 6.7 | 18 |
| 23 | Risk of acute liver injury in patients with diabetes. <i>Pharmacotherapy</i> , 2002 , 22, 1091-6 | 5.8 | 14 |
| 22 | Risk of clinical blood dyscrasia in a cohort of antibiotic users. <i>Pharmacotherapy</i> , 2002 , 22, 630-6 | 5.8 | 20 |
| 21 | Risk of irritable bowel syndrome among asthma patients. <i>Pharmacoepidemiology and Drug Safety</i> , 2002 , 11, 31-5 | 2.6 | 36 |

| | | | |
|----|--|------|-----|
| 20 | Selective serotonin reuptake inhibitors and the risk of stroke: a population-based case-control study. <i>Stroke</i> , 2002 , 33, 1465-73 | 6.7 | 115 |
| 19 | Incidence of serious upper gastrointestinal bleeding/perforation in the general population: review of epidemiologic studies. <i>Journal of Clinical Epidemiology</i> , 2002 , 55, 157-63 | 5.7 | 110 |
| 18 | Incidence of chronic atrial fibrillation in general practice and its treatment pattern. <i>Journal of Clinical Epidemiology</i> , 2002 , 55, 358-63 | 5.7 | 158 |
| 17 | Are users of lipid-lowering drugs at increased risk of peripheral neuropathy?. <i>European Journal of Clinical Pharmacology</i> , 2001 , 56, 931-3 | 2.8 | 76 |
| 16 | Risk of upper gastrointestinal bleeding and perforation associated with low-dose aspirin as plain and enteric-coated formulations. <i>BMC Clinical Pharmacology</i> , 2001 , 1, 1 | | 121 |
| 15 | Association between aspirin and upper gastrointestinal complications: systematic review of epidemiologic studies. <i>British Journal of Clinical Pharmacology</i> , 2001 , 52, 563-71 | 3.8 | 199 |
| 14 | The risk of obstructive airways disease in a glaucoma population. <i>Pharmacoepidemiology and Drug Safety</i> , 2001 , 10, 157-63 | 2.6 | 11 |
| 13 | Relative risk of upper gastrointestinal complications among users of acetaminophen and nonsteroidal anti-inflammatory drugs. <i>Epidemiology</i> , 2001 , 12, 570-6 | 3.1 | 219 |
| 12 | Incidence of newly diagnosed heart failure in UK general practice. <i>European Journal of Heart Failure</i> , 2001 , 3, 225-31 | 12.3 | 63 |
| 11 | Lipid-lowering drugs and risk of myopathy: a population-based follow-up study. <i>Epidemiology</i> , 2001 , 12, 565-9 | 3.1 | 176 |
| 10 | Association between nonsteroidal anti-inflammatory drugs and upper gastrointestinal tract bleeding/perforation: an overview of epidemiologic studies published in the 1990s. <i>Archives of Internal Medicine</i> , 2000 , 160, 2093-9 | | 490 |
| 9 | Differential effects of aspirin and non-aspirin nonsteroidal antiinflammatory drugs in the primary prevention of myocardial infarction in postmenopausal women. <i>Epidemiology</i> , 2000 , 11, 382-7 | 3.1 | 153 |
| 8 | Secondary prevention of upper gastrointestinal bleeding associated with maintenance acid-suppressing treatment in patients with peptic ulcer bleed. <i>Epidemiology</i> , 1999 , 10, 228-32 | 3.1 | 35 |
| 7 | Association between selective serotonin reuptake inhibitors and upper gastrointestinal bleeding: population based case-control study. <i>BMJ: British Medical Journal</i> , 1999 , 319, 1106-9 | | 289 |
| 6 | Increased risk of irritable bowel syndrome after bacterial gastroenteritis: cohort study. <i>BMJ: British Medical Journal</i> , 1999 , 318, 565-6 | | 278 |
| 5 | Health resource utilization and drug treatment pattern in a cohort of patients with a first episode of gastroesophageal reflux disease. <i>Pharmacoepidemiology and Drug Safety</i> , 1999 , 8, 493-500 | 2.6 | 5 |
| 4 | Use of the UK General Practice Research Database for pharmacoepidemiology. <i>British Journal of Clinical Pharmacology</i> , 1998 , 45, 419-25 | 3.8 | 431 |
| 3 | Risk of hospitalization for upper gastrointestinal tract bleeding associated with ketorolac, other nonsteroidal anti-inflammatory drugs, calcium antagonists, and other antihypertensive drugs. <i>Archives of Internal Medicine</i> , 1998 , 158, 33-9 | | 231 |

| | | |
|---|---|----|
| 2 | The UK General Practice Research Database375-385 | 18 |
| 1 | Cardiovascular Toxicities of NSAIDS: Epidemiologic Aspects287-312 | |