José A Pascual

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7581902/publications.pdf

Version: 2024-02-01

66 papers

1,772 citations

218381 26 h-index 301761 39 g-index

68 all docs 68 docs citations

68 times ranked 1657 citing authors

#	Article	IF	CITATIONS
1	Cigarettes vs. e-cigarettes: Passive exposure at home measured by means of airborne marker and biomarkers. Environmental Research, 2014, 135, 76-80.	3.7	159
2	Impact of the Spanish Smoking Law on Exposure to Second-Hand Smoke and Respiratory Health in Hospitality Workers: A Cohort Study. PLoS ONE, 2009, 4, e4244.	1.1	102
3	13C/12C Isotope ratio MS analysis of testosterone, in chemicals and pharmaceutical preparations. Journal of Pharmaceutical and Biomedical Analysis, 2001, 24, 645-650.	1.4	83
4	Structural analysis of the glycosylation of gene-activated erythropoietin (epoetin delta, Dynepo). Analytical Biochemistry, 2008, 383, 243-254.	1.1	78
5	Fast screening of anabolic steroids and other banned doping substances in human urine by gas chromatography/tandem mass spectrometry. Journal of Mass Spectrometry, 2002, 37, 1059-1073.	0.7	71
6	Recombinant Erythropoietin and Analogues. Therapeutic Drug Monitoring, 2004, 26, 175-179.	1.0	57
7	Fast screening method for diuretics, probenecid and other compounds of doping interest. Journal of Chromatography A, 1993, 655, 233-242.	1.8	50
8	Evaluation of protein <i>N</i> â€glycosylation in 2â€DE: Erythropoietin as a study case. Proteomics, 2007, 7, 4278-4291.	1.3	49
9	Assessment of chronic exposure to cigarette smoke and its change during pregnancy by segmental analysis of maternal hair nicotine. Journal of Exposure Science and Environmental Epidemiology, 2003, 13, 144-151.	1.8	47
10	Procedures for monitoring recombinant erythropoietin and analogues in doping control. Analytical and Bioanalytical Chemistry, 2007, 388, 1521-1529.	1.9	44
11	Assessment of exposure to secondhand smoke by questionnaire and salivary cotinine in the general population of Barcelona, Spain (2004–2005). Preventive Medicine, 2009, 48, 218-223.	1.6	44
12	Impact of the Spanish Smoke-Free Legislation on Adult, Non-Smoker Exposure to Secondhand Smoke: Cross-Sectional Surveys before (2004) and after (2012) Legislation. PLoS ONE, 2014, 9, e89430.	1.1	41
13	Prediction of LC retention of steroids using solvatochromic parameters. Chromatographia, 1995, 41, 573-580.	0.7	40
14	Direct Determination of Anabolic Steroids in Human Urine by On-line Solid-phase Extraction/Liquid Chromatography/Mass Spectrometry., 1996, 31, 309-319.		40
15	Quantitative Determination of Tricyclic Antidepressants and Their Metabolites in Plasma by Solid-Phase Extraction (Bond-Elut TCA) and Separation by Capillary Gas Chromatography With Nitrogen-Phosphorous Detection. Therapeutic Drug Monitoring, 1998, 20, 340-346.	1.0	38
16	Salivary cotinine concentrations in daily smokers in Barcelona, Spain: a cross-sectional study. BMC Public Health, 2009, 9, 320.	1.2	36
17	A simple and reliable method for the determination of nicotine and cotinine in teeth by gas chromatography/mass spectrometry. Rapid Communications in Mass Spectrometry, 2003, 17, 2853-2855.	0.7	35
18	Secondhand smoke exposure at home: Assessment by biomarkers and airborne markers. Environmental Research, 2014, 133, 111-116.	3.7	32

#	Article	IF	Citations
19	Rapid and sensitive determination of zidovudine and zidovudine glucuronide in human plasma by ion-pair high-performance liquid chromatography. Journal of Chromatography A, 1996, 721, 127-137.	1.8	30
20	Evaluation of immunoassays for the measurement of insulin-like growth factor-I and procollagen type III peptide, indirect biomarkers of recombinant human growth hormone misuse in sport. Clinical Chemistry and Laboratory Medicine, 2005, 43, 75-85.	1.4	30
21	Assessing the instability of the isoelectric focusing patterns of erythropoietin in urine. Electrophoresis, 2006, 27, 4387-4395.	1.3	30
22	Neurotoxic Thioether Adducts of 3,4-Methylenedioxymethamphetamine Identified in Human Urine After Ecstasy Ingestion. Drug Metabolism and Disposition, 2009, 37, 1448-1455.	1.7	30
23	Nicotine Concentrations in Deciduous Teeth and Cumulative Exposure to Tobacco Smoke During Childhood. JAMA - Journal of the American Medical Association, 2003, 290, 196-197.	3.8	28
24	Recombinant erythropoietin found in seized blood bags from sportsmen. Haematologica, 2008, 93, 313-314.	1.7	28
25	High-throughput wide dynamic range procedure for the simultaneous quantification of nicotine and cotinine in multiple biological matrices using hydrophilic interaction liquid chromatography-tandem mass spectrometry. Analytical and Bioanalytical Chemistry, 2015, 407, 8463-8473.	1.9	28
26	Assessment of tobacco specific nitrosamines (TSNAs) in oral fluid as biomarkers of cancer risk: A population-based study. Environmental Research, 2016, 151, 635-641.	3.7	27
27	Evaluation of immunoassays for the measurement of erythropoietin (EPO) as an indirect biomarker of recombinant human EPO misuse in sport. Journal of Pharmaceutical and Biomedical Analysis, 2004, 35, 1169-1177.	1.4	26
28	Liquid chromatography clean-up method to improve identification of anabolic agents in human urine by gas chromatography–mass spectrometry. Analytica Chimica Acta, 2004, 522, 79-88.	2.6	25
29	Growth Hormone in Sport: Beyond Beijing 2008. Therapeutic Drug Monitoring, 2009, 31, 3-13.	1.0	23
30	Impact of the Spanish smoking law in smoker hospitality workers. Nicotine and Tobacco Research, 2009, 11, 1099-1106.	1.4	22
31	Fully automated analytical method for codeine quantification in human plasma using on-line solid-phase extraction and high-performance liquid chromatography with ultraviolet detection. Biomedical Applications, 1999, 724, 295-302.	1.7	19
32	Monitoring Gene Therapy by External Imaging of mRNA: Pilot Study on Murine Erythropoietin. Therapeutic Drug Monitoring, 2007, 29, 612-618.	1.0	19
33	Development of a precolumn capillary liquid chromatography switching system for coupling to mass spectrometry. Journal of Separation Science, 1996, 8, 383-387.	1.0	18
34	Effect of Physical Fitness and Endurance Exercise on Indirect Biomarkers of Recombinant Erythropoietin Misuse. International Journal of Sports Medicine, 2007, 28, 9-15.	0.8	18
35	Evaluation of tobacco specific nitrosamines exposure by quantification of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in human hair of non-smokers. Scientific Reports, 2016, 6, 25043.	1.6	17
36	Evaluation of immunoassays for the measurement of soluble transferrin receptor as an indirect biomarker of recombinant human erythropoietin misuse in sport. Journal of Immunological Methods, 2004, 295, 89-99.	0.6	16

#	Article	IF	CITATIONS
37	Effect of Physical Fitness and Endurance Exercise on Indirect Biomarkers of Recombinant Growth Hormone Misuse: Insulin-Like Growth Factor I and Procollagen Type III Peptide. International Journal of Sports Medicine, 2006, 27, 976-983.	0.8	16
38	Stages of change, smoking characteristics, and cotinine concentrations in smokers: Setting priorities for smoking cessation. Preventive Medicine, 2011, 52, 139-145.	1.6	16
39	Secondhand exposure to aerosol from electronic cigarettes: pilot study of assessment of tobacco-specific nitrosamine (NNAL) in urine. Gaceta Sanitaria, 2019, 33, 575-578.	0.6	16
40	Immunoassays for the measurement of IGF-II, IGFBP-2 and -3, and ICTP as indirect biomarkers of recombinant human growth hormone misuse in sport. Journal of Pharmaceutical and Biomedical Analysis, 2008, 48, 844-852.	1.4	15
41	New urinary EPO drug testing method using two-dimensional gel electrophoresis. Clinica Chimica Acta, 2006, 373, 186-187.	0.5	14
42	Evaluation of immunoassays for the measurement of insulin and C-peptide as indirect biomarkers of insulin misuse in sport: Values in selected population of athletes. Journal of Pharmaceutical and Biomedical Analysis, 2009, 49, 793-799.	1.4	14
43	Passive smoking in babies: The BIBE study (Brief Intervention in babies. Effectiveness). BMC Public Health, 2010, 10, 772.	1.2	14
44	Nicotine depedence and salivary cotinine concentration in daily smokers. European Journal of Cancer Prevention, 2012, 21, 96-102.	0.6	14
45	Manufactured and roll-your-own cigarettes: A changing pattern of smoking in Barcelona, Spain. Environmental Research, 2017, 155, 167-174.	3.7	13
46	Effect of physical fitness and endurance exercise on indirect biomarkers of growth hormone and insulin misuse: Immunoassay-based measurement in urine samples. Journal of Pharmaceutical and Biomedical Analysis, 2010, 53, 1003-1010.	1.4	12
47	Third-hand exposure at homes: Assessment using salivary cotinine. Environmental Research, 2021, 196, 110393.	3.7	12
48	Prediction of LC retention of steroids using solvatochromic parameters. Chromatographia, 1995, 41, 573-580.	0.7	11
49	Intermittent hypoxia exposure in a hypobaric chamber and erythropoietin abuse interpretation. Journal of Sports Sciences, 2007, 25, 1241-1250.	1.0	11
50	Purification of erythropoietin from human plasma samples using an immunoaffinity well plate. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2010, 878, 2117-2122.	1.2	11
51	Association Between Time to First Cigarette After Waking Up and Salivary Cotinine Concentration. Nicotine and Tobacco Research, 2011, 13, 168-172.	1.4	10
52	Impact of a multi-level intervention to prevent secondhand smoke exposure in schoolchildren: A randomized cluster community trial. Preventive Medicine, 2013, 57, 585-590.	1.6	10
53	Assessment of salivary cotinine concentration among general non-smokers population: Before and after Spanish smoking legislations. Cancer Epidemiology, 2017, 51, 87-91.	0.8	10
54	Anti-EPO and anti-NESP antibodies raised against synthetic peptides that reproduce the minimal amino acid sequence differences between EPO and NESP. Analytical and Bioanalytical Chemistry, 2007, 388, 1531-1538.	1.9	9

#	Article	IF	CITATIONS
55	Current strategic approaches for the detection of blood doping practices. Forensic Science International, 2011, 213, 42-48.	1.3	9
56	Comparison of TSNAs concentration in saliva according to type of tobacco smoked. Environmental Research, 2019, 172, 73-80.	3.7	9
57	Clarification on the detection of epoetin delta and epoetin omega using isoelectric focusing. American Journal of Hematology, 2008, 83, 754-754.	2.0	8
58	New screening protocol for recombinant human erythropoietins based on differential elution after immunoaffinity purification. Journal of Pharmaceutical and Biomedical Analysis, 2010, 51, 255-259.	1.4	8
59	Environmental and individual exposure to secondhand aerosol of electronic cigarettes in confined spaces: Results from the TackSHS Project â€. Indoor Air, 2021, 31, 1601-1613.	2.0	4
60	Validity of self-reported intensity of exposure to second-hand smoke at home against environmental and personal markers. Gaceta Sanitaria, 2018, 32, 393-395.	0.6	3
61	Impact of the Spanish smoking legislation among smokers: A longitudinal study with biomarkers in Barcelona (Spain). Addictive Behaviors, 2018, 87, 101-108.	1.7	3
62	If you play with fire, you may get burned. Drug Testing and Analysis, 2020, 12, 582-587.	1.6	3
63	Toward a correct measure of third-hand exposure. Environmental Research, 2021, 194, 110686.	3.7	1
64	Association between biomarkers of tobacco consumption and lung cancer risk among daily smokers. Expert Review of Respiratory Medicine, 2022, 16, 247-252.	1.0	1
65	Research Spotlight: Bioanalysis and Analytical Services Research Group at The Municipal Institute for Medical Research IMIM-Hospital del Mar, Spain. Bioanalysis, 2009, 1, 1403-1409.	0.6	0
66	Exposición al humo ambiental del tabaco en el hogar y el tiempo libre según el dÃa de la semana (laborable y no laborable) en Barcelona. Revista De Psicologia De La Salud, 2012, 24, 173.	0.2	0