

Seong-Hoon Park

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7585133/seong-hoon-park-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

22
papers

156
citations

7
h-index

11
g-index

22
ext. papers

194
ext. citations

3.3
avg, IF

2.98
L-index

#	Paper	IF	Citations
22	Prevalence and Factors of Osteoporosis and High Risk of Osteoporotic Fracture in Patients with Ankylosing Spondylitis: A Multicenter Comparative Study of Bone Mineral Density and the Fracture Risk Assessment Tool. <i>Journal of Clinical Medicine</i> , 2022 , 11, 2830	5.1	0
21	Pathophysiology and MRI Findings of Infectious Spondylitis and the Differential Diagnosis. <i>Journal of the Korean Society of Radiology</i> , 2021 , 82, 1413	0.2	
20	Differentiation of Vertebral Metastases From Focal Hematopoietic Marrow Depositions on MRI: Added Value of Proton Density Fat Fraction. <i>American Journal of Roentgenology</i> , 2021 , 216, 734-741	5.4	2
19	Validity of the EOS-determined pelvic parameters and orientation with pelvic positional variation: a phantom study. <i>Scientific Reports</i> , 2021 , 11, 10468	4.9	1
18	Use of proton density fat fraction MRI to predict the radiographic progression of osteoporotic vertebral compression fracture. <i>European Radiology</i> , 2021 , 31, 3582-3589	8	5
17	Salt-and-Pepper Noise Sign on Fat-Fraction Maps by Chemical-Shift-Encoded MRI: A Useful Sign to Differentiate Bone Islands From Osteoblastic Metastases-A Preliminary Study. <i>American Journal of Roentgenology</i> , 2020 , 214, 1139-1145	5.4	2
16	Multivariate analyses of MRI findings for predicting osteomyelitis of the foot in diabetic patients. <i>Acta Radiologica</i> , 2020 , 61, 1205-1212	2	8
15	Comparison of proton density fat fraction, simultaneous R2*, and apparent diffusion coefficient for assessment of focal vertebral bone marrow lesions. <i>Clinical Radiology</i> , 2020 , 75, 123-130	2.9	4
14	Prevalence of osteoporosis in patients with systemic lupus erythematosus: A multicenter comparative study of the World Health Organization and fracture risk assessment tool criteria. <i>Osteoporosis and Sarcopenia</i> , 2020 , 6, 173-178	2.3	2
13	Absence of ATFL remnant does not affect the clinical outcomes of the modified broström operation for chronic ankle instability. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2020 , 28, 213-220	5.5	6
12	Clinical characteristics and role of whole-body bone scan in multifocal osteonecrosis. <i>BMC Musculoskeletal Disorders</i> , 2019 , 20, 23	2.8	5
11	Craniovertebral Junction Abnormalities in Surgical Patients With Congenital Muscular Torticollis. <i>Journal of Craniofacial Surgery</i> , 2018 , 29, e327-e331	1.2	3
10	Initial experience with synthetic MRI of the knee at 3T: comparison with conventional T weighted imaging and T mapping. <i>British Journal of Radiology</i> , 2018 , 91, 20180006	3.4	
9	Correlations between intravoxel incoherent motion diffusion-weighted MR imaging parameters and F-FDG PET/CT metabolic parameters in patients with vertebral bone metastases: initial experience. <i>British Journal of Radiology</i> , 2018 , 91, 20170889	3.4	7
8	Combination of whole-spine lateral radiograph and lateral scanogram in the assessment of global sagittal balance. <i>Spine Journal</i> , 2018 , 18, 255-260	4	1
7	Torticollis Caused by Nontraumatic Craniovertebral Junction Abnormalities. <i>Journal of Craniofacial Surgery</i> , 2018 , 29, 1266-1270	1.2	4
6	Intravoxel incoherent motion diffusion-weighted magnetic resonance imaging of focal vertebral bone marrow lesions: initial experience of the differentiation of nodular hyperplastic hematopoietic bone marrow from malignant lesions. <i>Skeletal Radiology</i> , 2017 , 46, 675-683	2.7	25

5	Initial experience with synthetic MRI of the knee at 3T: comparison with conventional T ₂ -weighted imaging and T ₂ mapping. <i>British Journal of Radiology</i> , 2017 , 90, 20170350	3.4	12
4	Supine Versus Standing Radiographs for Detecting Ischiofemoral Impingement: A Propensity Score-Matched Analysis. <i>American Journal of Roentgenology</i> , 2016 , 206, 1253-63	5.4	11
3	Correlation between clinical symptoms and magnetic resonance imaging findings in patients with temporomandibular joint internal derangement. <i>Journal of the Korean Association of Oral and Maxillofacial Surgeons</i> , 2015 , 41, 125-32	1.6	17
2	A superficial hyperechoic band in human articular cartilage on ultrasonography with histological correlation: preliminary observations. <i>Ultrasonography</i> , 2015 , 34, 115-24	4.3	3
1	An in vitro comparative study of T2 and T2* mappings of human articular cartilage at 3-Tesla MRI using histology as the standard of reference. <i>Skeletal Radiology</i> , 2014 , 43, 947-54	2.7	38