

Cristobal C Richart

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7589363/cristobal-c-richart-publications-by-citations.pdf>
Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

| | | | |
|--------------------|-------------------------|----------------|-----------------|
| 94 papers | 4,200 citations | 33 h-index | 62 g-index |
| 100 ext. papers | 4,652 ext. citations | 5.8 avg, IF | 4.35 L-index |

| # | Paper | IF | Citations |
|----|---|------|-----------|
| 94 | Circulating interleukin 6 levels, blood pressure, and insulin sensitivity in apparently healthy men and women. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 1154-9 | 5.6 | 418 |
| 93 | Resistin, adiponectin, ghrelin, leptin, and proinflammatory cytokines: relationships in obesity. <i>Obesity</i> , 2004 , 12, 962-71 | | 362 |
| 92 | Mirror extreme BMI phenotypes associated with gene dosage at the chromosome 16p11.2 locus. <i>Nature</i> , 2011 , 478, 97-102 | 50.4 | 322 |
| 91 | Substitution of nevirapine, efavirenz, or abacavir for protease inhibitors in patients with human immunodeficiency virus infection. <i>New England Journal of Medicine</i> , 2003 , 349, 1036-46 | 59.2 | 268 |
| 90 | Genetic polymorphism of alcohol dehydrogenase in europeans: the ADH2*2 allele decreases the risk for alcoholism and is associated with ADH3*1. <i>Hepatology</i> , 2000 , 31, 984-9 | 11.2 | 201 |
| 89 | The interleukin-6 (-174) G/C promoter polymorphism is associated with type-2 diabetes mellitus in Native Americans and Caucasians. <i>Human Genetics</i> , 2003 , 112, 409-13 | 6.3 | 145 |
| 88 | Circulating retinol-binding protein-4, insulin sensitivity, insulin secretion, and insulin disposition index in obese and nonobese subjects. <i>Diabetes Care</i> , 2007 , 30, 1802-6 | 14.6 | 118 |
| 87 | New adipokines vaspin and omentin. Circulating levels and gene expression in adipose tissue from morbidly obese women. <i>BMC Medical Genetics</i> , 2011 , 12, 60 | 2.1 | 113 |
| 86 | A polymorphism in the promoter of the tumor necrosis factor-alpha gene (-308) is associated with coronary heart disease in type 2 diabetic patients. <i>Atherosclerosis</i> , 2003 , 167, 257-64 | 3.1 | 93 |
| 85 | FABP 4 is associated with inflammatory markers and metabolic syndrome in morbidly obese women. <i>European Journal of Endocrinology</i> , 2011 , 164, 539-47 | 6.5 | 88 |
| 84 | Long-term changes in leptin, chemerin and ghrelin levels following different bariatric surgery procedures: Roux-en-Y gastric bypass and sleeve gastrectomy. <i>Obesity Surgery</i> , 2013 , 23, 1790-8 | 3.7 | 81 |
| 83 | CD14 monocyte receptor, involved in the inflammatory cascade, and insulin sensitivity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2003 , 88, 1780-4 | 5.6 | 79 |
| 82 | IgA nephropathy and polycystic kidney disease. <i>American Journal of Nephrology</i> , 1988 , 8, 477-8 | 4.6 | 67 |
| 81 | Genetic polymorphisms of ADH2, ADH3, CYP4502E1 Dra-I and Pst-I, and ALDH2 in Spanish men: lack of association with alcoholism and alcoholic liver disease. <i>Journal of Hepatology</i> , 2004 , 41, 744-50 | 13.4 | 66 |
| 80 | Upregulation of lipocalin 2 in adipose tissues of severely obese women: positive relationship with proinflammatory cytokines. <i>Obesity</i> , 2011 , 19, 2295-300 | 8 | 64 |
| 79 | Shedding of TNF-alpha receptors, blood pressure, and insulin sensitivity in type 2 diabetes mellitus. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2002 , 282, E952-9 | 6 | 57 |
| 78 | Ambulatory blood pressure and left ventricular mass in normotensive patients with autosomal dominant polycystic kidney disease. <i>Journal of the American Society of Nephrology: JASN</i> , 1999 , 10, 1020-6 | 12.7 | 52 |

| | | | |
|----|---|-----|----|
| 77 | Severe neutropenia and thrombocytopenia associated with infliximab. <i>Annals of Internal Medicine</i> , 2003 , 139, W-W63 | 8 | 50 |
| 76 | Plasma visfatin levels and gene expression in morbidly obese women with associated fatty liver disease. <i>Clinical Biochemistry</i> , 2013 , 46, 202-8 | 3.5 | 49 |
| 75 | Omega-3 docosahexaenoic acid and procyanidins inhibit cyclo-oxygenase activity and attenuate NF- κ B activation through a p105/p50 regulatory mechanism in macrophage inflammation. <i>Biochemical Journal</i> , 2012 , 441, 653-63 | 3.8 | 48 |
| 74 | Alcohol dehydrogenase of human and rat blood vessels. Role in ethanol metabolism. <i>FEBS Letters</i> , 1997 , 405, 26-30 | 3.8 | 47 |
| 73 | Mannose-binding lectin gene polymorphisms are associated with gestational diabetes mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004 , 89, 5081-7 | 5.6 | 44 |
| 72 | Retinol binding protein-4 circulating levels were higher in nonalcoholic fatty liver disease vs. histologically normal liver from morbidly obese women. <i>Obesity</i> , 2013 , 21, 170-7 | 8 | 43 |
| 71 | Progressive multifocal leukoencephalopathy in chronic lymphocytic leukemia after treatment with fludarabine. <i>Leukemia and Lymphoma</i> , 2002 , 43, 433-6 | 1.9 | 42 |
| 70 | Increased levels and adipose tissue expression of visfatin in morbidly obese women: the relationship with pro-inflammatory cytokines. <i>Clinical Endocrinology</i> , 2012 , 77, 691-8 | 3.4 | 41 |
| 69 | Polymorphisms in the interleukin-6 receptor gene are associated with body mass index and with characteristics of the metabolic syndrome. <i>Clinical Endocrinology</i> , 2006 , 65, 88-91 | 3.4 | 41 |
| 68 | Monocyte chemoattractant protein-1 in obesity and type 2 diabetes. Insulin sensitivity study. <i>Obesity</i> , 2007 , 15, 664-72 | 8 | 40 |
| 67 | Plasma levels of the soluble fraction of tumor necrosis factor receptors 1 and 2 are independent determinants of plasma cholesterol and LDL-cholesterol concentrations in healthy subjects. <i>Atherosclerosis</i> , 1999 , 146, 321-7 | 3.1 | 40 |
| 66 | Paraoxonase Gln-Arg(192) and Leu-Met(55) gene polymorphisms and enzyme activity in a population with a low rate of coronary heart disease. <i>Clinical Biochemistry</i> , 2002 , 35, 197-203 | 3.5 | 39 |
| 65 | Altered fatty acid metabolism-related gene expression in liver from morbidly obese women with non-alcoholic fatty liver disease. <i>International Journal of Molecular Sciences</i> , 2014 , 15, 22173-87 | 6.3 | 37 |
| 64 | Association of retinol-binding protein-4 (RBP4) with lipid parameters in obese women. <i>Obesity Surgery</i> , 2010 , 20, 1258-64 | 3.7 | 35 |
| 63 | Spanish HIV-1-infected long-term nonprogressors of more than 15 years have an increased frequency of the CX3CR1 249I variant allele. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2005 , 40, 527-31 | 3.1 | 35 |
| 62 | Lipoprotein(a) and the significance of the association between platelet glycoprotein IIIa polymorphisms and the risk of premature myocardial infarction. <i>Atherosclerosis</i> , 1998 , 140, 155-9 | 3.1 | 33 |
| 61 | Left ventricular hypertrophy in hypertensive patients with autosomal dominant polycystic kidney disease: influence of blood pressure and humoral and neurohormonal factors. <i>American Journal of Nephrology</i> , 2000 , 20, 193-200 | 4.6 | 31 |
| 60 | miR33a/miR33b* and miR122 as Possible Contributors to Hepatic Lipid Metabolism in Obese Women with Nonalcoholic Fatty Liver Disease. <i>International Journal of Molecular Sciences</i> , 2016 , 17, | 6.3 | 30 |

| | | | |
|----|---|------|----|
| 59 | Circulating microbiota-derived metabolites: a "liquid biopsy?". <i>International Journal of Obesity</i> , 2020 , 44, 875-885 | 5.5 | 30 |
| 58 | Adipocytokine levels in women with anorexia nervosa. Relationship with weight restoration and disease duration. <i>International Journal of Eating Disorders</i> , 2013 , 46, 855-61 | 6.3 | 29 |
| 57 | Influence of chronic alcohol abuse and liver disease on hepatic aldehyde dehydrogenase activity. <i>Alcohol</i> , 1998 , 15, 3-8 | 2.7 | 29 |
| 56 | Polymorphisms of alcohol-metabolizing enzymes and the risk for alcoholism and alcoholic liver disease in Caucasian Spanish women. <i>Drug and Alcohol Dependence</i> , 2006 , 84, 195-200 | 4.9 | 29 |
| 55 | Proximal tubular kidney damage and tenofovir: a role for mitochondrial toxicity?. <i>Aids</i> , 2004 , 18, 1741-2 | 3.5 | 26 |
| 54 | G protein beta3 gene variant, vascular function, and insulin sensitivity in type 2 diabetes. <i>Hypertension</i> , 2003 , 41, 124-9 | 8.5 | 26 |
| 53 | Downregulation of lipogenesis and fatty acid oxidation in the subcutaneous adipose tissue of morbidly obese women. <i>Obesity</i> , 2014 , 22, 2032-8 | 8 | 25 |
| 52 | Anti-inflammatory profile of FTO gene expression in adipose tissues from morbidly obese women. <i>Cellular Physiology and Biochemistry</i> , 2010 , 26, 1041-50 | 3.9 | 25 |
| 51 | Fatal Listeria meningitis, endocarditis and pericarditis in a patient with haemochromatosis. <i>Scandinavian Journal of Infectious Diseases</i> , 1997 , 29, 308-9 | | 25 |
| 50 | Effect of genetic variants of CCR2 and CCL2 on the natural history of HIV-1 infection: CCL2-2518GG is overrepresented in a cohort of Spanish HIV-1-infected subjects. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2007 , 44, 132-8 | 3.1 | 24 |
| 49 | Redundant roles of the phosphatidate phosphatase family in triacylglycerol synthesis in human adipocytes. <i>Diabetologia</i> , 2016 , 59, 1985-94 | 10.3 | 22 |
| 48 | Leptin and adiponectin, but not IL18, are related with insulin resistance in treated HIV-1-infected patients with lipodystrophy. <i>Cytokine</i> , 2012 , 58, 253-60 | 4 | 22 |
| 47 | A study on the TNF-alpha system in Caucasian Spanish patients with alcoholic liver disease. <i>Drug and Alcohol Dependence</i> , 2008 , 92, 91-9 | 4.9 | 22 |
| 46 | Effect of ascorbic acid on gastrointestinal aluminium absorption. <i>Lancet, The</i> , 1991 , 338, 1467 | 4.0 | 22 |
| 45 | Endocannabinoid receptors gene expression in morbidly obese women with nonalcoholic fatty liver disease. <i>BioMed Research International</i> , 2014 , 2014, 502542 | 3 | 21 |
| 44 | Polymorphism of RANTES chemokine gene promoter is not associated with long-term nonprogressive HIV-1 infection of more than 16 years. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2006 , 41, 17-22 | 3.1 | 21 |
| 43 | Genetic heterogeneity within ulcerative colitis determined by an interleukin-1 receptor antagonist gene polymorphism and antineutrophil cytoplasmic antibodies. <i>European Journal of Gastroenterology and Hepatology</i> , 1999 , 11, 413-20 | 2.2 | 21 |
| 42 | Clinical and adipocytokine changes after bariatric surgery in morbidly obese women. <i>Obesity</i> , 2014 , 22, 188-94 | 8 | 20 |

| | | | |
|----|---|------|----|
| 41 | Human subcutaneous adipose tissue LPIN1 expression in obesity, type 2 diabetes mellitus, and human immunodeficiency virus--associated lipodystrophy syndrome. <i>Metabolism: Clinical and Experimental</i> , 2007 , 56, 1518-26 | 12.7 | 20 |
| 40 | Proteomic Profile of Unstable Atheroma Plaque: Increased Neutrophil Defensin 1, Clusterin, and Apolipoprotein E Levels in Carotid Secretome. <i>Journal of Proteome Research</i> , 2016 , 15, 933-44 | 5.6 | 19 |
| 39 | Lipodystrophy and insulin resistance in combination antiretroviral treated HIV-1-infected patients: implication of resistin. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2011 , 57, 16-23 | 3.1 | 19 |
| 38 | Minimal-change glomerulopathy and carcinoma. Report of two cases and review of the literature. <i>American Journal of Nephrology</i> , 1993 , 13, 69-72 | 4.6 | 19 |
| 37 | Interleukin-17A Gene Expression in Morbidly Obese Women. <i>International Journal of Molecular Sciences</i> , 2015 , 16, 17469-81 | 6.3 | 18 |
| 36 | Adipogenic/lipid, inflammatory, and mitochondrial parameters in subcutaneous adipose tissue of untreated HIV-1-infected long-term nonprogressors: significant alterations despite low viral burden. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2012 , 61, 131-7 | 3.1 | 18 |
| 35 | Effect of TNF-alpha genetic variants and CCR5 Delta 32 on the vulnerability to HIV-1 infection and disease progression in Caucasian Spaniards. <i>BMC Medical Genetics</i> , 2010 , 11, 63 | 2.1 | 18 |
| 34 | The IL-6 system in HIV-1-infection and in HAART-related fat redistribution syndromes. <i>Aids</i> , 2008 , 22, 893-6 | 3.5 | 18 |
| 33 | Platelet count and interleukin 6 gene polymorphism in healthy subjects. <i>BMC Medical Genetics</i> , 2001 , 2, 6 | 2.1 | 18 |
| 32 | Decrease of skin and bronchial sensitization following short-intensive scheduled immunotherapy in mite-allergic asthma. <i>Chest</i> , 1993 , 103, 183-7 | 5.3 | 18 |
| 31 | Adipo/cytokines in atherosclerotic secretomes: increased visfatin levels in unstable carotid plaque. <i>BMC Cardiovascular Disorders</i> , 2016 , 16, 149 | 2.3 | 17 |
| 30 | Systemic vasculitis resembling periarteritis nodosa in the lupus-like syndrome induced by hydralazine. <i>American Journal of Nephrology</i> , 1987 , 7, 71-3 | 4.6 | 17 |
| 29 | PNPLA3 Expression Is Related to Liver Steatosis in Morbidly Obese Women with Non-Alcoholic Fatty Liver Disease. <i>International Journal of Molecular Sciences</i> , 2016 , 17, | 6.3 | 17 |
| 28 | Lack of association of SDF-1 3SA variant allele with long-term nonprogressive HIV-1 infection is extended beyond 16 years. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2005 , 40, 276-9 | 3.1 | 16 |
| 27 | Recovery of minimal change nephrotic syndrome and acute renal failure in a patient with Renal cell carcinoma. <i>American Journal of Nephrology</i> , 1998 , 18, 433-5 | 4.6 | 14 |
| 26 | Hepcidin in morbidly obese women with non-alcoholic fatty liver disease. <i>PLoS ONE</i> , 2017 , 12, e0187065 | 3.7 | 13 |
| 25 | Low Circulating Levels of Neurotensin in Women with Nonalcoholic Fatty Liver Disease Associated with Severe Obesity. <i>Obesity</i> , 2018 , 26, 274-278 | 8 | 12 |
| 24 | Parallel downregulation of retinol-binding protein-4 and adiponectin expression in subcutaneous adipose tissue of non-morbidly obese subjects. <i>European Journal of Endocrinology</i> , 2009 , 161, 87-94 | 6.5 | 12 |

| | | | |
|----|---|-----|----|
| 23 | Atypical liver alcohol dehydrogenase in the Spanish population: its relation with the development of alcoholic liver disease. <i>Alcoholism: Clinical and Experimental Research</i> , 1993 , 17, 782-5 | 3.7 | 12 |
| 22 | PPARgamma Pro12Ala polymorphism in HIV-1-infected patients with HAART-related lipodystrophy. <i>Current HIV Research</i> , 2009 , 7, 533-40 | 1.3 | 12 |
| 21 | Downregulation of de Novo Fatty Acid Synthesis in Subcutaneous Adipose Tissue of Moderately Obese Women. <i>International Journal of Molecular Sciences</i> , 2015 , 16, 29911-22 | 6.3 | 11 |
| 20 | Relationship between IL-8 Circulating Levels and TLR2 Hepatic Expression in Women with Morbid Obesity and Nonalcoholic Steatohepatitis. <i>International Journal of Molecular Sciences</i> , 2020 , 21, | 6.3 | 9 |
| 19 | No relationship between TNF- β genetic variants and combination antiretroviral therapy-related lipodystrophy syndrome in HIV type 1-infected patients: a case-control study and a meta-analysis. <i>AIDS Research and Human Retroviruses</i> , 2011 , 27, 143-52 | 1.6 | 9 |
| 18 | Circulating soluble CD14 monocyte receptor is associated with increased alanine aminotransferase. <i>Clinical Chemistry</i> , 2004 , 50, 1456-8 | 5.5 | 9 |
| 17 | Targeted metabolomic approach in men with carotid plaque. <i>PLoS ONE</i> , 2018 , 13, e0200547 | 3.7 | 8 |
| 16 | Polymorphisms in the interleukin-10 gene promoter and the risk of alcoholism and alcoholic liver disease in Caucasian Spaniard men. <i>Alcohol</i> , 2010 , 44, 211-6 | 2.7 | 8 |
| 15 | LMNA messenger RNA expression in highly active antiretroviral therapy-treated HIV-positive patients. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2007 , 46, 384-9 | 3.1 | 8 |
| 14 | Increased Circulating Levels of Alpha-Ketoglutarate in Morbidly Obese Women with Non-Alcoholic Fatty Liver Disease. <i>PLoS ONE</i> , 2016 , 11, e0154601 | 3.7 | 7 |
| 13 | Evaluation of the pharmacogenetics of immune recovery in treated HIV-infected patients. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2014 , 10, 81-101 | 5.5 | 6 |
| 12 | Asthma, mite sensitization, and sleeping in bunks. <i>Annals of Allergy, Asthma and Immunology</i> , 1999 , 82, 531-3 | 3.2 | 6 |
| 11 | Lamivudine in the treatment of hepatitis B virus-related polyarteritis nodosa. <i>Journal of Clinical Rheumatology</i> , 2007 , 13, 298-9 | 1.1 | 5 |
| 10 | Echocardiographic evaluation in patients with autosomal dominant polycystic kidney disease and end-stage renal disease. <i>American Journal of Kidney Diseases</i> , 1999 , 34, 264-72 | 7.4 | 4 |
| 9 | Retinol Binding Protein-4 Circulating Levels Were Higher in Nonalcoholic Fatty Liver Disease Vs. Histologically Normal Liver From Morbidly Obese Women. <i>Obesity</i> , | 8 | 4 |
| 8 | Obesity and Critical Care. <i>Clinical Pulmonary Medicine</i> , 2009 , 16, 202-211 | 0.3 | 3 |
| 7 | Granulomatous hepatitis and fever of unknown origin: influence of AIDS epidemics. <i>Journal of Clinical Gastroenterology</i> , 1993 , 17, 89 | 3 | 3 |
| 6 | Absence of CXCR4 C-terminal polymorphisms in HIV-1-infected and uninfected Spaniards. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2006 , 42, 382-3 | 3.1 | 2 |

| | | | |
|---|--|-----|---|
| 5 | More on unusual forms of meningococcal disease: septic polyarthrititis. <i>American Journal of Medicine</i> , 1986 , 81, 561-2 | 2.4 | 2 |
| 4 | Hepatocyte Notch Signaling Deregulation Related to Lipid Metabolism in Women with Obesity and Nonalcoholic Fatty Liver. <i>Obesity</i> , 2020 , 28, 1487-1493 | 8 | 2 |
| 3 | The Potential Protective Role of RUNX1 in Nonalcoholic Fatty Liver Disease. <i>International Journal of Molecular Sciences</i> , 2021 , 22, | 6.3 | 1 |
| 2 | Clinical Characteristics and Outcome of Infective Endocarditis in Individuals of the General Population Managed at a Teaching Hospital Without Cardiac Surgery Facilities. Study of 120 Cases. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2006 , 59, 1131-1139 | 0.7 | 0 |
| 1 | Lipocalin, Resistin and Gut Microbiota-Derived Propionate Could Be Used to Predict Metabolic Bariatric Surgery Selected Outcomes. <i>Processes</i> , 2022 , 10, 143 | 2.9 | |